



Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions

(Posted on clinicaltrials.gov as NCT02728102)

BMT CTN PROTOCOL 1401 Version 4.0

Study Co-Chairs

David Avigan, MD¹, Nina Shah, MD², David Chung, MD³

Protocol Team

Devine, Steven⁴
Fay, Joseph⁵
Geller, Nancy⁶
Holstein, Sarah⁷
Lazarus, Hillard⁸

Logan, Brent⁹
Mendizabal, Adam¹⁰
Nelson, Courtney¹⁰
O'Brien, Kelly¹⁰
Pasquini, Marcelo⁹

Poland, Nancy¹¹
Rosenblatt, Jacalyn¹
Rapoport, Aaron¹²
Soiffer, Robert¹³
Waller, Edmund¹⁴

Manufacturing Team:

Choi, Chris⁷
Copland, Ian¹⁴
Hematti, Peiman¹⁵
O'Donnell, Lynn⁴

Reese, Jane⁸
Keever-Taylor, Carolyn¹⁶
Young, James³
Zurawski, Gerard⁵

Uhl, Lynne¹
Stroopinsky, Dina¹
Herman, Michele¹

Sponsored by the National Institutes of Health

National Heart, Lung, and Blood Institute and National Cancer Institute
FDA IND # 16655

¹ Beth Israel Deaconess Medical Center

² University of California- San Francisco

³ Memorial Sloan-Kettering Cancer Center

⁴ The Ohio State University

⁵ Baylor Institute for Immunology Research, Baylor Research Institute

⁶ National Heart, Lung, and Blood Institute

⁷ Roswell Park Cancer Institute

⁸ University Hospitals Case Medical Center, Case Western Reserve University

⁹ Center for International Blood and Marrow Transplant Research, Medical College of Wisconsin

¹⁰ The EMMES Corporation

¹¹ National Marrow Donor Program

¹² University of Maryland


¹³ Dana Farber Cancer Institute

¹⁴ Emory University

¹⁵ University of Wisconsin

¹⁶ Medical College of Wisconsin


Approval Signature (Protocol Chair/Officer)


Approval Signature (Protocol Chair/Officer)

Core Study Participants:

University Hospitals Case Medical Center,
Case Western Reserve University
Emory University
Fred Hutchinson Cancer Research Center
Memorial Sloan Kettering Cancer Center
Mount Sinai Medical Center
Ohio State University
Roswell Park Cancer Institute
University of Minnesota
University of Nebraska
University of California-San Francisco
Medical College of Wisconsin

Affiliate Study Participants:

Beth Israel Deaconess Medical Center
University of Maryland
University of Wisconsin
University of Texas, MD Anderson Cancer
Center
Thomas Jefferson University

PROTOCOL SYNOPSIS – BMT CTN 1401 PROTOCOL

Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell (DC)/Myeloma Fusions

Co- Principal Investigators: David Avigan, MD, Nina Shah, MD, David Chung, MD

Study Design: The study is designed as a Phase II, multicenter trial of vaccination with DC/myeloma fusions with granulocyte macrophage colony-stimulating factor (GM-CSF) adjuvant plus lenalidomide maintenance therapy versus maintenance therapy alone or with GM-CSF following autologous transplant as part of upfront treatment of multiple myeloma (MM).

Primary Objective: The primary objective of this randomized trial is to compare the proportion of patients alive and in complete response (defined as CR or sCR) at one year post transplant (or approximately 10 months post randomization) between patients receiving DC/myeloma vaccine/GM-CSF with lenalidomide maintenance therapy to those receiving lenalidomide maintenance therapy with or without GM-CSF.

Clinical Secondary Objectives: To compare DC/myeloma vaccine/GM-CSF with lenalidomide maintenance therapy to lenalidomide maintenance therapy with or without GM-CSF with respect to: myeloma response (sCR, CR, VGPR, PR, and SD), conversion of partial to complete response, disease progression, treatment-related mortality, progression-free survival, overall survival, toxicities according to the Common Terminology Criteria for Adverse Events (CTCAE) v4.0, incidence of infections, and measures of minimal residual disease at one year post transplant (or approximately 10 months from randomization). Secondary exploratory analyses will also be conducted to compare in a pairwise fashion the vaccine arm, lenalidomide/GM-CSF arm, and lenalidomide alone arm, to describe the proportion of patients with collection of tumor cells who reach randomization, compliance with vaccine, and to describe the reproducibility of the vaccine manufacturing based on the release criteria.

Immunologic Objectives: The primary immunologic endpoint is to compare the effect of DC/myeloma vaccine/GM-CSF with lenalidomide maintenance therapy to lenalidomide maintenance therapy alone or with GM-CSF on treatment-induced expansion of myeloma-specific

T cells, defined as a 2.4-fold increase from pre-therapy to peak post-treatment levels.

Secondary immunologic endpoints will compare each of the treatment arms for: a) the percent of patients achieving at least a 10-fold expansion of myeloma-specific T cells; b) expansion of myeloma antigen-specific T cells by tetramer analysis; c) quantification of T cell subsets and PD-1 expressing lymphocytes; d) quantification of NK cells and identification of activation and inhibitory ligands; e) assessment of NK-mediated killing of myeloma targets; f) assessment of humoral response against whole myeloma cells and myeloma-associated antigens by SEREX and by characterizing myeloma-specific plasmablast responses.

Eligibility:

Patients must be considered transplant eligible by treating physician at the time of study. Patients will be enrolled prior to tumor cell collection for vaccine manufacturing. Eligible patients are ≥ 18.00 and < 71.00 years of age with symptomatic multiple myeloma requiring treatment, without history of prior disease progression or prior HCT, Karnofsky performance score $\geq 70\%$, evidence of at least 20% plasma cells in a bone marrow differential within 60 days of enrollment, and have received ≤ 1 cycle of systemic anti-myeloma therapy. If patient receives anti-myeloma therapy treatment after bone marrow aspirate to assess eligibility and before bone marrow aspirate for tumor cell collection, a repeat bone marrow evaluation will be required to confirm $\geq 20\%$ plasma cells in the bone marrow aspirate differential prior to enrollment and tumor cell collection. Patients with prior auto or allo transplant are not eligible. Patients with active autoimmune diseases will be excluded.

Treatment Plan:

After meeting the initial eligibility criteria, patients will undergo harvesting of myeloma cells for assessment of immune response and for vaccine generation. Patients will have 30 mL of marrow harvested and cryopreserved after undergoing immunophenotypic analysis per the BMT CTN 1401 Manufacturing SOPs. Patients will then complete initial systemic anti-myeloma therapy and autologous hematopoietic stem cell mobilization and collection according to institutional practices.

Approximately 2 months post-transplant, patients will undergo disease response assessment and will be randomized 2:1:1 to vaccine/GM-CSF plus lenalidomide maintenance, lenalidomide alone, or lenalidomide/GM-CSF. Patients randomized to the vaccine plus lenalidomide arm will undergo leukapheresis for

dendritic cell generation for vaccine manufacturing. All patients will start lenalidomide maintenance Day 90-100 post transplant. Patients randomized to the vaccine arm will receive vaccine the first day of cycles 2, 3, and 4 of maintenance. Vaccine will be administered by subcutaneous injection with 100 ug GM-CSF given subcutaneously at the vaccine site on day of vaccination and daily for a total of 4 days. Patients randomized to lenalidomide/GM-CSF will receive 100 ug GM-CSF given subcutaneously daily for a total of 4 days starting the first day of cycles 2, 3 and 4 of maintenance. Lenalidomide maintenance will continue for 2 years for patients who continue free of progression.

Vaccine Production and Characterization:

Vaccine Production will consist of the following steps per the BMT CTN 1401 Manufacturing SOPs:

- 1) Bone marrow aspiration of 30 mL, isolation of mononuclear cells by ficoll centrifugation, quantification of CD138+ or CD38+ cells by immunocytochemistry, and cryopreservation.
- 2) Leukapheresis collection post transplant and after randomization.
- 3) Mononuclear cells isolated by ficoll density centrifugation. Adherent cells segregated following 1 hour culture and then cultured for 5-7 days with GM-CSF and IL-4. Cultures are re-fed with cytokines after 3 days. Maturation with TNF α for 48-72 hours. DCs are harvested and undergo characterization by immunocytochemistry for expression of DR, CD86, and CD83.
- 4) Tumor cells are thawed and re-characterized.
- 5) DC and tumor cells are pelleted and co-cultured in the presence of PEG at a ratio of 3:1.
- 6) Fusion cells are quantified by determining the percentage of dual expressing cells by immunocytochemistry.

Vaccine Release criteria:

- 1) At least 20% of tumor cell prep must express CD38 or CD138 by staining
- 2) Tumor cell prep must yield ≥ 10 million plasma cells¹
- 3) 50% of DC prep express CD86
- 4) Viability of DC prep > 50%
- 5) Fusion efficiency > 15%
- 6) Fusion viability > 50%

¹ Plasma cell count is determined by multiplying the total mononuclear cell yield by the percent of CD38 and CD138 staining reported on the Immunohistochemistry Report


7) Sterility, mycoplasma, and endotoxin assays are negative

Accrual Objective:	Target accrual is 132 patients randomized to vaccine/GM-CSF/lenalidomide (n=66), lenalidomide alone (n=33), lenalidomide/GM-CSF (n=33), with an estimated total enrollment of 188, assuming about 30% of patients are unable to proceed with post-transplant immunotherapy.
Accrual Period:	36 months
Study Duration:	Patients will be followed for approximately 3 years after maintenance initiation
Interim Analysis:	There will be no interim analysis for efficacy or futility.

STUDY SCHEMA

<p>Initial Inclusion Criteria</p> <ol style="list-style-type: none"> 1. Transplant Eligible 2. Meet criteria for symptomatic multiple myeloma (Appendix A) 3. Age $\geq 18.00 - < 71.00$ 4. Karnofsky Performance of $\geq 70\%$ 5. 20% plasma cells in the bone marrow ≤ 60 days prior to enrollment¹ 6. Received ≤ 1 cycles of systemic anti-myeloma therapy² 7. Creatinine clearance ≥ 40 mL/min (estimated or calculated) <p>¹ If patient receives anti-myeloma therapy treatment after bone marrow aspirate to assess eligibility and before bone marrow aspirate for tumor cell collection, a repeat bone marrow evaluation will be required to confirm $\geq 20\%$ plasma cells in the bone marrow aspirate differential prior to enrollment and tumor cell collection.</p> <p>² Anti-myeloma therapy is defined as systemic treatment intended to treat the underlying myeloma disease. Treatments intended to alleviate pain and other symptoms of disease and/or administration of ≤ 160mg of dexamethasone or equivalent alternative steroid dose within a 30 day period are not considered anti-myeloma therapy.</p>	<p>Initial Exclusion Criteria</p> <ol style="list-style-type: none"> 1. Prior Autologous or Allogeneic HCT 2. Non secretory multiple myeloma (defined as normal serum and urine immunofixation and normal serum free light chain assay). 3. Patients with a history of Plasma Cell Leukemia at any time prior to enrollment 4. Patients with prior disease progression at any time prior to enrollment 5. Patients seropositive for the human immunodeficiency virus (HIV). 6. Patients receiving treatment with Daratumumab or other CD38 Monoclonal Antibodies. 7. Patient receiving other investigational or anti-myeloma drugs within 14 days of enrollment. 8. Patients with active clinically significant autoimmune disease 9. Patients who have received mid-intensity melphalan (>50 mg IV) as part of prior therapy. 10. Additional exclusion criteria outlined in section 2.3.2
<p><u>Inclusion Criteria for Randomization</u></p> <ol style="list-style-type: none"> 1. No disease progression since initiation of systemic anti-myeloma therapy 2. Received autologous transplant with melphalan 200mg/m² with a minimum cell dose of 2×10^6 CD34+ cells/kg ≤ 12 months from enrollment onto BMT CTN 1401 Tumor cell preparation must (1) Express $\geq 20\%$ CD38 or CD138 by staining (2) yield ≥ 10 million plasma cells (<i>Plasma cell count is determined by multiplying the total mononuclear cell yield by the percent of CD38 and CD138 staining reported on the Immunohistochemistry Report</i>) (3) Have a negative microbiology assessment 3. No evidence of uncontrolled infection requiring systemic therapy 4. Platelet count $\geq 75,000/\text{mm}^3$ (without transfusion in previous 7 days). 5. Absolute neutrophil count (ANC) $\geq 1,500/\text{mm}^3$ without filgrastim administration within 7 days, or pegfilgrastim within 14 days of measurement. 6. Bilirubin $< 2 \times$ the upper limit of normal 7. ALT and AST $< 2.5 \times$ the upper limit of normal 8. Creatinine clearance of ≥ 40 mL/min. Patients with creatinine clearance ≥ 30 but ≤ 40 will be considered with review/approval from the protocol chairs or officer if the cause of renal insufficiency is associated with multiple myeloma 9. Patients must be willing to receive DVT prophylaxis 10. Patient must be willing to follow birth control and pregnancy testing practices outlined in the protocol 	<p><u>Primary Clinical Endpoint:</u></p> <p>Complete response (CR or sCR) at one year between patients receiving DC/myeloma vaccine/GM-CSF with lenalidomide maintenance therapy to those receiving lenalidomide maintenance therapy with or without GM-CSF.</p> <p><u>Primary Immunologic Endpoint:</u></p> <p>Treatment-induced expansion of myeloma-specific T cells as defined by the 2.4-fold increase from pre-therapy to peak post-treatment levels comparing the vaccine arm with patients receiving lenalidomide maintenance therapy with and without GM-CSF.</p> <p><u>Secondary Endpoints:</u></p> <p><u>Clinical:</u> Myeloma Response; Rate of conversion to CR; Disease Progression; Treatment-Related Mortality; Progression-Free Survival; Overall Survival; Toxicity and rates of infection; Minimal Residual Disease; CR and sCR rates will be compared between the vaccine arm and each of the no-vaccine arms; proportion of patients with tumor cell collection who reach randomization; compliance with vaccine; and reproducibility of the vaccine.</p> <p><u>Immunologic:</u> Secondary immunologic endpoint :a) the percent of patients achieving at least a 10 fold expansion of myeloma specific T-cells cells; b) expansion of myeloma antigen-specific T cells by tetramer analysis; c) quantification of T-cells subsets and PD-1 expressing lymphocytes; d) quantification of NK cells and identification of activation and inhibitory ligands; e) assessment of NK mediated killing of myeloma targets; f) assessment of humoral response against whole myeloma cells and myeloma associated antigens by SEREX and by characterizing myeloma-specific plasmablast responses.</p>

OUTLINE OF TREATMENT PLAN



Note: Len = Maintenance lenalidomide

- Accrual targets 188 patients to be enrolled with a target of 132 patients to be randomized, assuming about 30% of patients are unable to proceed with post-transplant immunotherapy.
 - Arm A: Maintenance lenalidomide + vaccine + GM-CSF (n=66)
 - Arm B: Maintenance lenalidomide + GM-CSF (n=33)
 - Arm C: Maintenance lenalidomide alone (n=33)
- Patients will be stratified according to disease status at time of randomization between CR/sCR and VGPR/PR/stable disease.

TABLE OF CONTENTS

1.	OVERVIEW	1-1
1.1.	Background	1-1
1.1.1.	Vaccination with DC/Myeloma fusion cells following autologous transplant	1-1
1.1.2.	Post-transplant lenalidomide maintenance	1-2
1.2.	Study Specific Information	1-3
1.3.	Rationale for Study	1-3
2.	STUDY DESIGN.....	2-1
2.1.	Study Overview	2-1
2.2.	Hypotheses and Specific Objectives	2-1
2.2.1.	Hypothesis	2-1
2.2.2.	Study Objectives	2-1
2.2.3.	Clinical Secondary Objectives	2-1
2.2.4.	Immunologic Secondary Objectives	2-2
2.3.	Patient Eligibility	2-2
2.3.1.	Initial Inclusion Criteria	2-2
2.3.2.	Initial Exclusion Criteria	2-3
2.3.3.	Patient Eligibility Criteria for Randomization	2-4
2.4.	Treatment Plan.....	2-6
2.4.1.	Treatment Prior to Enrollment	2-6
2.4.2.	Tumor Cell Collection.....	2-6
2.4.3.	Systemic Anti-myeloma Therapy	2-6
2.4.4.	Stem Cell Mobilization	2-6
2.4.5.	Autologous Stem Cell Transplantation	2-7
2.4.6.	Maintenance Therapy.....	2-7
2.4.7.	Vaccine Manufacturing and Administration.....	2-9
2.4.8.	GM-CSF Administration.....	2-11
2.5.	Supportive Care	2-12
2.5.1.	Post Autologous Transplantation(s).....	2-12
2.5.2.	Deep Vein Thrombosis (DVT) and Pulmonary Embolism (PE) Prophylaxis	2-12
2.6.	Risks and Toxicities	2-12
2.6.1.	Therapy Toxicities	2-12
2.6.2.	Lenalidomide.....	2-12
2.6.3.	DC/Fusion Vaccine - Manufacturing and Administration	2-13
2.6.4.	GM-CSF	2-14
2.7.	Study Drug Supply.....	2-14
2.7.1.	Lenalidomide.....	2-14
2.8.	Patient Withdrawal from Study and Off Study Criteria	2-18
3.	STUDY ENDPOINTS AND DEFINITIONS.....	3-1
3.1.	Definition of Disease Status.....	3-1
3.1.1.	Response Categories	3-1

3.1.2.	Progressive Disease (PD)	3-2
3.2.	Primary Endpoint	3-3
3.3.	Clinical Secondary Endpoints.....	3-3
3.3.1.	Response to treatment	3-3
3.3.2.	Myeloma Progression	3-4
3.3.3.	Minimal Residual Disease Assessment.....	3-4
3.3.4.	Treatment-related Mortality	3-4
3.3.5.	Incidence of Toxicities Grade ≥ 3 per CTCAE vesion 4.0	3-4
3.3.6.	Incidence of Infections per BMT CTN Technical MOP.....	3-4
3.3.7.	Progression-free survival	3-4
3.3.8.	Overall Survival	3-5
3.4.	Primary Immunologic Endpoints.....	3-5
3.4.1.	Myeloma-Reactive T-cells	3-5
3.5.	Secondary Immunologic Endpoints:	3-5
3.5.1.	Myeloma-Reactive T-cells	3-5
3.5.2.	Assessment of antigen specific reactivity:	3-5
3.5.3.	Quantification of T-cell subsets:	3-5
3.5.4.	Assessment of effect on NK cell populations:	3-6
3.5.5.	Assessment of humoral immune response:	3-6
4.	PATIENT REGISTRATION, ENROLLMENT, AND EVALUATION.....	4-1
4.1.	Enrollment Procedures.....	4-1
4.2.	Study Monitoring.....	4-1
4.2.1.	Follow - Up Schedule	4-1
4.3.	Patient Evaluations	4-3
4.3.1.	Pre- Enrollment Segment A Evaluations and Requirements	4-3
4.3.2.	Tumor Cell Collection Evaluations and Requirements at Enrollment.....	4-4
4.3.3.	Pre- Randomization Evaluations and Requirements.....	4-4
4.3.4.	Pre- Maintenance Evaluations and Requirements.....	4-5
4.3.5.	Evaluations during Maintenance.....	4-6
4.4.	Data Reporting	4-11
4.4.1.	Criteria for Forms Submission	4-11
4.4.2.	Reporting Patient Deaths.....	4-11
4.4.3.	Center for International Blood and Marrow Transplant Research (CIBMTR) Data Reporting.....	4-11
4.4.4.	Adverse Event Reporting	4-11
4.4.5.	Pregnancy Reporting.....	4-13
4.4.6.	Celgene Drug Safety Contact Information:.....	4-14
4.4.7.	Reporting of other Special Events of Interest	4-14
4.5.	Ancillary Correlative Laboratory Studies Procedures	4-14
4.5.1.	Obtaining Informed Consent.....	4-14
4.5.2.	Data Entry Requirements	4-15
5.	STATISTICAL CONSIDERATIONS	5-1

5.1.	Study Design	5-1
5.1.1.	Accrual	5-1
5.1.2.	Randomization	5-1
5.1.3.	Primary Endpoint	5-1
5.1.4.	Primary Hypothesis	5-1
5.1.5.	Duration of follow-up	5-2
5.2.	Sample Size and Power Considerations	5-2
5.2.1.	Primary Endpoint (Clinical)	5-2
5.2.2.	Power Considerations for Primary Immunologic Endpoint	5-2
5.3.	Interim Analysis and Stopping Guidelines	5-2
5.3.1.	Interim Analysis for Efficacy	5-2
5.3.2.	Interim Analysis for Futility	5-2
5.3.3.	Guidelines for Safety Monitoring	5-3
5.4.	Demographic and Baseline Characteristics	5-4
5.5.	Analysis of Primary Clinical Endpoint	5-4
5.6.	Analysis of Secondary Clinical Endpoints	5-4
5.6.1.	Response to treatment	5-4
5.6.2.	Myeloma Progression	5-4
5.6.3.	Treatment-related Mortality	5-4
5.6.4.	Progression-free survival	5-5
5.6.5.	Overall survival	5-5
5.6.6.	Incidence of toxicities grade ≥ 3 per CTCAE version 4.0	5-5
5.6.7.	Incidence of infections	5-5
5.6.8.	Minimal Residual Disease assessment	5-5
5.7.	Analysis of Immunological Endpoints	5-5
5.7.1.	Primary immunologic endpoint: Myeloma reactive T-cells	5-5
5.7.2.	Secondary immunologic endpoints:	5-6
5.8.	Endpoint Review Committee	5-6

LIST OF APPENDICES

APPENDIX A	CRITERIA FOR SYMPTOMATIC MULTIPLE MYELOMA
APPENDIX B1	INFORMED CONSENT FORM FOR BMT CTN 1401
APPENDIX B2	INFORMED CONSENT DOCUMENTS FOR BMT CTN 1401 OPTIONAL CORRELATIVE LABORATORY STUDY PARTICIPATION
APPENDIX C	LABORATORY PROCEDURES
APPENDIX D	LENALIDOMIDE RISKS OF FETAL EXPOSURE, PREGNANCY TESTING GUIDELINES AND ACCEPTABLE BIRTH CONTROL METHODS
APPENDIX E	Revlimid REMS® PROGRAM
APPENDIX F	HUMAN SUBJECTS
APPENDIX G	KARNOFSKY PERFORMANCE STATUS SCALE
APPENDIX H	NEW YORK HEART ASSOCIATION (NYHA) CLASSIFICATION OF CARDIAC DISEASE

APPENDIX I	ADVERSE EVENTS
APPENDIX J	VCAN ANCILLARY STUDY
APPENDIX K	T CELL/MYELOMA GENOMIC SIGNATURE ANCILLARY STUDY
APPENDIX L	REFERENCES

CHAPTER 1

1. OVERVIEW

1.1. Background

Immune therapy has emerged as a leading area of cancer therapeutics due to its potential to recruit multiple effectors that broadly target malignant cells and overcome mechanisms of resistance. Multiple myeloma (MM) is characterized by the loss of critical mediators of immune surveillance, resulting in the suppression of antigen-presenting and effector cell function and the development of an immunologic milieu that fosters disease progression¹⁻⁶. We have developed a tumor vaccine in which patient-derived myeloma cells are fused with autologous dendritic cells (DCs), such that a broad array of myeloma antigens are presented in the context of DC-mediated costimulation⁷⁻¹⁰. In diverse tumor models that include MM, vaccination of animals with DC/tumor fusions results in protection from an otherwise lethal challenge of malignant cells and, more significantly, eradication of disease in the setting of advanced metastatic involvement^{11,12}. In a phase I study, 17 patients with advanced myeloma (median of 4 prior regimens) underwent vaccination with DC/MM fusions in conjunction with 4 days of GM-CSF administered at the vaccine site. As previously described, MM cells were isolated from bone marrow aspirates and fused with autologous DCs generated from adherent mononuclear cells cultured with GM-CSF, IL-4, and TNF α ¹³. Vaccine production was successful in all patients and was well-tolerated without evidence of clinically significant autoimmunity. Vaccination resulted in a mean 10-fold expansion of CD4 and CD8 myeloma-specific T cells, as determined by the percentage of cells expressing IFN γ in response to ex vivo exposure to autologous tumor lysate. Similarly, vaccination resulted in the development of myeloma-specific antibody responses as documented by SERAX analysis¹³.

1.1.1. Vaccination with DC/Myeloma fusion cells following autologous transplant

Vaccination with DC/Myeloma fusion cells following autologous transplant is associated with potent immune response and conversion from PR to CR in a subset of patients¹⁴. We completed a clinical trial in which MM patients were vaccinated with DC/MM fusion cells in conjunction with autologous transplantation (ASCT). In the first cohort of the study, twenty-four patients received serial vaccinations following post-transplant hematopoietic recovery. A second cohort of 12 patients received a pre-transplant vaccine followed by post-transplant vaccinations. Vaccine preparation was successful in all patients. Mean yield of the DC and MM cells was 1.74×10^8 and 6.5×10^7 cells, respectively. Mean fusion efficiency, as determined by the percentage of cells that co-expressed unique DC (CD80, CD86, and/or CD83) and MM (CD38 and/or CD138) antigens, was 38%. The mean dose administered was 3.6×10^6 fusion cells. Mean viability of the DC, myeloma, and fusion preparations was 87%, 87%, 79%, respectively. The DC/MM fusion preparations exhibited potent antigen presenting capacity, as evidenced by their stimulation of allogeneic T cell proliferation, with a mean stimulation index of 36.6, similar to that observed with the DC preparation prior to fusion (mean stimulation index of 52.3). In contrast, the unfused myeloma cells demonstrated minimal capacity for T cell stimulation (mean stimulation index of 13.7). All vaccine-associated toxicities were of grade I-II intensity. The most common toxicity was erythema and induration at the vaccine injection site associated with T cell infiltration on biopsy. The other common side effects were transient pruritus, rash, fatigue, fever, and myalgias. One patient had a transiently elevated ANA level without clinical evidence of autoimmunity. One patient developed transient grade 2 leukopenia, but no evidence of graft compromise was observed following vaccination. Consistent with prior reports, the post-transplant period was associated with

the relative suppression of general measures of cellular immunity as manifested by decreased T cell proliferative responses to mitogen or recall antigens. In contrast, the period of post-transplant lymphopoietic reconstitution was associated with the expansion of myeloma-specific T cells that was further boosted following vaccination, as determined by percentage of CD4+ and CD8+ T cells expressing IFN γ in response to ex vivo exposure to autologous tumor lysate. In the first cohort that received only post-transplant vaccination, the mean log10 fold increase in myeloma-specific CD4+ T cells from pre-mobilization to post-transplant and from pre-mobilization to peak post-vaccination was 3.55 (95% CI 0.81;15.49) and 10.72 (95% CI 3.89;29.51), respectively. Similarly, the mean log10 fold increase in myeloma specific CD8+ T cells from pre-mobilization to post-transplant and from pre-mobilization to peak post-vaccination was 6.76 (95% CI 3.02;15.49) and 11.48 (95% CI 4.17;32.36), respectively. A smaller subset of patients underwent a single pre-transplant vaccination followed by post-transplant boosting. Of note, no difference in peak levels of CD4+ or CD8+ circulating myeloma-specific T cells was observed between the cohort receiving a pre-transplant vaccine and that undergoing post-transplant vaccination alone ($p=0.185$ and $p=0.689$, respectively). For the entire study population, the mean log10 fold increase in myeloma-specific CD4+ T cells from pre-mobilization to post-transplant and from pre-mobilization to peak post-vaccination was 3.55 (95% CI 1.58;8.13) and 9.55 (95% CI 5.37;16.98), respectively. The mean log10 fold increase in myeloma-specific CD8+ T cells from pre-mobilization to post-transplant and from pre-mobilization to peak post-vaccination was 4.37 (95% CI 2.40;7.76) and 8.32 (95% CI 4.68;15.14), respectively. Notably, vaccination was associated with the expansion of T cells targeting the myeloma-specific antigen, MUC1, as determined by tetramer analysis. In a cohort of patients who were HLA-A2, the median percentage of MUC1-specific T cells was 0.12 pre-transplant and increased to 1.84 at 3 months after completion of vaccination ($p<0.05$), representing a median fold increase of 17.5. The expansion of T cells targeting MUC1 in response to vaccination is of particular interest, as we have identified a population of CD34+/MUC1+ cells in the bone marrow of patients with myeloma that demonstrate myeloma-initiating capacity.

Seventy-eight percent of patients achieved a CR or VGPR (47% CR/nCR; 31% VGPR). Thirty-one percent achieved a CR/nCR in the early post-transplant period, whereas an additional 17% (6 patients: 4 from VGPR, 2 from PR) achieved CR/nCR as best response only after day 100 post-transplant and after undergoing vaccination. The presence of late responses several months after ASCT is consistent with an impact of vaccine therapy on post-transplant residual disease.

1.1.2. Post-transplant lenalidomide maintenance

Post-transplant lenalidomide maintenance prolongs disease-free and overall survival¹⁵. This is thought to be potentially due to its immunomodulatory characteristics. In pre-clinical studies, we demonstrated that lenalidomide potently enhances immune response to vaccination¹⁶. T cells were stimulated in vitro with DC/MM fusions in the absence or presence of lenalidomide. After 5–7 days of stimulation by the vaccine, the percentage of CD8+ T cells expressing IFN γ increased significantly in the presence of lenalidomide (from 6.37 to 9.9 %; $n = 7$, $p = 0.03$). A concomitant statistically significant decrease in the proportion of regulatory T cells was also observed (4.4 % in the lenalidomide group as opposed to 9.6 % in the control group; $n = 3$, $p = 0.02$). As determined by a fluorochrome assay measuring granzyme B release in target cells, CTL-mediated lysis of autologous MM tumor targets was increased by 50% when T cells were stimulated by DC/MM fusions in the presence of lenalidomide, as compared to those cultured with fusions alone (mean 29.2% vs. 19.7%; $n=3$, $p=0.004$).

1.2. Study Specific Information

Based on these encouraging data, the DC/MM fusion vaccine was chosen to be the first cellular immunotherapy to be evaluated in a multicenter cooperative group study involving several nationally prominent cancer centers (BMT CTN protocol 1401). We will conduct a multicenter phase II randomized study comparing post-transplant vaccination with DC/MM fusions/GM-CSF and lenalidomide maintenance to lenalidomide maintenance with and without GM-CSF, with vaccine manufacturing performed locally at each participating center. The primary objective is to assess the impact of post-transplant immunotherapy on the percentage of patients in CR at 1 year between the vaccine arm and both no-vaccine arms combined. Secondary endpoints include determining the effect of therapy on progression-free survival, myeloma response, minimal residual disease, toxicities, transplant-related mortality, and overall survival. Additionally, this study includes several immunologic endpoints to understand the effect of the vaccine on the anti-myeloma immunity, the immune effects of lenalidomide, and general immune reconstitution after transplant. The primary immunologic endpoint is to compare the expansion of myeloma-specific T cells across all three arms.

1.3. Rationale for Study

MM is associated with immune suppression, including defects in antigen presentation and effector cell function that are thought to play a role in disease progression. Although the outcomes for myeloma patients have improved greatly over the past decade, the majority of patients relapse even after autologous stem cell transplant and post-transplant maintenance therapy with lenalidomide. Early phase clinical data demonstrate that post-transplant vaccination with a DC-MM fusion product with GM-CSF adjuvant increases the number of MM-specific T cells. Furthermore, it appears that lenalidomide can specifically augment vaccine-stimulated induction of MM-specific T cells. We are therefore interested in combining standard immunomodulatory therapy (lenalidomide) with a potential immune stimulant (DC-MM vaccination) to increase post-transplant clinical response. Finally, pre-clinical data has suggested that immunomodulatory agents augment NK cell activity¹⁷ and that NK cells interact with and can be activated by DCs¹⁸. Thus, we are interested in further defining the NK immunophenotype in the setting of post-transplant maintenance lenalidomide and the changes to this phenotype in the setting of DC-MM fusion vaccination. We postulate that post-transplant lenalidomide maintenance will augment post-transplant anti-myeloma immunity, thus providing an ideal platform for DC/MM fusion vaccination. Additionally, to understand the effect of vaccine versus the adjuvant in the post-transplant setting, we split the non-vaccine arms into lenalidomide alone and lenalidomide plus GM-CSF, with GM-CSF administered in the exact same schedule and dosing as in the vaccine arm. Although there is no evidence that GM-CSF has an anti-myeloma effect, this additional control arm is important to understand whether any effect of the vaccine is due to its adjuvant and whether GM-CSF influences any of the immunologic outcomes. We further hypothesize that vaccination in this setting will effectively target post-transplant residual disease, increase the percentage of patients achieving CR, and create durable responses.

The results of this trial will provide information on whether this patient-specific vaccine administered after transplant is promising for further comparison with current standard of care treatment. Furthermore, this trial will address several important aspects of anti-myeloma immunity, including how lenalidomide maintenance influences post-transplant recovery and myeloma control. Lastly, isolating the effect of the GM-CSF adjuvant will address whether any effect observed is related solely to the vaccine.

CHAPTER 2

2. STUDY DESIGN

2.1. Study Overview

The study is a three-arm, phase II randomized, open-labeled clinical trial to evaluate vaccination with DC/myeloma fusions/GM-CSF plus lenalidomide maintenance therapy compared to lenalidomide maintenance therapy with or without GM-CSF following autologous transplant as part of upfront treatment for patients diagnosed with multiple myeloma. Patients meeting initial eligibility will be enrolled, and tumor cells will be harvested. Patients will proceed or continue with systemic anti-myeloma therapy and stem cell mobilization according to institutional practices. Patients with available cryopreserved tumor cells who are eligible to proceed to transplantation will be randomized in the study around Day 56 post-transplant.

Approximately 2 months post-transplant, patients will undergo their first disease response assessment and will be randomized 2:1:1 to the vaccine/GM-CSF plus lenalidomide arm, lenalidomide/GM-CSF arm, or lenalidomide alone arm. Patients assigned to vaccine plus lenalidomide maintenance will subsequently undergo leukapheresis for collection of dendritic cell precursors and vaccine production (DC/myeloma vaccine). For all patients, lenalidomide maintenance will start at Day 90-100 post-transplant and continue for two years or until disease progression, whichever comes first. Vaccines will be administered on the first day of cycles 2, 3, and 4 of lenalidomide maintenance, followed by four daily doses of GM-CSF for patients randomized to the vaccine arm. Patients randomized to lenalidomide/GM-CSF will receive GM-CSF in the same schedule and dose as patients in the vaccine arm. Primary endpoint assessment will occur at 1 year post-transplant.

2.2. Hypotheses and Specific Objectives

2.2.1. Hypothesis

Vaccination with DC/myeloma fusions plus lenalidomide maintenance therapy will result in improved responses in patients with multiple myeloma after autologous HCT.

2.2.2. Study Objectives

Compare the proportion of patients alive and in complete response (CR or sCR) at one year post-transplant, corresponding to 1 year post transplant (approximately 10 months after randomization) between patients receiving DC/myeloma vaccine/GM-CSF with lenalidomide maintenance therapy to those receiving lenalidomide maintenance therapy with or without GM-CSF.

2.2.3. Clinical Secondary Objectives

Compare myeloma response, rate of conversion to CR, myeloma progression, treatment-related mortality, progression-free survival, overall survival, incidence of grade ≥ 3 toxicity according to the Common Terminology Criteria for Adverse Events (CTCAE) v4.0, incidence of infections, measures of minimal residual disease, between the vaccine and the combined no-vaccine treatment arms. A secondary pairwise analysis of the proportion of patients alive and in CR at one year post-transplant will also be conducted between the vaccine arm, lenalidomide/GM-CSF arm, and the

lenalidomide alone arm. Additionally, the study will describe the proportion of patients with collection of tumor cells who reach randomization, compliance with vaccine, and the reproducibility of the vaccine manufacturing based on the release criteria.

2.2.4. Immunologic Secondary Objectives

Primary immunologic endpoints:

- Compare the expansion of treatment-induced myeloma-specific T cells, defined as a 2.4-fold increase from pre-therapy to peak post-treatment levels between the vaccine arm and each of the no-vaccine arms separately.

Secondary immunologic endpoints:

- Determine the percentage of patients achieving at least a 10-fold expansion of myeloma-specific T cells in each treatment arm.
- Quantify myeloma antigen-specific T cells by tetramer analysis.
- Quantify T cell subsets and PD-1 expressing lymphocytes.
- Quantify NK cells and identify activation and inhibitory ligands.
- Assess NK cell-mediated killing of myeloma targets.
- Assess humoral response against whole myeloma cells and myeloma-associated antigens by SEREX and by characterizing myeloma-specific plasmablast responses.

2.3. Patient Eligibility

Patients must meet specified eligibility criteria outlined in section 2.3.1 and 2.3.2 to be registered on the study. Additional criteria must also be met to continue to successive stages of the protocol. All questions regarding eligibility criteria should be directed to the protocol coordinator at 301-251-1161.

2.3.1. Initial Inclusion Criteria

1. Patients must be considered transplant eligible by the treating physician at time of study entry.
2. Patients must meet the criteria for symptomatic multiple myeloma (Appendix A) prior to initiating systemic anti-myeloma treatment.
3. Age ≥ 18.00 years and < 71.00 years at the time of enrollment
4. Karnofsky Performance status of $\geq 70\%$
5. Patients must have $\geq 20\%$ plasma cells in the bone marrow aspirate differential ≤ 60 days prior to enrollment. If patient receives anti-myeloma therapy¹ treatment after bone marrow aspirate to assess eligibility and before bone marrow aspirate for tumor cell collection, a repeat bone marrow evaluation will be required to confirm $\geq 20\%$

¹ Anti-myeloma therapy is defined as systemic treatment intended to treat the underlying myeloma disease. Treatments intended to alleviate pain and other symptoms of disease and/or administration of $\leq 160\text{mg}$ of dexamethasone or equivalent alternative steroid dose within a 30-day period are not considered anti-myeloma therapy.

plasma cells in the bone marrow aspirate differential prior to enrollment and tumor cell collection.

6. Patients must have received ≤ 1 cycle of systemic anti-myeloma therapy¹.
7. Renal: Creatinine clearance of ≥ 40 mL/min, estimated or calculated.

2.3.2. Initial Exclusion Criteria

1. Patients with a prior autologous or allogeneic HCT
2. Patients with purely non-secretory MM [absence of a monoclonal protein (M protein) in serum as measured by electrophoresis and immunofixation and the absence of Bence Jones protein in the urine defined by use of conventional electrophoresis and immunofixation techniques and the absence of involved serum free light chain >100 mg/L]. Patients with light chain MM detected in the serum by free light chain assay are eligible.
3. Patients with a history of Plasma Cell Leukemia at any time prior to enrollment.
4. Patients with disease progression at any time prior to enrollment (see Section 3.1.3 for disease progression definition).
5. Patients seropositive for the human immunodeficiency virus (HIV).
6. Myocardial infarction within 6 months prior to enrollment or New York Heart Association (NYHA) Class III or IV heart failure (see Appendix H), uncontrolled angina, severe uncontrolled ventricular arrhythmias, or electrocardiographic evidence of acute ischemia or active conduction system abnormalities. Prior to study entry, any ECG abnormality at screening will be documented by the investigator as not medically relevant.
7. Patients with active clinically significant autoimmune disease, defined as a history of requiring systemic immunosuppressive therapy and at ongoing risk for potential disease exacerbation. Patients with a history of autoimmune thyroid disease, asthma, or limited skin manifestations are potentially eligible.
8. Patients receiving other investigational immunotherapy or anti-myeloma drugs within 14 days before enrollment.
9. Patients receiving prior treatment with daratumumab, elotuzumab, and/or any other monoclonal antibody-based therapy at any time prior to enrollment.

¹ Anti-myeloma therapy is defined as systemic treatment intended to treat the underlying myeloma disease. Treatments intended to alleviate pain and other symptoms of disease and/or administration of ≤ 160 mg of dexamethasone or equivalent alternative steroid dose within a 30-day period are not considered anti-myeloma therapy.

10. Patients with prior malignancies except resected basal cell carcinoma or treated cervical carcinoma in situ. Cancer treated with curative intent ≤ 5 years prior to enrollment will not be allowed unless approved by the Protocol Officer or one of the Protocol Chairs. Cancer treated with curative intent > 5 years prior to enrollment is allowed.
11. Female patients who are pregnant (positive β -HCG) or breastfeeding.
12. Females of childbearing potential (FCBP)¹ or men who have sexual contact with FCBP unwilling to use contraceptive techniques (Appendix D) during the length of lenalidomide maintenance therapy.
13. Patients who have received mid-intensity melphalan (>50 mg IV) as part of prior therapy.
14. Prior organ transplant requiring immunosuppressive therapy.
15. Patients who previously received lenalidomide and have experienced toxicities resulting in treatment discontinuation.
16. Patients who experienced thromboembolic events while on full anticoagulation during prior therapy with lenalidomide or thalidomide.
17. Patients unwilling to take DVT prophylaxis.
18. Patients unable or unwilling to provide informed consent.
19. Patients unable or unwilling to return to the transplant center for their assigned treatments.

2.3.3. Patient Eligibility Criteria for Randomization

1. No disease progression since initiation of systemic anti-myeloma therapy as determined ≤ 10 days prior to randomization/enrollment.
2. Received an autologous cell transplant with melphalan $200\text{mg}/\text{m}^2$ with a minimum cell dose of 2×10^6 CD34+ cells/kg (actual body weight) ≤ 12 months of enrollment onto BMT CTN 1401.
3. Tumor cell preparation must:
 - a. Express $\geq 20\%$ CD38 or CD138 by staining
 - b. Yield ≥ 10 million plasma cells²
 - c. Have a negative microbiology assessment

¹ A female of childbearing potential is a sexually mature female who: 1) has not undergone a hysterectomy or bilateral oophorectomy; or 2) has not been naturally postmenopausal for at least 24 consecutive months (i.e., has had menses at any time in the preceding 24 consecutive months).

² Plasma cell count is determined by multiplying the total mononuclear cell yield by the percent of CD38 and CD138 staining reported on the Immunohistochemistry Report

4. Mucositis and gastrointestinal symptoms resolved, off hyperalimentation and intravenous hydration.
5. No evidence of uncontrolled infection requiring systemic therapy. Patients who completed treatment for an infection but are continuing antibiotics, anti-viral, or anti-fungal therapy for prophylaxis are eligible to continue on protocol.
6. Platelet count $\geq 75,000/\text{mm}^3$ (without transfusion in previous 7 days).
7. Absolute neutrophil count (ANC) $\geq 1,500/\text{mm}^3$ without filgrastim administration within 7 days, or pegfilgrastim within 14 days of measurement.
8. Hepatic: bilirubin $< 2\times$ the upper limit of normal and ALT and AST $< 2.5\times$ the upper limit of normal. (Patients who have been diagnosed with Gilbert's Disease are allowed to exceed the defined bilirubin value of $2\times$ the upper limit of normal.)
9. Renal: Creatinine clearance of ≥ 40 mL/min, estimated or calculated. Patients with creatinine clearance ≥ 30 but < 40 will be considered with review/approval from the protocol chairs or officer if the cause of renal insufficiency is associated with multiple myeloma.
10. All study participants must be registered into the mandatory Revlimid REMs program, and be willing and able to comply with the requirements.
11. Females of childbearing potential (FCBP) as defined in section 2.7.1.1 must have a negative serum pregnancy test with a sensitivity of at least 50 mIU/mL within 10 – 14 days prior to and again within 24 hours of prescribing lenalidomide (prescriptions must be filled within 7 days)
12. FCBP must either commit to abstain continuously from sexual intercourse or use TWO acceptable methods of birth control, one highly effective method and one additional effective method AT THE SAME TIME, at least 4 weeks before she starts taking lenalidomide, during therapy, during dose interruptions, and continuing for 4 weeks following discontinuation of lenalidomide.
13. FCBP must agree to ongoing pregnancy testing as required by the RevlimidREMs program.
14. Men must agree to use a latex condom during sexual contact with females of child bearing potential even if they have had a successful vasectomy while taking lenalidomide, during dose interruptions and for 28 days after discontinuing lenalidomide.
15. Patients must be willing to receive DVT prophylaxis.

2.4. Treatment Plan

2.4.1. Treatment Prior to Enrollment

Patients must have ≤ 1 cycle of anti-myeloma therapy prior to enrolling on the study. It is recommended whenever clinically feasible to minimize the treatment administered prior to enrollment to ensure adequate tumor cell collection and preparation for vaccine production. Patients receiving treatment with daratumumab or other CD38 monoclonal antibodies prior to enrollment should not be considered for this study.

2.4.2. Tumor Cell Collection

An initial eligibility assessment will be performed at time of study entry prior to tumor collection for immune monitoring and subsequent vaccine production. Eligible patients will undergo aspiration of 30 mL of bone marrow from which myeloma cell preparations will be generated (Appendix C and BMT CTN 1401 Manufacturing, Clinical and Reference SOPs). Bone marrow will be aspirated into sterile heparinized syringes with approximately 5-8 mL aspirated per syringe to avoid significant hemodilution of bone marrow mononuclear cells. Bone marrow aspirate samples will be transferred in sterile syringes to the cell manipulation facility, and mononuclear cells will be isolated by ficoll density gradient centrifugation. Autologous plasma will be obtained by harvesting supernatant following centrifugation of 50 mL of peripheral blood. An aliquot of the tumor cells will be sent to BIDMC for centralized immunohistochemical staining as detailed in Appendix C. Tumor lysate will be prepared by freeze/thawing or sonication of an aliquot of tumor cells for immunological analysis. The remaining myeloma cells will be frozen in 10% DMSO/90% autologous plasma stored in liquid nitrogen for potential subsequent vaccine generation.

2.4.2.1. Second Tumor Cell Collection

If the initial bone marrow harvest does not yield a sufficient cell count as required by the BMT CTN 1401 Vaccine Product Release criteria, a second collection may be considered if the following items are completed *prior* to the second collection:

- Submission and review of an occurrence report in AdvantageEDC
- Written approval from the BMT CTN 1401 Protocol Chairs/Officer and Medical Monitor is received
- All corrective actions are completed and appropriate documentation is submitted to the BMT CTN DCC
- Final written approval from the BMT CTN DCC is received by the site

2.4.3. Systemic Anti-myeloma Therapy

Pre-transplant systemic anti-myeloma therapy will be administered per investigator discretion following enrollment and tumor cell collection.

2.4.4. Stem Cell Mobilization

Stem cell mobilization and collection should be done according to institutional guidelines.

2.4.5. Autologous Stem Cell Transplantation

Autologous hematopoietic cell transplant will be done with high-dose melphalan (200mg/m²) at the schedule and timing according to institutional practices and must be within 12 months of enrollment on to the study

2.4.5.1. Peripheral blood stem cell infusion

All patients will receive an autologous graft with a minimum cell dose of 2.0×10^6 CD34+ cells/kg patient actual body weight per autologous transplantation. The graft may not be CD34+ selected or otherwise manipulated to remove tumor or other cells. Cryopreservation and thawing of product will comply with FACT standards and local institutional practice.

2.4.6. Maintenance Therapy

All patients who meet the eligibility criteria to proceed to randomization and who have completed the post-transplant disease response assessment will be randomized between day 53 and day 80 post-transplant and within 10 days of a disease response assessment.

Patients will begin maintenance therapy with lenalidomide between 90 and 100 days after stem cell infusion. Lenalidomide will be administered initially at a dose of 10 mg per day continuously. Cycle duration during maintenance therapy is 28 days. Patients will continue lenalidomide for two years from initiation of therapy or until one of the criteria in section 2.8 applies. For patients receiving GM-CSF and/or Vaccine, lenalidomide is recommended to be taken in the evening at least 4 hours after vaccination to avoid confusion with immediate vaccine and/or GM-CSF site reactions.

Patients who stop maintenance therapy for any reason prior to two years from initiation of maintenance will continue to be followed per protocol for 3 years post maintenance initiation. Upon completion of protocol specified maintenance therapy, patients will be treated at the investigator's discretion. However, it is recommended that patients continue receiving lenalidomide as per standard of care.

2.4.6.1. Lenalidomide maintenance dose modifications

Lenalidomide dose will be adjusted for toxicity. Table 2.4.5B describes the dose reduction steps to be utilized during maintenance therapy. In the presence of lenalidomide-related toxicities (Table 2.4.5A), the study drug will be held until the toxicity resolves then restarted at a reduced dose as described in Table 2.4.5B. If lenalidomide is held for >8 weeks for any reason, study chair approval must be obtained prior to re-starting lenalidomide. Patients will be given DVT prophylaxis. DVT prophylaxis for patients will be given per institutional guidelines. See section 2.5.2 for suggested dose and more information.

Table 2.4.6A – Required Treatment Modification Guidelines for Lenalidomide Maintenance Therapy for Non-Hematologic Toxicities

Grade by NCI CTCAE# ¹	Action
Grade 3 neutropenia associated with fever (temperature $\geq 38.5^{\circ}\text{C}$) or Grade 4 neutropenia	Hold lenalidomide. Follow CBC weekly. If the toxicity resolves to \leq grade 2 restart lenalidomide at next lower dose level. If neutropenia is the only toxicity for which a dose reduction is required, granulocyte growth factors could be considered at the discretion of the treating physician and the lenalidomide dose maintained. .
Thrombocytopenia (platelet count $< 30,000/\text{mm}^3$)	Hold lenalidomide. Follow CBC weekly. If the toxicity resolves to \leq grade 2 restart lenalidomide at next lower dose level.
Non-blistering rash Grade 3 (Generalized rash $\geq 25\%$ BSA)	Hold lenalidomide: Follow weekly. Treatment with antihistamine may be initiated at the discretion of the investigator. If the toxicity resolves to \leq grade 1 restart lenalidomide at next lower dose level.
Non-blistering rash Grade 4	Discontinue lenalidomide permanently. Continue to follow the patient per-protocol
Desquamating (blistering) rash any Grade	Discontinue lenalidomide permanently. Continue to follow the patient per-protocol
Venous thrombosis/embolism \geq Grade 3 (DVT or cardiac thrombosis; intervention indicated)	Hold lenalidomide and start anticoagulation per institutional guidelines. Restart therapy and maintain dose level.
Other non-hematologic toxicity assessed as Lenalidomide-related \geq Grade 3	Hold lenalidomide. If the toxicity resolves to \leq grade 2 restart lenalidomide at next lower dose level.
Hyperthyroidism or Hypothyroidism	Hold lenalidomide. Evaluate etiology, and initiate appropriate therapy. Restart lenalidomide at next lower dose level .
Pregnancy³	Discontinue lenalidomide study drug.

¹Please consult NCI CTCAE version 4 <http://ctep.cancer.gov/reporting/> for complete **Grade** descriptions. The “ \geq **Grade 3**” descriptions listed above are minimums

³If a subject, or the partner of a male study subject, misses her period or if her pregnancy test or her menstrual bleeding is abnormal, pregnancy testing and counseling must be performed (Section 4.4.5).

Table 2.4.6B – Lenalidomide Dose Reduction Steps During Lenalidomide Maintenance Therapy

Lenalidomide Dose Reduction Steps for Non-Hematologic Toxicity	
Dose at Time of Toxicity	Dose reduction
10 mg daily	5 mg daily
5 mg daily	5 mg daily for 21 days every 28 days
5 mg daily for 21 days every 28 days	2.5 mg daily for 21 days every 28 days
2.5 mg daily for 21 days every 28 days	Discontinue lenalidomide

2.4.6.2. Lenalidomide maintenance dose re-escalation

If a dose reduction has occurred and ANC $\geq 1000/\mu\text{L}$ and platelet count is $\geq 75,000/\mu\text{L}$, the study drug dose may be re-escalated as shown on Table 2.4.5c, one step per cycle to a maximum of 10 mg daily.

Table 2.4.6C – Lenalidomide Dose Re-Escalation Steps During Lenalidomide Maintenance Therapy

Current Patient Dose	Dose Re-Escalation
2.5 mg daily for 21 days every 28 days	5 mg daily for 21 days every 28 days
5 mg daily for 21 days every 28 days	5 mg daily
5 mg daily	10 mg daily
10 mg daily	No more dose escalations permitted

2.4.7. Vaccine Manufacturing and Administration

Vaccine preparation is outlined in detail in the BMT CTN 1401 SOPs and will be performed locally at each participating institution. Patients randomized to receive the DC/MM fusion vaccine will undergo leukapheresis for DC and vaccine generation per the BMT CTN 1401 manufacturing, clinical, and reference SOPs. Leukapheresis should occur by day 85 post-transplant to ensure sufficient time for vaccine production and sterility testing. The leukapheresis product will be transferred in a sterile container to the cell manipulation facility, and PBMC will be isolated from the leukapheresis product and cultured in the presence of autologous plasma for 1 hour and then cultured for 5-7 days with GM-CSF and IL-4. Cultures will be re-fed with cytokines, GM-CSF, and IL-4 after 5-7 days. Twenty-five ng/mL of TNF α will be added for 48-72 hours to induce DC maturation after 5-7 days. An aliquot of the DC preparation will be sent to BIDMC for immunocytochemical staining for immunophenotypic analysis. Tumor cells and DC preparations will be co-cultured at ratio of 1:3 and washed in serum-free medium. After low-speed centrifugation, the cell pellet will be re-suspended in 50% solution of polyethylene glycol (PEG) and will be progressively diluted by the slow addition of serum-free medium. An aliquot of the DC, tumor, and fusion cell preparations will be sent to BIDMC for immunocytochemical analysis for vaccine characterization at the timepoints indicated in the BMT CTN 1401 manufacturing SOPs. The vaccine doses will be prepared per the Vaccine Dose Guidelines in section 2.4.7.1. The fusion cells will then be irradiated at 30 Gy, and frozen in 10% DMSO/90% autologous plasma in liquid nitrogen. An aliquot of the vaccine product will be sent for microbiological assessment consisting of endotoxin, mycoplasma and sterility testing as per the BMT CTN 1401 manufacturing SOPs. Release criteria for vaccine administration are:

- 1) At least 20% of tumor cell prep must express CD38 or CD138 by staining
- 2) Tumor cell prep must yield ≥ 10 million plasma cells¹
- 3) > 50% of DC prep express CD86
- 4) Viability of DC prep > 50%
- 5) Fusion efficiency > 15%
- 6) Fusion viability > 50%
- 7) Sterility, mycoplasma, and endotoxin assays are negative

¹ Plasma cell count is determined by multiplying the total mononuclear cell yield by the percent of CD38 and CD138 staining reported on the Immunohistochemistry Report

At time of administration, fusion cells will be thawed. Patients randomized to the vaccine arm will receive the DC/myeloma fusion vaccine/GM-CSF (vaccine) on day 1 of cycles 2, 3, and 4 of lenalidomide maintenance. Vaccine will be administered by subcutaneous injection with 100 ug GM-CSF given subcutaneously at the vaccine site on day of vaccination and daily for a total of 4 days of each cycle. Patients who are randomized to the vaccine arm and are unable to receive vaccine for any reason, will receive lenalidomide alone for the duration of the study per section 2.4.6.

All vaccine related procedures are detailed in the BMT CTN 1401 manufacturing, clinical, and reference SOPs which must be adhered to by all sites for each patient enrolled on the study. Patients who are unable to receive vaccine for any reason (i.e. toxicity, failure to meet release criteria, etc) should receive lenalidomide only for the duration of the study.

2.4.7.1. Vaccine Dose Guidelines

The target dose is 3×10^6 fusion cells per vaccine. A minimum of 3×10^6 fusion cells will be required to proceed with vaccine administration. Patients who have $< 3 \times 10^6$ fusion cells will not proceed with vaccination.

Table 2.4.7A – Vaccine Dose Guidelines

Total Fusion Cell Yield	Dose Guidance
$\geq 9 \times 10^6$ fusion cells	<ul style="list-style-type: none"> - 3 doses of 3×10^6 fusion cells are cryopreserved for patient administration - Excess product should be cryopreserved in aliquots of 3×10^6 fusion cells when possible.
$\geq 3 \times 10^6$ fusion cells and $< 9 \times 10^6$ fusion cells	<ul style="list-style-type: none"> - Divide product in to 3 equal doses for cryopreservation and patient administration
$< 3 \times 10^6$ fusion cells	<ul style="list-style-type: none"> - Patient will not receive vaccine.

2.4.7.2. Vaccine Production Occurrences

Deviations from the manufacturing process outlined in the BMT CTN 1401 manufacturing, clinical, and reference SOPs will be documented on the Occurrence Report form in AdvantageEDC as per the Occurrence Report SOP (BMT 1401-RF 04). Following submission of an occurrence report, sites will be required to complete the following items *prior* to enrolling additional patients and proceeding with vaccine production:

- All queries and corrective actions associated with the report must be completed
- The BMT CTN 1401 protocol chairs/officer and BMT CTN DCC will provide approval for continuing enrollment on the study.

2.4.7.2.1. Failure to Meet Release Criteria

Sites experiencing difficulties with production as evidenced by failure to meet release criteria for 3 products will have vaccine production temporarily halted while undergoing onsite review of procedures by a member from the protocol team and central site. Additional training or guidance will be provided if needed.

Table 2.4.7B –Vaccine Administration Guidelines for Vaccine, GM-CSF and/or Lenalidomide Toxicities

Toxicity	Action
Hold/delay in lenalidomide therapy	Hold/delay vaccine. When lenalidomide is restarted, adjust date of vaccine administration to coincide with Day 1 of cycles 2, 3 and 4 of lenalidomide maintenance.
Discontinuation in lenalidomide therapy	Administer vaccine at monthly intervals for a total of 3 doses
Grade 4 toxicity judged to be related to vaccine alone	No additional vaccines.
Grade 3 toxicity judged to be related to vaccine alone	Hold Vaccine Dose. If the toxicity resolves to \leq grade 2 by time of subsequent vaccine administration, vaccine may continue.
Grade 3-4 toxicity judged to be related to GM-CSF alone	Omit GM-CSF administration for subsequent vaccine injections or subsequent scheduled administrations in the non-vaccine arm. Re-challenging GM-CSF administered when toxicity resolves to \leq grade 2 can be considered at the discretion of the treating investigator. Vaccine administration may continue.
Grade 3-4 toxicity possibly related to combination of lenalidomide and vaccine	Follow lenalidomide dose adjustments. Vaccine administration may continue.

2.4.8. GM-CSF Administration

For patients randomized to the vaccine and GM-CSF plus lenalidomide maintenance and lenalidomide maintenance plus GM-CSF arms, GM-CSF will be administered at cycles 2, 3, and 4 of maintenance therapy. 100 ug GM-CSF will be given subcutaneously (at the vaccine site for patients receiving vaccine) on days 1-4 of cycles 2, 3, and 4 per the BMT CTN 1401 manufacturing SOPs.

On the first day of GM-CSF administration, the clinical research nurse will administer 100 ug of GM-CSF subcutaneously. The patient will be trained to inject the remaining three GM-CSF injections (100ug dose once a day at the vaccination site) for self-administration subcutaneously at home. Alternatively, if patients are unwilling or unable to self-administer the GM-CSF, they can return to the transplant center for subsequent injections. For patients receiving vaccine, tumor vaccine will be administered before the GM-CSF injection. GM-CSF is given on Day 1 at time of vaccination and around the same time of day (+/- an hour whenever feasible) for subsequent doses. If a patient experiences a grade 2 or higher (graded according to CTCAE v4.0) injection site reaction during the days when GM-CSF is administered, subsequent doses of GM-CSF may be held until the injection site reaction improves to grade 1 or less at the discretion of the treating investigator. For subsequent administrations, patients will receive pre-medication with diphenhydramine (Benadryl) 25-50 mg and/or acetaminophen (Tylenol) 650-1000 mg to minimize potential allergic-related symptoms.

2.5.Supportive Care

2.5.1. Post Autologous Transplantation(s)

All supportive care will be given at the investigators discretion.

2.5.2. Deep Vein Thrombosis (DVT) and Pulmonary Embolism (PE) Prophylaxis

Prophylaxis for DVT/PE will be done according to institutional guidelines. Upon initiation of lenalidomide maintenance it is recommended that patients without a history of thromboembolic events should be on aspirin daily (81mg, 165mg or 350 mg); low molecular weight heparin or warfarin is usually administered in patients with a history of thromboembolic events or those that cannot tolerate aspirin.

2.6.Risks and Toxicities

Recipients of autologous transplantation incur risks from pre-transplant conditioning and post-transplant therapy, which must be weighed against the risk of the disease for which the transplant is prescribed. Major risks following transplantation include: 1) Infection, which can be bacterial, viral, parasitic, or fungal. Often, these infections are life-threatening, particularly when caused by viral or fungal agents, and are associated with high mortality in the transplant population; 2) Graft Failure can occur and is associated with a high-risk of mortality; 3) End Organ Damage of one or more major organs may occur as a result of reactions to drugs (e.g., melphalan, antibiotics, anti-fungal medications, etc.), and as a result of destructive processes (e.g., infection, etc.) and may have a fatal outcome; 4) Progression of MM may occur; 5) Unknown Toxicities may occur in any individual patient due to multiple events and cumulative effects which may involve any and all organs, including the brain. Brain damage can result in severe loss of cognitive or neurologic function; and, 6) Death. These risks may or may not be increased by the post-transplant therapies to be evaluated in this protocol.

2.6.1. Therapy Toxicities

All toxicities will be graded using the Common Terminology Criteria for Adverse Events (CTCAE) Version 4.0 with BMT CTN 1401 specific definitions when appropriate.

2.6.2. Lenalidomide

Common toxicities described for lenalidomide include:

- Neurologic: Somnolence, dizziness, headache, tremor, asthenia, paresthesia, and numbness
- Hematologic: anemia, neutropenia, leucopenia, lymphopenia and thrombocytopenia; thromboembolic events (deep vein thrombosis and pulmonary embolism).
- Gastrointestinal: Constipation, dehydration, dry mouth, diarrhea, dyspepsia, nausea, vomiting and stomatitis.
- Constitutional: Weakness, insomnia, rigors, chills, sweating, weight loss and fever.
- Reproductive: teratogenicity and miscarriage.
- Musculoskeletal: arthralgia, back/neck pain, joint pain, muscle cramp and weakness.

- Cardiac: hypotension.
- Dermatologic: rash, dry skin, itching.
- Endocrine: hypothyroidism.
- Infection.
- Pulmonary: cough, dyspnea.
- Metabolic: hypokalemia, liver damage.
- Renal: increased creatinine, renal failure.
- Second Primary Malignancies

Pregnancy reporting:

See Section 4.2.3, Adverse Event Reporting.

Other instructions related to lenalidomide:

During maintenance, only one cycle (maximum 28-day supply) of therapy may be dispensed to the patient each month. If a dose of lenalidomide is missed, it should be taken as soon as possible on the same day. If it is missed for the entire day, it should not be made up. Patients taking more than the prescribed dose of lenalidomide should be instructed to seek emergency medical care if needed and contact study staff immediately. See Section 2.7.1.1, Chapter 4, Appendices D (Risks of Fetal Exposure, Pregnancy Testing Guidelines and Acceptable Birth Control Methods) and E (Revlimid REMS program).

2.6.3. DC/Fusion Vaccine - Manufacturing and Administration**2.6.3.1. Tumor Collection**

All eligible patients will undergo a bone marrow biopsy and collection of 30 mL of bone marrow aspirate. The risks of this procedure are not different than bone marrow aspirates performed for diagnosis or disease status assessments. The risks include: bleeding, pain in the site of marrow collection, claudication and bruising.

2.6.3.2. Dendritic Cell Precursor Collection

Collection of dendritic cell precursors will occur in patients randomized to receive vaccine post-transplant. Collection will be done through leukapheresis in a single session. Risks of this collection are related to placement of an intravenous leukapheresis catheter that can be associated with bleeding and pain. The leukapheresis procedure has a risk of hypocalcemia, thrombocytopenia, anemia, leukopenia and bleeding diathesis.

2.6.3.3. Vaccine Administration

Vaccines will be administered at three time points approximately one month apart. GM-CSF will also be administered at the site of vaccine for 4 consecutive days. Toxicities related to the vaccine are:

- Hematologic: Leucopenia, Neutropenia
- Neurologic: Headache, light headedness
- Gastrointestinal: Diarrhea,
- Constitutional: Weakness, chills and fever. Pain at tumor site, fatigue, night sweats
- Musculoskeletal: arthralgia and myalgia.
- Dermatologic: rash, itching, edema and local reaction in the site of infusion (swelling, pain, bruising, erythema, and pruritus).
- Endocrine: elevated TSH.
- Other: ANA positivity, candida infection

In prior studies, one participant with myeloma and a history of deep vein thrombosis developed a pulmonary embolus, which was determined to be serious and possibly related to the vaccine and GM-CSF.

2.6.4. GM-CSF

Potential side effects of GM-CSF treatment include fever, chills, nausea, vomiting, diarrhea, fatigue, weakness, headache, decreased appetite, facial flushing, bone and muscle pain, local reaction at the site of injections, rashes, low blood pressure, shortness of breath and low blood counts. Rarely patients may develop blood clots, rapid or irregular heartbeats, feeling of faintness, and allergic reactions or fluid retention, including potential fluid retention in lungs or around the heart.

2.7. Study Drug Supply

2.7.1. Lenalidomide

Lenalidomide will be provided by Celgene and distributed by Biologics, Incorporated. Patients must be registered in the Revlimid REMS program in order to receive lenalidomide through the program (please see Appendix E).

2.7.1.1. Lenalidomide (NSC 703813)

NOTE:

Before lenalidomide is dispensed, patients must 1) have a negative pregnancy test (if applicable) and 2) study patients must be counseled through the Revlimid REMS program. A maximum 28-day supply may be dispensed to a patient at one time. For more information please refer to www.revlimidrems.com. Only use the study specific drug request form per patient for ordering through Biologics.

Chemical Name: 3-(4'-amino-1,3-dihydro-1-oxo-2*H*-isoindol-2-yl)-2,6-piperidinedione

Other Names: CC-5013, Revlimid™, CDC-501

Classification: Immunomodulatory Agent

CAS Registry Number: 191732-72-6

Molecular Formula: $C_{13}H_{13}N_3O_3$

M.W.: 259.25

Mechanism of Action:

Lenalidomide, a thalidomide analog, is an immunomodulatory agent with a spectrum of activity that is still under investigation. Some of its effects include inhibition of inflammation, inhibition of angiogenesis, inhibition of hematopoietic tumor cell proliferation, modulation of stem cell differentiation and up regulating responses of T cells and NK cells.

Drug Supply and Storage:

Celgene supplies and Biologics, Inc. distributes lenalidomide 5 mg (size 2) and 2.5 mg hard gelatin capsules in tamper-evident, child-resistant, opaque, high density polyethylene (HDPE) bottles with HDPE caps. Bottles will contain a sufficient number of capsules per container for one cycle of dosing.

The capsules also contain anhydrous lactose, microcrystalline cellulose, croscarmellose sodium, and magnesium stearate.

Lenalidomide (Revlimid®) will be provided to research subjects for the duration of their participation in this trial at no charge to them or their insurance providers. Lenalidomide will be provided in accordance with the Revlimid REMS™ program of Celgene Corporation. Per standard Revlimid REMS™ requirements all physicians who prescribe lenalidomide for research subjects enrolled into this trial, and all research subjects enrolled into this trial, must be registered in and must comply with all requirements of the Revlimid REMS™. Prescriptions must be filled within 7 days for FCBP and 14 days for all other risk categories.

All study participants must be registered into the mandatory Revlimid REMS™ program, and be willing and able to comply with the requirements of the Revlimid REMS™ program.

Any unused lenalidomide supplies distributed through the Revlimid REMS™ program must be returned to the clinical site and destroyed per institutional guidelines.

Administration:

Take lenalidomide by mouth with or without food. Do not crush, chew or open capsules.

Dispensing:

Only enough lenalidomide for one cycle may be dispensed at one time.

Patient Care Implications and Counseling:**Risks Associated with Pregnancy**

Lenalidomide is structurally related to thalidomide. Thalidomide is a known human teratogenic active substance that causes severe life-threatening birth defects. An embryofetal development study in animals indicates that lenalidomide produced malformations in the offspring of female monkeys who received the drug during pregnancy. The teratogenic effect of lenalidomide in humans cannot be ruled out. Therefore, a risk minimization plan to prevent pregnancy must be observed.

Definition of female of childbearing potential (FCBP)

This protocol defines a female of childbearing potential as a sexually mature female who: 1) has not undergone a hysterectomy or bilateral oophorectomy or 2) has not been naturally postmenopausal (amenorrhea following cancer therapy does not rule out childbearing potential) for at least 24 consecutive months (i.e., has had menses at any time in the preceding 24 consecutive months).

Before starting study drug:

Female Subjects:

- FCBP must have two negative pregnancy tests (minimum sensitivity of 50 mIU/mL) prior to prescribing study drug. The first pregnancy test must be performed within 10-14 days prior to prescribing study drug and the second pregnancy test must be performed within 24 hours prior to prescribing study drug (prescriptions must be filled within 7 days as required by Revlimid REMS™). The subject may not receive study drug until the Investigator has verified that the results of these pregnancy tests are negative.
- FCBP must commit either to abstain continuously from heterosexual intercourse or to use TWO methods of reliable birth control AT THE SAME TIME—one highly effective form of contraceptions and one additional effective contraceptive method as defined below. Contraception must begin 4 weeks prior to initiating treatment with lenalidomide, during therapy, during dose interruptions, and continuing for 4 weeks following discontinuation of lenalidomide.
 - HIGHLY EFFECTIVE:
 - Tubal Ligations
 - IUD
 - Hormonal contraceptives (birth control pills, injections, hormonal patches, vaginal rings or implants)
 - Partners Vasectomy
 - EFFECTIVE:
 - Male latex or synthetic condom
 - Diaphragm
 - Cervical cap

Male Subjects:

- Must agree to use a latex or synthetic condom during sexual contact with females of childbearing potential while participating in the study, during dose interruptions and for at least 28 days following discontinuation from the study even if he has undergone a successful vasectomy.

All Subjects:

- Only enough lenalidomide for one cycle of therapy may be dispensed with each cycle of therapy.
- If pregnancy or a positive pregnancy test does occur in a study subject or the partner of a male study subject during study participation, lenalidomide must be immediately discontinued.

Potential Drug Interactions:

Periodic monitoring of digoxin levels is recommended during co-administration with lenalidomide.

Monitor patients receiving concomitant warfarin per standard practice guidelines.

Lenalidomide is not a substrate of human CYP enzymes, nor is it an inhibitor or inducer.

Drug Ordering and Accountability:

The REMs program provides education and counseling on the risks of fetal exposure, blood clots and reduced blood counts. Counseling will be provided by Biologics, Inc. prior to drug distribution. Please refer to Appendix E (Lenalidomide REMs program). The patient will be required to receive counseling every 28 days during treatment with lenalidomide, follow the pregnancy testing and birth control requirements of the program that are appropriate in order to take the telephone surveys regarding compliance with the program. All physicians must be registered prescribers of Revlimid® in the REMs program. Physician registration allows access to the REMs software to enroll patients in the REMs program. The prescriber should submit the Registration Form via fax number 919-256-0794 or REMsOnline (**RAO**) for Revlimid®) to Celgene Customer Care. Please reference Appendix E (REMsprogram) and follow the directions for submitting the registration. Biologics, the distributor of the lenalidomide, will not dispense or ship Revlimid® prior to Celgene's receipt of registration. Prescription information **MUST BE** entered using the REMs study specific electronic prescription form referenced in Appendix E (REMs program). An authorization **number** must be on the prescription form at the time of faxing. Prescriptions for Revlimid® must be sent to Biologics Clinical Trial Division at the following FAX number: 919-256-0794. A maximum of a 28-day supply of Revlimid® may be dispensed per cycle sent to the actual address noted on the REMs electronic prescription form. Biologics will verify the authorization number and complete the patient counseling. Patients will be provided with instructions from Biologics with each new dispense on the procedures for return of any unused Revlimid® capsules. Refer to Appendix E (REMsprogram).

2.8. Patient Withdrawal from Study and Off Study Criteria

Patients on the BMT CTN 1401 study have voluntarily agreed to the study and may withdraw at anytime. Additionally, the treating physician may discontinue therapy with reasonable justification. The following criteria would result in discontinuation of study therapy:

- Disease progression
- Unacceptable adverse event(s)
- Participant decides to withdraw from the study
- General or specific changes in the participant's condition render the participant unacceptable for further treatment in the opinion of the treating investigator.

After randomization, patients who stop therapy for any reason prior to two years from initiation will continue to be followed per protocol for 3 years post maintenance initiation unless the patient withdraws consent for study follow up. In the event of patient withdrawal, written documentation of the withdrawal must be provided to the BMT CTN.

CHAPTER 3

3. STUDY ENDPOINTS AND DEFINITIONS

3.1. Definition of Disease Status

Patients' disease status at each data collection period will be evaluated based on the International Uniform Response Criteria¹⁹. Until disease progression, all disease classifications are relative to the patient's disease status prior to autologous transplant. At time of disease progression, disease classifications are relative to the patient's best response since time of study entry.

3.1.1. Response Categories

Stringent Complete Response (sCR):

sCR requires, in addition to CR (defined below), all of the following:

- Normal free light chain ratio (FLC).
- Absence of clonal cells in bone marrow by immunohistochemistry or immunofluorescence.

Complete Response (CR):

CR requires *all* of the following:

- Absence of the original monoclonal paraprotein in serum and urine by routine electrophoresis and by immunofixation. The presence of new monoclonal bands consistent with oligoclonal immune reconstitution does not exclude CR.
- Less than 5% plasma cells in a bone marrow aspirate and also on trephine bone biopsy, if biopsy is performed.
- No increase in size or number of lytic bone lesions on radiological investigations (development of a compression fracture does not exclude CR)*.
- Disappearance of soft tissue plasmacytomas.

**If not clinically indicated, radiographs are not required to document CR.*

Patients in whom some, but not all, the criteria for CR are fulfilled are classified as partial responses (see below), providing the remaining criteria satisfy the requirements for partial response. This includes patients in whom routine electrophoresis is negative but in whom immunofixation has not been performed.

Very Good Partial Remission (VGPR)

VGPR requires, in addition to PR (defined below), all of the following:

- Serum or urine paraprotein detectable by immunofixation but not on electrophoresis.

OR

- Greater than or equal to 90% reduction in serum paraprotein plus urine paraprotein <100 mg/24hrs.

- For free light chain only disease, VGPR requires a 90% reduction of involved light chain

Partial Response (PR)

PR requires one of the following:

- Greater than or equal to 50% reduction in the level of the serum monoclonal paraprotein and reduction in 24 hour urinary monoclonal paraprotein either by greater than or equal to 90% or to <200 mg/24 hours in light chain disease.
- If the only measurable non-bone marrow parameter is FLC, greater than or equal to 50% reduction in the difference between involved and uninvolved FLC levels or a 50% decrease in level of involved FLC with 50% decrease in ratio,
- If the bone marrow is the only measurable parameter, greater than or equal to 50% reduction in bone marrow plasma cells given that the baseline count was greater or equal to 30%,
- Greater than or equal to 50% reduction in the size of soft tissue plasmacytomas if present at baseline (by radiography or clinical examination).

Stable Disease (SD)

- Patients who do not meet criteria for sCR, CR, VGPR, partial response or progressive disease (section 3.1.1.2) are considered to have stable disease (SD).

3.1.2. Progressive Disease (PD)**Disease Progression (PD)**

Progression from CR or sCR requires one or more of the following:

- A reappearance of serum monoclonal paraprotein, with a level of at least 0.5 g/dL.
- 24-hour urine protein electrophoresis with at least 200 mg paraprotein/24 hours.
- Abnormal FLC levels of >10 mg/dl, only in patients without measurable paraprotein in the serum and urine.
- At least 10% plasma cells in a bone marrow aspirate or on trephine biopsy.
- Definite increase in the size of existing bone lesions or soft tissue plasmacytomas.
- Development of new bone lesions or soft tissue plasmacytomas.
- Development of hypercalcemia (corrected serum Ca >11.5 mg/dL or >2.8 mmol/L) not attributable to any other cause.

Progressive Disease (PD)

For patients not in CR or sCR, progressive disease requires one or more of the following measured from the time of randomization:

- >25% increase in the level of the serum monoclonal paraprotein, which must also be an absolute increase of at least 0.5 g/dL.
- >25% increase in 24-hour urine protein electrophoresis, which must also be an absolute increase of at least 200 mg/24 hours.
- Absolute increase in the difference between involved and uninvolved FLC levels (absolute increase must be >10 mg/dl), only in patients without measurable paraprotein in the serum and urine.
- >25% increase in plasma cells in a bone marrow aspirate or on trephine biopsy, which must also be an absolute increase of at least 10%.
- Definite increase in the size of existing bone lesions or soft tissue plasmacytomas.
- Development of new bone lesions or soft tissue plasmacytomas.
- Development of a compression fracture does not exclude continued response and may not indicate progression.
- Development of hypercalcemia (corrected serum Ca >11.5 mg/dL or >2.8 mmol/L) not attributable to any other cause.

3.2.Primary Endpoint

The primary objective of this randomized trial is to compare the proportion of patients alive and in complete response (CR or sCR) at one year post transplant between patients receiving DC/myeloma vaccine/GM-CSF with lenalidomide maintenance therapy to those receiving lenalidomide maintenance therapy with or without GM-CSF.

3.3.Clinical Secondary Endpoints**3.3.1. Response to treatment**

The trial will assess the rates of VGPR or better (VGPR, nCR, CR and sCR) responses) according to the International Uniform Response Criteria (Section 3.2) at specific time points in both arms from the time of randomization. Assessment of disease response will be done immediately prior to randomization, prior to cycles 3, 6, and then every 3 months until approximately 22 months from randomization or evidence of disease progression. Additionally, assessments of disease response will be conducted at 6, 12, and 24 months post-transplant. Rates of VGPR or better responses will be compared between the two arms. For patients not in CR at randomization, the rate of CR conversion (CR and sCR) will be compared between the vaccine and no-vaccine arms combined. A secondary pairwise analysis will compare the CR rates at 1 year between the vaccine arm, lenalidomide/GM-CSF arm and lenalidomide alone arm.

3.3.2. Myeloma Progression

Myeloma progression defined according to Section 3.2 or initiation of off protocol anti-myeloma therapy is the event for this endpoint. Death without documentation of disease progression is the competing event. Patients alive without disease progression at last contact are considered censored for this event. The cumulative incidence of myeloma progression will be compared between vaccine and no-vaccine arms combined. A secondary pairwise analysis will compare the cumulative incidence of myeloma progression between the vaccine arm, lenalidomide/GM-CSF arm and lenalidomide alone arm.

3.3.3. Minimal Residual Disease Assessment

Minimal residual disease (MRD) is defined as the presence of malignant plasma cells detected by multicolor flow cytometry among patients who are in complete remission. Multichannel flow cytometry will be used to establish MRD based on the presence of malignant plasma cells that are CD45 (-/dim), CD38+, CD138+, CD19-, CD56+ kappa or lambda restricted. The result will determine whether that patients has MRD (MRD +) or not (MRD -). The proportion of patients who are MRD negative at one year will be compared between vaccine and no-vaccine arms combined. A secondary pairwise analysis will be conducted comparing the proportion of patients who are MRD negative at one year between the vaccine arm, lenalidomide/GM-CSF arm and lenalidomide alone arm.

3.3.4. Treatment-related Mortality

TRM is defined as death occurring in a patient from causes other than disease relapse or progression. Disease progression is the competing event for TRM. Patients alive without disease progression at last contact are considered censored for this event. TRM from time of randomization will be compared between vaccine and no-vaccine arms combined starting at time of randomization. A secondary pairwise analysis will compare TRM between the vaccine arm, lenalidomide/GM-CSF arm and lenalidomide alone arm.

3.3.5. Incidence of Toxicities Grade ≥ 3 per CTCAE vesion 4.0

All Grade ≥ 3 toxicities will be tabulated for treatment arms. The proportion of patients developing Grade ≥ 3 toxicity will be compared between the vaccine and no-vaccine arms combined until disease progression or end of follow up. A secondary pairwise analysis will compare the incidence of toxicities between the vaccine arm, lenalidomide/GM-CSF arm and lenalidomide alone arm.

3.3.6. Incidence of Infections per BMT CTN Technical MOP

The incidence of definite and probable viral, fungal and bacterial infections will be tabulated for each patient. The proportion of patients in each treatment arm with these infections will be compared from randomization until disease progression or end of follow up.

3.3.7. Progression-free survival

Patients are considered a failure of this endpoint if they die or suffer from disease progression. The time to this event is the time from randomization to progression, death, initiation of non-protocol anti myeloma therapy, loss to follow up or end of study whichever comes first and it will be compared between the vaccine and no-vaccine arms combined from time of randomization.

A secondary pairwise analysis will compare progression-free survival between the vaccine arm, lenalidomide/GM-CSF arm and lenalidomide alone arm.

3.3.8. Overall Survival

The event is death from any cause. The time to this event is the time from randomization to death, loss to follow-up or the end of the study, whichever comes first. Patients alive at the time of last observation are considered censored. Overall survival will be compared between the vaccine and no-vaccine arms combined from time of randomization. A secondary pairwise analysis will compare overall survival between the vaccine arm, lenalidomide/GM-CSF arm and lenalidomide alone arm.

3.4.Primary Immunologic Endpoints

3.4.1. Myeloma-Reactive T-cells

The effect of DC/Myeloma Fusion Vaccine/GM-CSF with lenalidomide maintenance; lenalidomide maintenance; and lenalidomide maintenance with GM-CSF on levels of circulating myeloma reactive T cells will be quantified. Myeloma-reactive T cells, as defined by the percentage of circulating CD4+ and CD8+ T cells that express IFN γ following ex vivo exposure to autologous tumor lysate will be assessed in both treatment arms. The percent of patients achieving at least a 2.4-fold increase in tumor reactive T cells will be compared between the vaccine and each of the no-vaccine arms separately.

3.5.Secondary Immunologic Endpoints:

3.5.1. Myeloma-Reactive T-cells

The percent of patients achieving at least a 10-fold increase in tumor-reactive T cells will be compared between the vaccine and no-vaccine arms separately as described above.

3.5.2. Assessment of antigen specific reactivity:

The effect of DC/Myeloma Fusion Vaccine/GM-CSF + lenalidomide maintenance; lenalidomide maintenance; and lenalidomide maintenance + GM-CSF on immunologic responses directed against previously identified myeloma antigens will be assessed. In HLA-A2.1 patients, circulating CD8+ T cells binding tetramer constructed from MUC1, SOX2, NY-ESO, and XBP-1 will be quantified. At serial time points PBMC and BMMC will undergo dual staining for CD8 and the appropriate tetramer, and analyzed by multichannel flow cytometry. When cell yields allow, functional characteristics of the tetramer positive population will be determined by measuring IFN γ , granzyme B, IL-4, and IL-10 expression in the tetramer positive cells by intracellular staining.

3.5.3. Quantification of T-cell subsets:

Quantification of regulatory T cells and activated memory effector cells will be performed at the serial time points (Appendix C). T cell subsets will be quantified with respect to the relative proportion of CD4 and CD8+ T cells and the presence of naïve, effector memory, and central memory cells will be determined by staining for CD45RA, and CD45RO, CD27, and CD62L by multichannel flow cytometry. Regulatory T cells will be quantified as CD4/CD25/FOXP3 cells.

T cells expressing PD-1 will be quantified at the time points indicated above. Quantification of granulocytic and monocytic myeloid derived suppressor cells will be performed from peripheral blood and bone marrow samples undergoing multichannel flow cytometry for CD11b, HLA-DR, CD14, CD15 and CD33.

3.5.4. Assessment of effect on NK cell populations:

The effect of lenalidomide alone and in combination with DC/MM vaccination on the phenotype and function of circulating NK cell populations will be assessed as detailed in Appendix C. The expression of a panel of NK receptors including activating receptors (NKp30, NKp44, NKp46, NKG2D, NKG2C, DNAM1), inhibitory receptors (NKG2A and KIRs) and markers of NK function (CD16, CD25, CD132) will be assessed. The effector function of NK cells against MHC class I-deficient leukemia targets and primary MM cells (where available) from the recipient will be assessed by intracellular cytokine assay for interferon-gamma (IFN- γ) production, and cytotoxicity, including CD107a degranulation and 51chromium release assay. The expression of receptors important in the regulation of human NK cell trafficking and homing in vivo, including CD62L, CXCR4, CXCR3, CCR6 and CCR7 will be measured. The effect of treatment on NK function will be assessed by examining expression of the canonical transcription factors eomesodermin (Eomes) and T-bet prior to and serially following initiation of therapy. Surface expression of CD57 and KLRG1, both markers of mature NK cells, will be evaluated.

3.5.5. Assessment of humoral immune response:

Serological Analysis of Recombinant cDNA Expression Library (SEREX) will be performed on serum samples to assess humoral response targeting myeloma antigens in patients treated with lenalidomide maintenance alone or in conjunction with DC/MM fusion vaccination as detailed in Appendix C. Peripheral blood will be transported for a cell based assay to assess the kinetics, magnitude and specificity of vaccine-induced myeloma specific plasmablast responses, as detailed in Appendix C.

CHAPTER 4

4. PATIENT REGISTRATION, ENROLLMENT, AND EVALUATION

4.1. Enrollment Procedures

Patients are enrolled in to the Segment A prior to initial systemic therapy (≤ 1 cycle) and first autologous transplant. At the time of enrollment, an authorized user at the clinical center completes the demographics and primary eligibility form which includes questions confirming that the patient signed the consent and meets the eligibility criteria for study entry. Patients must be successfully enrolled in Segment A prior the collection of tumor cells.

Once the patient is enrolled in the study (Eligibility Form, Segment A competed):

1. A visit schedule based on enrollment date will be available for printing
2. Following enrollment and tumor cell collection the patient should proceed with initial systemic therapy and autologous transplant. Patients should be transplanted within one year of enrollment into Segment A.
3. Patient will be seen 14 days or less prior to randomization for eligibility assessments prior to randomization/enrollment in Segment B. The pre-randomization disease response assessment must be performed ≤ 10 days prior to randomization. Randomization should occur between day 53 and day 80 after the autologous HCT. An authorized user at the clinical center completes a checklist confirming that the patient is eligible to proceed to the next stage of treatment on the study.
4. If the patient is eligible, they will be enrolled in Segment B and randomized to their treatment assignment and the treatment plan is continued.
5. A visit schedule based on the date of randomization will be available for printing.

4.2. Study Monitoring

4.2.1. Follow - Up Schedule

The follow up schedule for study visits after Segment A enrollment and subsequently after randomization are outlined below in tables 4.2.1A and 4.2.1B, respectively. The follow up schedule is the intended schedule; however, the date of the follow up evaluations may vary due to delays in treatment or treatment interruption related to Adverse Events and toxicities. The visit window for the scheduled evaluations is included in each table. Patients will be followed for three years post maintenance initiation.

**4.2.1A: FOLLOW UP SCHEDULE
AFTER ENROLLMENT (SEGMENT A)**

Study Visit	Target Day
Pre- Enrollment Screening	≤ 8 weeks prior to enrollment
Tumor cell collection	7 days post enrollment ± 7 days ¹
Pre- Randomization Screening	≤ 14 days prior to randomization
Randomization	Day 53- Day 80 post transplant ²
Dendritic Cell Collection ³	≤ 14 days after randomization and prior to Day 85 post transplant
Pre- Maintenance Screening	≤ 14 days prior to maintenance initiation
Maintenance Start	90-100 days post transplant

¹ Tumor cell collection by bone marrow aspirate can occur the same day of enrollment and up to 14 days after enrollment.

² Randomization must also occur within 10 days of a disease response assessment.

³ Leukapheresis for dendritic cell collection is only required for patients randomized to the vaccine/GM-CSF/Lenalidomide arm.

**4.2.1B: FOLLOW UP SCHEDULE
POST RANDOMIZATION TO MAINTENANCE**

Study Visit	Target Day Post Maintenance (± 7 days) ¹
Baseline	≤ 14 days prior to randomization
Prior to initiation of maintenance	≤ 14 days prior to maintenance
Start of Maintenance ² Cycle 1	Day 1
Cycle 2	29 days
Cycle 2 + 7 days	35 days
Cycle 3	57 days
Cycle 4	85 days
Cycle 4 + 7 days	91 days
Cycle 6	141 days
Cycle 9 (Start)	225 days
Cycle 9 (End)	253 days
Cycle 12	309 days
Cycle 15	393 days
Cycle 18	477 days
Cycle 21	561 days
Cycle 24	645 days

¹ After 9 cycles of maintenance therapy, patients can be seen ± 14 days from the target date.

**4.2.1C: FOLLOW UP SCHEDULE
ENDPOINT ASSESSMENTS¹**

Endpoints	Days From Transplant
6 months post transplant (6T)	Day 180 \pm 14 days
12 months post transplant (12T) ²	Day 365 \pm 14 days
24 months post transplant (24T) ³	Day 730 \pm 14 days

¹Endpoint assessments will occur at 6 months posttransplant and yearly post-transplant for 2 years or until disease progression. These assessments may be conducted at the required visits outlined in table 4.2.1B if they fall within 14 days of the target date for the post randomization visit.

² The 6 month posttransplant visit should correspond to the start of the 4th cycle of maintenance unless the patient has been delayed for any reason.

The 12 month posttransplant visit should correspond to the start of the 11th cycle of maintenance unless the patient has been delayed for any reason.

³The 24 month post transplant visit should correspond to the start of the 24th cycle of maintenance unless the patient has been delayed for any reason.

4.3.Patient Evaluations**4.3.1. Pre- Enrollment Segment A Evaluations and Requirements**

The following evaluations must be determined \leq 8 weeks prior enrollment on BMT CTN 1401. See table 4.3.1A.

1. History, physical examination, height and weight
2. Karnofsky Performance Score
3. CBC with differential, platelet count
4. Liver Functions and blood chemistries: Serum creatinine, total bilirubin, alkaline phosphatase, AST, ALT
5. Evaluation of creatinine clearance
6. Laboratory Disease Evaluation:
 - a. Quantitative serum immunoglobulin levels.
 - b. Serum protein electrophoresis (SPEP).
 - c. 24 hour urine collection to determine protein excretion, urine electrophoresis (UPEP).
 - d. Immunofixation of urine and serum protein regardless of SPEP and UPEP results.
 - e. Serum free light chain ratios (FLC)

7. Bone Marrow Evaluation: unilateral bone marrow biopsy and aspirate are required to assess plasma cell involvement in the marrow as per eligibility criteria outlined in section 2.3.1
8. Skeletal bone survey to include cranium, axial skeleton and proximal long bones or other imaging study to evaluate for bone involvement.

4.3.2. Tumor Cell Collection Evaluations and Requirements at Enrollment

1. Tumor Cell Collection¹
 - a. Bone marrow aspiration (30mL) is required for vaccine production and eligibility validation.
 - b. Peripheral Blood (50mL) collection
2. Protocol-required Samples for Correlative Studies (Table 4.3B)²
 - a. Bone marrow aspirate (4mL) for immunologic endpoints (see Appendix C for details)
 - b. Peripheral Blood (40mL) collection for immunologic endpoints (see Appendix C for details)
3. Optional Samples for Future Research²
 - a. Bone marrow aspirate (3mL) (see Appendix C for details)
 - b. Peripheral Blood (6mL) collection (see Appendix C for details)

4.3.3. Pre- Randomization Evaluations and Requirements

The following evaluations to determine eligibility for randomization must be performed ≤ 14 days prior to randomization unless otherwise noted below.

1. History and physical examination
2. Assessment for toxicities
3. Laboratory Disease Evaluation (≤ 10 days from randomization):
 - a. Quantitative serum immunoglobulin levels.
 - b. Serum protein electrophoresis (SPEP).
 - c. 24 hour urine collection to determine protein excretion, urine electrophoresis (UPEP).
 - d. Immunofixation of urine and serum protein regardless of SPEP and UPEP results.
 - e. Serum free light chain ratios (FLC)
4. Bone Marrow Evaluation: unilateral bone marrow biopsy and aspirate are required only to confirm CR in patients unless required for BMT CTN 1401 correlative studies.

¹ Must be collected ≤ 14 days after enrollment

² Must be collected ≤ 14 days after enrollment

5. Protocol-required Samples for Correlative Studies (Table 4.3B)³
 - a. Bone marrow aspirate (10mL) for immunologic endpoints (see Appendix C for details)
 - b. Peripheral Blood collection (30mL) for immunologic endpoints (see Appendix C for details)
6. Optional Samples for Future Research
 - a. Bone marrow aspirate (3mL) (see Appendix C for details)
 - b. Peripheral Blood (6mL) collection (see Appendix C for details)
7. Optional Samples for Ancillary Research: To be collected only on patients providing consent for ancillary research per Appendix J and/or K (see Appendix C for details)
 - a. Bone marrow aspirate (4mL) - additional sample volume collected at the time of required sample for immunologic endpoints
 - b. Four (4) Paraffin Embedded Slides from Core Bone Marrow Biopsy required to confirm CR in some study patients prior to randomization
8. Skeletal bone survey to include cranium, axial skeleton and proximal long bones or other imaging study to evaluate for bone involvement.
9. CBC with differential, platelet count
10. Liver Functions and blood chemistries: Serum creatinine, total bilirubin, alkaline phosphatase, AST, ALT
11. Evaluation of creatinine clearance
12. Pregnancy test, serum HCG (sensitivity of at least 50 mIU/mL) for FCBP.

4.3.4. Pre- Maintenance Evaluations and Requirements

The following evaluations to determine eligibility for maintenance initiation must be performed \leq 14 days prior to the initiation of lenalidomide maintenance therapy unless otherwise noted below.

1. History and Physical Exam
2. Assessment for toxicities
3. CBC with differential, platelet count
4. Liver Functions and blood chemistries: Serum creatinine, total bilirubin, alkaline phosphatase, AST, ALT
5. Required evaluation prior Lenalidomide: Pregnancy test, serum HCG (sensitivity of at least 50 mIU/mL) for FCBP. Women are required to have 2 pregnancy tests prior to the initiation of lenalidomide as follows (Appendix D)

³ Should be collected \pm 14 days from date of randomization however, randomization samples are preferred to be collected after the patient is randomized. If samples(s) need to be collected prior to randomization, contact the protocol coordinator.

- a. The first is required within 10 to 14 days prior to prescribing lenalidomide,
- b. The second is required within 24 hours of prescribing lenalidomide.

4.3.5. Evaluations during Maintenance

1. The following evaluations will be done monthly for the first 4 cycles and then at cycles 6, 9, 12, 15, 18, 21, 24:
 - a. History and physical exam
 - b. CBC with differential, platelet count⁴
 - c. Liver Functions and blood chemistries: Serum creatinine, total bilirubin, alkaline phosphatase, AST, ALT
2. Assessment for Toxicities monthly for the first 4 cycles and then at cycles 6, 9, 15, 21, and 24⁵.
3. Required evaluation for Lenalidomide: Pregnancy test, serum HCG (sensitivity of at least 50 mIU/mL) for female of childbearing potential (FCBP):
 - a. FCBP with regular or no menstruation must have a pregnancy test weekly for the first 28 days and then every 28 days while on therapy (including breaks in therapy); at discontinuation of lenalidomide and at Day 28 post the last dose of lenalidomide. Females with irregular menstruation must have a pregnancy test weekly for the first 28 days and then every 14 days while on therapy (including breaks in therapy), at discontinuation of lenalidomide and at Day 14 and Day 28 post the last dose of lenalidomide (Appendix D).
 - b. The subject must follow the requirements of the Revlimid REMS® program of the Celgene Corporation. This program provides education and counseling on the risks of fetal exposure, blood clots and reduced blood counts. The patient will be required to receive counseling every 28 days during treatment with lenalidomide, follow the pregnancy testing and birth control requirements of the program that are appropriate in order to take the telephone surveys regarding compliance with the program.
4. Laboratory Disease Evaluation at prior to cycles 3, and 6, and every 3 cycles thereafter. Disease evaluations will also be collected at 6 months, 1 and 2 years post-transplant:
 - a. Quantitative serum immunoglobulin levels.
 - b. Serum protein electrophoresis (SPEP).
 - c. 24 hour urine collection to determine protein excretion, urine electrophoresis (UPEP).
 - d. Immunofixation of urine and serum protein regardless of SPEP and UPEP results.
 - e. Serum free light chain ratios (FLC)

⁴ Patients receiving lenalidomide should receive monthly CBCs per the requirements of the RevlimidREMs program.

⁵ Patients experiencing adverse events outside of the toxicity assessment timelines are still required to adhere to the adverse event reporting requirements in section 4.4.4 and Appendix I.

5. Bone Marrow Evaluation: unilateral bone marrow biopsy and aspirate are required only to confirm CR in patients
6. Skeletal bone survey to include cranium, axial skeleton and proximal long bones or other imaging study to evaluate for bone involvement at 1 and 2 years post transplant.
7. Samples for Correlative Studies (Table 4.3B)
 - a. Bone marrow aspirate (10mL) just prior to the start of cycle 9 of maintenance for immunologic endpoints (see Appendix C for details).
 - b. Peripheral Blood collection prior to cycles 1, 2, 3, 4, and 9 and at day 7 of cycles 2 and 4 for immunologic endpoints (see Appendix C for details).
8. Optional Samples for Future Research
 - a. Bone marrow aspirate (3mL) collection just prior to the start of cycle 9 of maintenance (see Appendix C for details)
 - b. Peripheral Blood (6mL) collection just prior to the start of cycle 9 of maintenance (see Appendix C for details)
9. Optional Samples for Ancillary Research: To be collected only on patients providing consent for ancillary research per Appendix K (see Appendix C for details)
 - a. Bone marrow aspirate (4mL) (see Appendix C for details) – additional volume collected just prior to the start of cycle 9 of maintenance along with required research sample being collected for the primary trial, and a (5mL) marrow aspirate sample collected at time of disease progression determination.
 - b. Peripheral Blood Collection: additional volume of blood collected prior to cycles 1, 2, and 4 (10mL) along with the required research samples being collected for the primary trial, and (40mL) blood collected at the time of disease progression determination.

TABLE 4.3A: PATIENT CLINICAL ASSESSMENTS

Study Assessments	Pre-Enrollment	Pre-Randomization	Pre maintenance	Cycles of maintenance										Post-transplant visits		
			1	2	3	4	6	9	12	15	18	21	24	6T	12T	24T
History, physical exam, weight and height ¹	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Karnofsky performance score	X															
CBC, differential, platelet count, and blood chemistries ²	X	X	X	X	X	X	X	X	X	X	X	X	X			
Evaluation of Creatinine Clearance	X	X														
Pregnancy test ³		X	X	X	X	X	X	X	X	X	X	X	X			
Quantitative serum immunoglobulins	X	X			X		X	X	X	X	X	X	X	X	X	X
SPEP and immunofixation	X	X			X		X	X	X	X	X	X	X	X	X	X
24 Hour Urine for UPEP, protein excretion and immunofixation	X	X			X		X	X	X	X	X	X	X	X	X	X
Serum free light chain ratio	X	X			X		X	X	X	X	X	X	X	X	X	X
Skeletal Survey or other imaging study	X	X													X	X
Bone marrow aspirate and biopsy	X ⁴	X ⁵			X ₅		X ⁵	X ⁵	X ⁵	X ⁵	X ⁵	X ⁵	X ⁵	X ⁵	X ⁵	X ⁵
Toxicity assessment ⁶		X	X	X	X	X	X	X		X		X	X			

¹Height and Weight are only required at pre-enrollment and pre-randomization visits.

²Blood chemistries include: serum creatinine, bilirubin, alkaline phosphatase, AST and ALT as required per protocol.

³For female of childbearing potential: A female of childbearing potential (FCBP) is a sexually mature female who: 1) has not undergone a hysterectomy or bilateral oophorectomy; or 2) has not been naturally postmenopausal for at least 24 consecutive months (i.e., has had menses at any time in the preceding 24 consecutive months).

⁴Bone Marrow aspirate and biopsy required within 60 days of enrollment; must be repeated if patient receives additional therapy between initial pre-treatment BM and enrollment

⁵Bone Marrow aspirate and biopsy required only to confirm CR in patients

⁶The toxicity assessment will include a review of **all** toxicities and appropriate lab evaluations experienced **during the entire assessment period** and the **highest grade** for each toxicity during the assessment period will be recorded on the Toxicity form in AdvantageEDC using CTCAE version 4.

TABLE 4.3B: SAMPLE COLLECTION TIME POINTS

Sample Collection	Tumor Cell Collection (At Enrollment) ⁴	Pre-Randomization ⁶	Cycles of maintenance								Progression ¹⁰
			1 ⁷	2 ⁷	2 + 7d ⁸	3 ⁷	4 ⁷	4 + 7d ⁸		9 ⁷ (prior)	
Bone Marrow Aspirate											
Bone marrow aspirate for tumor collection (30mL)	X ¹										
MRD Assessment (6 mL) ²		X								X	
Myeloma Reactive T Cells (4 mL)	X ¹	X								X	
T Cell/Genomic Signature Study (4mL unless otherwise noted)		X								X	X (5mL)
VCAN Assessment ⁹		X									
Optional Repository Sample for Future Research (3mL)	X ¹	X								X	
Total Maximum Marrow Volume (mL)⁵	37	17								17	5
Peripheral Blood											
Plasma for Vaccine Production (50mL)	X ⁵										
Plasmablast Responses (30mL)		X		X ³	X ³		X ³	X			
Myeloma Reactive T Cells (40mL)	X ⁵		X	X		X	X			X	
NK cell reconstitution (30mL)			X		X	X		X		X	
T Cell/Genomic Signature Study (10mL unless otherwise noted)			X	X			X				X (40mL)
Optional Repository Sample for Future Research (6mL)	X ⁵	X								X	
Total Maximum Blood Volume (mL)	96⁵	36	80	80³	60³	70	80³	60		76	40

¹ Bone marrow aspirate for collection of tumor cells should be done with 1-2 pull(s) per site until the total volume is achieved. The first 30mL will be dedicated for vaccine production, which will be cryopreserved at the transplant center. The remainder volume (7mL) will be aliquoted into two samples, which will be shipped to the BMT CTN repository and to BIDMC.

² Minimal residual disease (MRD) assessment will be done at the center and centrally at RPCI, the MRD sample will be aliquot in two 3 mL samples, one will be sent to RPCI and the other for analysis at the transplant center. Patients with overt evidence of disease progression are not required to submit sample for MRD assessment.

³ Assessment for plasmablast only applies to patients who were randomized to the vaccine arm.

⁴ The Tumor Cell Collection Visit should occur ≤ 14 days post enrollment in Segment A.

⁵ Blood collection at study entry can be collected over a 10 day period to avoid collection of excessive volumes in a single day.

⁶The Pre-Randomization collection should be collected ± 14 days from date of randomization however, randomization samples are preferred to be collected after a patient is randomized. If samples(s) need to be collected prior to randomization, contact the protocol coordinator.

⁷Samples should be collected ≤ 3 days from target date (If target date is a Friday, Saturday or Sunday, collect the previous Thursday)

⁸Samples should be collected -2 days to +1 day from target (If target date is a Friday or, Saturday, collect the previous Thursday. If Sunday, collect Monday.)

⁹No additional sample is required. Four (4) Paraffin embedded slides prepared at the clinical site from the core biopsy required for clinical assessment prior to randomization. ¹⁰Collected at the time of confirmation of disease progression per Chapter 3.

4.4.Data Reporting

4.4.1. Criteria for Forms Submission

Forms that are not entered into AdvantageEDCSM within the specified time will be considered delinquent. A missing form will continue to be requested either until the form is entered into the AdvantageEDCSM and integrated into the Data and Coordinating Center's (DCC) master database, or until an exception is granted and entered into the Missing Form Exception File.

4.4.2. Reporting Patient Deaths

Recipient death information must be entered into AdvantageEDCSM within 24 hours of knowledge of the patient's death. If the cause of death is unknown at that time, it need not be recorded at that time. However, once the cause of death is determined, the form must be updated in AdvantageEDCSM.

4.4.3. Center for International Blood and Marrow Transplant Research (CIBMTR) Data Reporting

Centers participating in BMT CTN trials must register pre- and post-transplant outcomes on all consecutive hematopoietic stem cell transplants done at their institution during their time of participation to the Center for International Blood and Marrow Transplant Research (CIBMTR). Registration is done using procedures and forms of the Stem Cell Transplant Outcomes Database (SCTOD). (Note: Federal legislation requires submission of these forms for all US allotransplant recipients.) Enrollment in BMT CTN 1401 must be indicated on the SCTOD pre-transplant registration form. Additionally, CIBMTR pre- and post- transplant Comprehensive Report Forms must also be submitted for all patients enrolled on this trial. CIBMTR forms will be submitted directly to the CIBMTR at the times specified on the Form Submission Schedule.

4.4.4. Adverse Event Reporting

Reporting of adverse events on the BMT CTN 1401 protocol has unique requirements due to the addition of lenalidomide as part of the protocol. Adverse event reporting requirements are summarized below and further described in Appendix I.

4.4.4.1.Definitions

Adverse Event: An adverse event (AE) is any undesirable sign, symptom or medical condition or experience that develops or worsens in severity after starting the first dose of study treatment or any procedure specified in the protocol, even if the event is not considered to be related to the study.

Expectedness: An adverse event can be Expected or Unexpected

- **Expected adverse events** are those that have been previously identified as resulting from administration of the agent. For the purposes of this study, an adverse event is considered expected when it appears in the current adverse event list, the Investigator's Brochure, the package insert or is included in the informed consent document as a potential risk.
- **Unexpected adverse events** are those that vary in nature, intensity or frequency from information in the current adverse event list, the Investigator's Brochure, the package insert, or when it is not included in the informed consent document as a potential risk.

4.4.4.2. BMT CTN Adverse Event Reporting Guidelines

It is BMT CTN policy that AEs must be reported even if the investigator is unsure whether a relationship exists between the adverse event and the use of the study treatment. Reporting of AEs for BMT CTN 1401 will be consistent with the BMT CTN Administrative Manual of Procedures. Additional requirements specific to this protocol are outlined below and in Appendix I.

In BMT CTN studies, expected adverse events are reported via the web-based electronic data capture system, AdvantageEDC. Events are captured on calendar-driven case report forms (e.g., Toxicity) or event-driven case report forms (e.g., Progression, and Death).

From the time of enrollment until randomization, only deaths, secondary primary malignancies and progressions are required to be reported. From time of randomization until 3 years post maintenance initiation, unexpected, grades 3-5 AEs, irrespective of the attribution of the event to the study drug /procedure/treatment, will be reported through the expedited AE reporting system via AdvantageEDC, and will be graded according to the NCI Common Terminology Criteria for Adverse Events (CTCAE), Version 4.0. The BMT CTN 1401 protocol has two distinct interventions: transplant and maintenance therapy. Determination of expectedness for events occurring post-transplant and/or post-maintenance therapy will be at the discretion of the investigator as described in Appendix I. **Unexpected, grades 4-5 AEs must be reported within 24 hours of knowledge of the event. Unexpected, grade 3 AEs must be reported within three business days of knowledge of the event.** The NHLBI Data and Safety Monitoring Board will receive summary reports of all adverse experiences at least twice yearly.

Since this study is under an FDA Investigational New Drug, all suspected and unexpected fatal or life-threatening adverse events are reported to the FDA within seven calendar days after receipt of the information, following FDA guidelines. All suspected and unexpected Grade 3 serious adverse events are reported to the FDA via a written report within fifteen days of receipt of the information (21 CFR 312.32). If the Medical Monitor assesses the event to be unrelated to the study, then the event will not require expedited reporting but will be included in a summary report issued annually. All expected adverse events (i.e., those listed in the informed consent, product inserts, or study materials) not covered under the above requirements need not be reported via the AE Form. They will be captured on other CRFs. Although death and graft failures are not considered unexpected post-transplant events, they are reported to the FDA via annual reports. Safety reporting to the FDA will be done by the BMT CTN based upon date of initial entry on the Adverse Event Forms and requires no additional action from the participating site.

All second primary malignancies (SPM), excluding non-melanoma skin cancers, experienced by patients enrolled on the study will be reported using the expedited AE reporting system in AdvantageEDC and must be reported within three business days of knowledge of the event. The Event Description entered in AdvantageEDC should include histologic type.

4.4.4.3. Additional Adverse Event Reporting

Celgene Corporation (Celgene) is supplying lenalidomide for this study and a description of additional reporting requirement for this study is detailed in **Appendix I**. The additional adverse

event reporting period for lenalidomide begins with the first dose and continues until 28 days after permanent discontinuation of lenalidomide.

4.4.4.4. Adverse Event Reporting Following Progression

If a patient meets the protocol defined definition of progression (Chapter 3), Unexpected Grade 3-5 Adverse Events and events listed in Appendix I are no longer required to be reported on the Adverse Event Form once the patient is more than 28 days from their last dose of lenalidomide. However, SPMs should continue to be reported within three business days of the knowledge of the event through 3 years post-maintenance initiation.

4.4.4.5. Adverse Event Reporting Following an SPM

Adverse Event reporting following an SPM is dependent on the treatment received for the reported SPM.

- If a patient experiences an SPM resulting in permanent discontinuation of lenalidomide and initiation of non-protocol systemic therapy, Unexpected Grade 3-5 Adverse Events and events listed in Appendix I are no longer required to be reported on the Adverse Event Form once the patient is more than 28 days from their last dose of lenalidomide.
- If a patient experiences an SPM that does *not* result in permanent discontinuation of lenalidomide, Adverse Events will continue to be reported as per section 4.2.3 and appendix I of the protocol.
- Requests to discontinue Adverse Event Reporting for events that do not meet the criteria above will be considered on a case by case basis.

4.4.5. Pregnancy Reporting

Pregnancies, suspected exposure of a pregnant woman (including female partner of a male patient and exposure to lenalidomide while dispensing) and suspected pregnancies (including a positive pregnancy test regardless of age or disease state) of a female subject occurring while the subject is on lenalidomide, or within 28 days of the subject's last dose of lenalidomide, are considered immediately reportable events. Lenalidomide is to be discontinued immediately. The pregnancy, suspected pregnancy, or positive pregnancy test must be reported through the expedited AE reporting system in AdvantageEDC. The female subject should be referred to an obstetrician-gynecologist, preferably one experienced in reproductive toxicity for further evaluation and counseling.

The Investigator will follow the female subject until completion of the pregnancy, and must notify Celgene Drug Safety immediately about the outcome of the pregnancy (either normal or abnormal outcome) using the expedited AE reporting system in AdvantageEDC. If the outcome of the pregnancy was abnormal (e.g., spontaneous or therapeutic abortion), the Investigator should report the abnormal outcome as an AE.

All neonatal deaths that occur within 28 days of birth should be reported, without regard to causality through the expedited AE reporting system in AdvantageEDC. In addition, any infant death after 28 days that the Investigator suspects is related to the inutero exposure to lenalidomide

should also be reported through the expedited AE reporting system within 24 hours of the Investigator's knowledge of the event

Male Subjects

If a female partner of a male subject taking investigational product becomes pregnant, the male subject taking lenalidomide should notify the Investigator, and the pregnant female partner should be advised to call her healthcare provider immediately.

4.4.6. Celgene Drug Safety Contact Information:

Celgene Corporation
Global Drug Safety and Risk Management
Connell Corporate Park
300 Connell Dr. Suite 6000
Berkeley Heights, NJ 07922
Fax: (908) 673-9115
E-mail: drugsafety@celgene.com

4.4.7. Reporting of other Special Events of Interest

There are other special events of interest to Celgene that require reporting through the AE expedited reporting system in AdvantageEDC. If any of the following events occur, they are required to be reported within 3 business days of the site's knowledge of the event:

- overdose, abuse and misuse that meets one of the criteria of a serious adverse event
- medication errors that meets one of the criteria of a serious adverse event
- occupational exposure of any kind
- public health emergency
- Celgene product technical complaint that leads to an adverse event

4.5. Ancillary Correlative Laboratory Studies Procedures

Selected study patients will be presented with the opportunity to participate in the ancillary studies outlined in Appendix J and K. This section describes procedures associated with the ancillary studies including data entry and obtaining informed consent. Additional details including eligibility requirements for each ancillary study can be found in the protocol appendices.

4.5.1. Obtaining Informed Consent

The Optional Research Sample Consent form should be presented to eligible patients \leq 30 days prior to randomization on BMT CTN 1401.

The following 3 documents are provided for documentation of informed consent to participate on the ancillary studies in appendix J and K:

- **Informed Consent Form to Participate in Optional Ancillary Correlative Laboratory Research:** The purpose of this document is to obtain consent to participate in the Optional Ancillary Research for BMT CTN 1401. This consent provides an overview of the required components required to obtain informed consent. However, it must be presented in conjunction with at least one of the two supplementary consent documents.
- **Supplementary Consent Document for the VCAN Study:** The purpose of this document is to be presented alongside the Informed Consent Form to Participate in Optional Ancillary Research. This consent document should be presented to patients eligible for the VCAN study per appendix J. Patients must sign this Supplementary Consent Document along with the Informed Consent Form to Participate in Optional Ancillary Research in order to move forward with providing the required sample for the VCAN Study.
- **Supplementary Consent Document for T Cell/Genomic Signature Study:** The purpose of this document is to be presented alongside the Informed Consent Form to Participate in Optional Ancillary Research. This consent document should be presented to patients eligible for the T Cell/Genomic Signature Study per appendix K. Patients must sign this Supplementary Consent Document along with the Informed Consent Form to Participate in Optional Ancillary Research in order to move forward with providing the required samples for the T Cell/Genomic Signature Study.

4.5.2. Data Entry Requirements

Consent and Eligibility for ancillary research participation will be documented on the BMT CTN 1401 Segment B Enrollment Form.

Sample collection and shipment will be documented in GlobalTrace and on the Specimen Acquisition Form in AdvantageEDC. Further information regarding sample collection, shipment, and documentation can be found in the Research Sample Information Guide and the BMT CTN 1401 Forms Guide.

Additional information on the data entry and laboratory procedures are outlined in appendix J and K.

CHAPTER 5

5. STATISTICAL CONSIDERATIONS

5.1. Study Design

The study is designed as a Phase II randomized, open label, multicenter trial to evaluate the use of vaccination with DC/myeloma fusions/GM-CSF plus lenalidomide maintenance therapy compared to lenalidomide maintenance therapy with and without GM-CSF following autologous transplant as part of upfront treatment for patients diagnosed with multiple myeloma. Patients meeting initial eligibility will be enrolled for the purpose of harvesting tumor cells. Randomization will then occur approximately 2 months after autologous transplant and within 10 days of disease assessment. The target enrollment is 132 patients randomized; assuming that 30% will drop out between tumor collection and randomization, the study will target 188 patients for initial enrollment. As the impact of GM-CSF without vaccine on clinical outcomes is expected to be minimal, analysis of clinical outcomes will focus on comparing the vaccine arm vs. the combined no-vaccine arms. This will help maintain power for these primary analyses. Further randomization of the no-vaccine arms into maintenance vs. maintenance +GM-CSF is done to isolate the effect of the GM-CSF adjuvant on immunological outcomes and address whether any observed effect of vaccine is truly related to the vaccine.

5.1.1. Accrual

It is estimated that 36 months of accrual will be necessary to enroll the targeted sample size. Accrual will be reported by race, ethnicity, gender, and age.

5.1.2. Randomization

Randomization will occur approximately 2 months after autologous transplant and within 10 days of disease assessment. Randomization will be performed in a 2:1:1 ratio using random block sizes for the vaccine/GM-CSF (n=66), lenalidomide/GM-CSF (n=33) and lenalidomide alone (n=33) treatment arms, respectively. Randomization will be stratified according to disease response at time of randomization between sCR/CR and VGPR/PR/Stable disease.

5.1.3. Primary Endpoint

The primary endpoint is the proportion of patients alive and in CR or sCR at one year post transplant, corresponding to approximately 10 months post randomization. The primary analysis will be performed using the intent-to-treat principle so that all randomized patients will be included in the analysis.

5.1.4. Primary Hypothesis

The primary objective of the study is to determine whether vaccination with DC/myeloma fusions/GM-CSF in combination with lenalidomide maintenance therapy increases the proportion of patients alive and in CR or sCR at one year compared to the combined maintenance groups without vaccine. Letting P denote the proportion of patients alive and in CR and sCR at one year, the hypotheses of interest are:

$H_0: P_{\text{VACCINE/GMCSF+MAINT}} = P_{\text{MAINT+GMCSF}}$ vs. $H_a: P_{\text{VACCINE/GMCSF+MAINT}} > P_{\text{MAINT+GMCSF}}$.

5.1.5. Duration of follow-up

All patients will be followed for approximately 3 years post-maintenance initiation for primary and secondary endpoints.

5.2. Sample Size and Power Considerations**5.2.1. Primary Endpoint (Clinical)**

Based on the prior cooperative group studies, we anticipate approximately 40% of patients will achieve CR and sCR at 1 year. In the prior phase II study of patients undergoing post-transplant vaccination in the absence of lenalidomide maintenance, CR was achieved in 29% and 54% of patients at 100 days and 1 year post-transplant respectively. Given that we anticipate that lenalidomide maintenance will enhance vaccine efficacy, we will consider the experimental arm of vaccine/GM-CSF + maintenance promising if the CR rate at 1 year is improved from 40% to 60%. A sample size of 66 patients (132 total) randomized to the vaccine arm and the combined no-vaccine arms will have 85% power to detect this improvement in the CR rate at 1 year from 40% to 60%, using a two-sample comparison of proportions with a one-sided type I error of 10%. Because randomization occurs post collection of tumor sample and post-transplant, we anticipate needing to enroll 188 at the time of initial tumor collection pre-transplant assuming that 30% will drop out between tumor collection and randomization.

Secondary comparisons of CR rate at 1 year post HCT between the vaccine/GM-CSF arm vs. either lenalidomide/GM-CSF or lenalidomide alone separately will have 80% power to detect a 23% increase in the CR rate, using the same one-sided alpha level of 10%. Comparison between the two no-vaccine arms will have 80% power to detect a 26% improvement in the CR rate.

5.2.2. Power Considerations for Primary Immunologic Endpoint

The primary immunologic endpoint will be to quantify the expansion of myeloma reactive T cells as determined by the pretreatment and peak post-treatment levels. The anticipated standard deviation of the log 10 peak change in tumor reactive T cells from baseline is 0.61, based on prior studies. Assuming 15% of patients will not be evaluable for immune reconstitution studies, the expected evaluable sample sizes will be 56 for the vaccine group and 28 for each of the no-vaccine groups. Comparisons between the vaccine group and either of the no-vaccine groups will have 80% power to detect a log 10 fold change of 0.4, corresponding to a 2.5 fold expansion of myeloma specific T cells, using a two-sample t-test with a 5% two-sided significance level. Comparisons between the two no-vaccine groups will have 80% power to detect a log 10 fold change of 0.46, corresponding to a 2.9 fold expansion of myeloma specific T cells. Note that a stricter alpha level is used for immunologic endpoints to reduce the risk of type I errors while still being adequately powered to detect larger quantitative differences likely to translate into clinical improvements.

5.3. Interim Analysis and Stopping Guidelines**5.3.1. Interim Analysis for Efficacy**

There will be no interim analyses for efficacy.

5.3.2. Interim Analysis for Futility

There will be no interim analyses for futility.

5.3.3. Guidelines for Safety Monitoring

Monitoring of a key safety endpoint will be conducted monthly, and if rates significantly exceed pre-set thresholds, the NHLBI will be notified in order that the DSMB can be advised. Policies and composition of the DSMB are described in the BMT CTN's Manual of Procedures. The stopping guideline serves as trigger for consultation with the DSMB for additional review.

The key safety endpoint for this study is treatment limiting toxicity (TLT) within the first month of combined therapy, defined as any grade V event regardless of attribution to combined therapy, grade IV hematologic (lasting greater than 7 days) and grade III-IV non-hematologic toxicity judged to be related to combined therapy. At least three events must be observed in order to trigger review. The rate of TLT within the first month is expected to be no higher than 25%. Each month, the null hypothesis that the 1 month TLT rate is 25% will be tested. This outcome will be monitored using a truncated Sequential Probability Ratio Test (SPRT) for binary data as described below. The SPRT conserves type I error at 5% across all of the monthly examinations.

The SPRT can be represented graphically. At each interim analysis, the total number of patients enrolled and started combined therapy is plotted against the total number of patients who have experienced treatment limiting toxicities. The continuation region of the SPRT is defined by two decision boundaries. Only the upper boundary will be used for monitoring the study to protect against high incidences of TLT. If the graph falls above the upper boundary, the SPRT rejects the null hypothesis, and concludes that the TLT incidence is higher than predicted by the observed number of patients enrolled on study. Otherwise, the SPRT continues until enrollment reaches the target goal. The SPRT for TLT was developed from the following SPRT:

A SPRT contrasting 25% versus 40% 1 month incidence, with nominal type I and II errors of 12% and 15%, respectively, which results in decision boundaries with a common slope of 0.322 and an upper intercept of 2.824.

The actual operating characteristics of this truncated test, shown in Table 5.3.3, were determined in a simulation study that assumed uniform accrual of 66 individuals over a three-year time period.

TABLE 5.3.3: OPERATING CHARACTERISTICS OF SEQUENTIAL TESTING PROCEDURE FOR 1 MONTH TLT FROM A SIMULATION STUDY WITH 10,000 REPLICATIONS

True 1 month Incidence	25%	35%	40%	45%
Probability Reject Null	0.100	0.595	0.836	0.957
Mean Month Stopped	34.8	25.0	18.3	13.5
Mean # Endpoints in 1 month	15.5	15.4	13.1	10.6
Mean # Patients Enrolled	62.1	43.8	32.7	23.7

For example, the testing procedure rejects the null hypothesis in favor of the alternative 10% of the time when the true 1 month TLT rate is 25%, and 84% of the time when the rate is 40%. This corresponds to a type I error rate of $\alpha = 0.1$ and a type II error rate of $\beta = 0.16$. When the true

1 month TLT rate is 40%, on average, the DSMB will be consulted 18 months after opening, when 13 events have been observed in 33 patients.

5.4. Demographic and Baseline Characteristics

Demographics and baseline characteristics will be summarized for all patients. Characteristics to be examined are: age, gender, race/ethnicity, performance status, pre-randomization disease response. Between group comparisons will be performed for continuous variables via a Kruskal-Wallis test and for categorical variables, via the chi-square test.

5.5. Analysis of Primary Clinical Endpoint

The intent-to-treat principle will be followed, so that all patients who are randomized will be included in the analysis. The proportion of patients alive and in CR/sCR at 1 year post transplant will be described in the vaccine and no vaccine groups with 80% confidence intervals and compared between groups using a two-sample Z test comparing binomial proportions. A one-sided significance level of 0.1 will be used for this comparison to indicate whether the vaccine plus maintenance treatment is promising compared to the combined no-vaccine groups in this phase II trial. An 80% two-sided confidence interval (consistent with one-sided significance level of 0.1) will be constructed for the difference in proportions. A secondary analysis stratified on disease response prior to randomization will be conducted using a Cochran-Mantel-Haenszel test, and a stratified odds ratio along with 80% confidence intervals will be estimated. A secondary pairwise analysis of CR rates comparing the vaccine arm, lenalidomide/GM-CSF arm and the lenalidomide alone arm will also be conducted.

5.6. Analysis of Secondary Clinical Endpoints

5.6.1. Response to treatment

The distribution of disease response will be tabulated at each time point. The proportion of patients alive and with a response of VGPR or better will be compared between the vaccine arm with the no-vaccine arms combined at 6 months, 1 year, and 2 years post-transplant using a chi-square test. A secondary pairwise analysis of response to treatment comparing the vaccine arm, lenalidomide/GM-CSF arm and the lenalidomide alone arm will also be conducted. Rates of conversion to CR will also be compared using a chi-square test among the subset of patients who are not in CR at the time of randomization.

5.6.2. Myeloma Progression

Incidence of myeloma progression will be estimated using cumulative incidence function, treating death as a competing risk. Incidence of myeloma progression will be compared between the vaccine arm with the no-vaccine arms combined using Gray's test. A secondary pairwise analysis of myeloma progression comparing the vaccine arm, lenalidomide/GM-CSF arm and the lenalidomide alone arm will also be conducted.

5.6.3. Treatment-related Mortality

Incidence of TRM will be estimated using the cumulative incidence function, treating myeloma progression as a competing risk. Incidence of TRM will be compared between the vaccine arm with the no-vaccine arms combined using Gray's test. A secondary pairwise analysis of TRM

comparing the vaccine arm, lenalidomide/GM-CSF arm and the lenalidomide alone arm will also be conducted.

5.6.4. Progression-free survival

Progression-free survival curves will be estimated using the Kaplan-Meier estimator, and compared between the vaccine arm with the no-vaccine arms combined using the log-rank test. Hazard ratios, along with confidence intervals, will be estimated from a Cox model with treatment group as a covariate. A secondary pairwise analysis of PFS comparing the vaccine arm, lenalidomide/GM-CSF arm and the lenalidomide alone arm will also be conducted.

5.6.5. Overall survival

Overall survival curves will be estimated using the Kaplan-Meier estimator, and compared between the vaccine arm with the no-vaccine arms combined using the log-rank test. Hazard ratios, along with confidence intervals, will be estimated from a Cox model with treatment group as a covariate. A secondary pairwise analysis of overall survival comparing the vaccine arm, lenalidomide/GM-CSF arm and the lenalidomide alone arm will also be conducted.

5.6.6. Incidence of toxicities grade ≥ 3 per CTCAE version 4.0

All Grade ≥ 3 toxicities will be tabulated by grade for each treatment arm, by type of toxicity as well as the peak grade overall. Toxicity frequencies will be described for each time interval as well as cumulative over time.

5.6.7. Incidence of infections

The number of infections and the number of patients experiencing infections will be tabulated by type of infection, severity, and time period after transplant. The cumulative incidence of infections, treating death as a competing event, will be compared between the vaccine arm with the no-vaccine arms combined using Gray's test. A secondary pairwise analysis of the incidence of infections comparing the vaccine arm, lenalidomide/GM-CSF arm and the lenalidomide alone arm will also be conducted.

5.6.8. Minimal Residual Disease assessment

The proportions of patients with MRD present (MRD+) will be described using frequencies at each timepoint and compared between the vaccine arm with the no-vaccine arms combined using the chi-square test. A secondary pairwise analysis of MRD comparing the vaccine arm, lenalidomide/GM-CSF arm and the lenalidomide alone arm will also be conducted.

5.7. Analysis of Immunological Endpoints

5.7.1. Primary immunologic endpoint: Myeloma reactive T-cells

Log transformations will be used to induce normality in quantitative measurements, and if still non-normal, then nonparametric tests will be used. The primary endpoint of log 10 peak change in tumor reactive T cells from baseline will be compared across the 3 groups using analysis of variance, and if significant then two-sample t-tests will be conducted for each pairwise comparison to determine which groups are different from one another. A secondary analysis of peak change in tumor reactive T cells will be done by comparing the proportions of patients who experience >10 -fold increase in IFN γ expression, using a chi-squared test. Myeloma reactive T-cell response

profiles over time will be compared between the treatment groups using linear mixed models for repeated measures data. Peak immune response as well as most recent immune response will be compared between those in CR and those not in CR at each assessment, using a two-sample t-test, to assess the relationship between immune response and clinical response. Peak and most recent immune response will also be considered as time-dependent covariates in a Cox proportional hazards model to assess their impact on progression-free survival.

5.7.2. Secondary immunologic endpoints:

Additional exploratory analyses will be conducted in a similar fashion to examine a number of secondary immunologic endpoints, including: 1) antigen specific reactivity by tetramer analysis, 2) quantification of T-cell subsets and PD-1 expressing lymphocytes by flow cytometry 3) quantification of NK cell populations with inhibitor and activating markers, 4) NK-cell cytotoxic function as measured by IFN γ and CD107a degranulation in response to ex vivo exposure to autologous MM cells, and 5) humoral response against autologous MM cells and myeloma associated antigens such as MUC1, and vaccine-induced myeloma-specific plasmablast responses. In each case, log transformations will be considered to induce normality, and if still non-normal then nonparametric tests will be used. Profiles of these secondary immunologic endpoints will be described at each time point for each group using summary statistics and compared between the groups using mixed models for repeated measures data. The correlation between the immune environment measures and the myeloma specific T-cell response will be assessed and reported using Spearman rank correlation at each time point.

5.8.Endpoint Review Committee

An Endpoint Review Committee of appointed physicians will convene to adjudicate the primary endpoint of complete response as defined by a protocol endpoint review charter. The charter outlines the primary and secondary endpoints to be evaluated and details the endpoints to be adjudicated, process for adjudication, procedures for discrepancy resolution, and recording of results. This review will be done in a blinded manner.

APPENDIX A

CRITERIA FOR SYMPTOMATIC MULTIPLE MYELOMA

APPENDIX A**CRITERIA FOR SYMPTOMATIC MULTIPLE MYELOMA****All three required:**

1. Monoclonal plasma cells in the bone marrow $\geq 10\%$ and/or presence of a biopsy-proven plasmacytoma
2. Monoclonal protein present in the serum and/or urine^a
3. Myeloma defining events (1 or more)
 - a. Myeloma- related organ dysfunction^b:
 - i. Calcium elevation in the blood (serum calcium > 10.5 mg/l or upper limit of normal)
 - ii. Renal insufficiency (serum creatinine > 2 mg/dl)
 - iii. Anemia (hemoglobin < 10 g/dL)
 - iv. Lytic bone lesions or osteoporosis^c
 - b. Any one or more of the following biomarkers of malignancy:
 - i. Clonal bone marrow plasma cell percentage $\geq 60\%$
 - ii. Involved: uninvolved serum free light chain ratio ≥ 100
 - iii. > 1 focal lesion of MRI

*Note: These criteria identify Stage IB and Stages II and III A/B myeloma by Durie/Salmon stage. Stage IA becomes smoldering or indolent myeloma.

- ^a If no monoclonal protein is detected (nonsecretory disease), then $\geq 30\%$ monoclonal bone marrow plasma cells and/or a biopsy-proven plasmacytoma required.
- ^b A variety of other types of end organ dysfunctions can occasionally occur and lead to a need for therapy. Such dysfunction is sufficient to support classification as myeloma if proven to be myeloma related.
- ^c If a solitary (biopsy-proven) plasmacytoma or osteoporosis alone (without fractures) are the sole defining criteria, then $\geq 30\%$ plasma cells are required in the bone marrow.

*Myeloma management guidelines: a consensus report from the Scientific Advisors of the International Myeloma Foundation. The Hematology Journal (2003) 4, 379–398

APPENDIX B

INFORMED CONSENT DOCUMENTS

APPENDIX B-1

INFORMED CONSENT FORM FOR BMT CTN 1401

Informed Consent to Participate in Research

BMT CTN 1401 v4.0

Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions

Your Name: _____

Study Title: Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions

Protocol: BMT CTN 1401 v4.0

Principal Investigator: *Insert local PI information*

Sponsor: The National Institutes of Health (NIH) is sponsoring this study by providing financial support for the coordination of this study through the Blood and Marrow Transplant Clinical Trials Network (BMT CTN).

1. Introduction

We invite you to join this clinical trial, also known as a research study. You are being asked to join because:

- You're ≥ 18 and < 71 years old
- You have multiple myeloma (MM)
- Your doctor thinks an **autologous transplant** is a possible treatment option for you.

The standard of care for patients with MM is an autologous transplant and **maintenance treatment**. Maintenance treatment is chemotherapy after transplant. There's no cure for MM, so patients get maintenance treatment to slow the return of MM after a transplant.

We're doing this study to learn if maintenance treatment works better alone or with a vaccine made from your own blood cells.

This study will take at least 3 years and will include 203 participants. **Your participation will last for a maximum of 4 years from when you are enrolled.**

This consent form will tell you about the purpose of the study, the possible risks and benefits, other options available to you, and your rights as a participant in the study.

Everyone who takes part in research at [*insert facility name*] should know that:

- Being in any research study is voluntary.
- You may or may not benefit from being in the study. Knowledge we gain from this study may benefit others.
- If you join the study, you can quit the study at any time.
- If you decide to quit the study, it will not affect your care at [*insert name of facility or institution*].
- Please ask the study staff questions about anything that you do not understand, or if you would like to have more information.
- You can ask questions now or any time during the study.

- Please take the time you need to talk about the study with your doctor, study staff, and your family and friends. It is your decision to be in the study. If you decide to join, please sign and date the end of the Consent Form.

You and your doctor will discuss other treatment choices if you do not want to participate in this study.

2. Study Background

The National Institutes of Health (NIH), through the Blood and Marrow Transplant Clinical Trials Network (BMT CTN), are providing staff support and money for this research study. The BMT CTN and the NIH will make decisions about how to manage the study.

For this study, you will receive an **autologous transplant** (transplant). An autologous transplant uses blood-forming cells that are collected from your blood stream. After they are collected and frozen, you will get a high dose of chemotherapy. Chemotherapy before a transplant is called the “conditioning regimen”.

The goal of chemotherapy is to kill as many cancer cells in your body as possible. It also destroys most of the normal cells in your bone marrow. To restore your marrow, your frozen blood-forming cells are thawed and transplanted back into your blood stream. The cells find their way into the bone marrow where they start making healthy blood-forming cells.

The standard of care for patients with MM is an autologous transplant and **maintenance treatment**. Maintenance treatment is chemotherapy after transplant. There’s no cure for MM, so patients get maintenance treatment to slow the return of MM after a transplant.

Some patients may receive a **vaccine, and/or GM-CSF (also known as Leukine)** with their maintenance therapy. GM-CSF helps your body make new white blood cells after transplant.

The vaccine is made from dendritic cells and MM tumor cells in a laboratory. We will collect a sample of your cells through your central line (or catheter). This process is called leukapheresis. If the vaccine is made successfully, it will be frozen and stored to give to you later.

We’re doing this study to learn if maintenance treatment works better alone, or with a vaccine made from your own blood cells.

3. Study Purpose

We are inviting you to take part in this study because you have multiple myeloma (MM) and an autologous transplant is a treatment option for you. We are doing this study to learn more about ways to prevent or slow the return of MM after transplant.

We will use 3 treatments to see which one works best to prevent or slow the return of MM. See **Table 1** for information on each treatment group.

Table 1. Study Treatment Groups

Treatment Group A (maintenance treatment with GM-CSF <u>and</u> vaccines)	Treatment Group B (maintenance treatment with GM-CSF)	Treatment Group C (only maintenance treatment)
<ul style="list-style-type: none"> • Melphalan (chemotherapy drug) • Autologous transplant • Lenalidomide (maintenance treatment) • GM-CSF • Vaccine 	<ul style="list-style-type: none"> • Melphalan (chemotherapy drug) • Autologous transplant • Lenalidomide (maintenance treatment) • GM-CSF 	<ul style="list-style-type: none"> • Melphalan (chemotherapy drug) • Autologous transplant • Lenalidomide (maintenance treatment) •

This study will help doctors make the best choice about treatment after transplant for patients with MM. See section **5: Study Tests and Treatments** for information on how your treatment group will be determined and for information about the drugs listed in **Table 1**.

4. Rights to Ask Questions and/or Withdraw

You have the right to ask questions about the study at any time. If you have questions about your rights as a participant or you want to leave the study, please contact:

[insert contact info]

Being in this study is voluntary. You can choose not to be in this study or leave this study at any time. If you choose not to take part or leave this study, it will not affect your regular medical care in any way.

Your study doctor and study staff will be available to answer any questions that you may have about taking part in or leaving this study.

5. Study Treatment and Tests

We will check your health before you start treatment, during your treatment, and for 3 years after you begin your maintenance treatment.

Before You Start Your Treatment

You will need to have several check-ups and tests to see if you can be in the study. These check-ups and tests are part of your regular cancer care and would be done even if you were not part of this study. These tests include:

- Medical history and physical exam
- Blood tests for cell counts
- Liver and kidney function tests
- Tests to measure your disease
- Bone marrow tests
- Skeletal survey
- A pregnancy test (if you are a woman able to have children)

During Your Treatment

Tumor Cell Collection

After you join the study, we will collect about 30mL (approximately 2 tablespoons) of bone marrow and 50mL (approximately 3 tablespoons) of blood. The bone marrow and blood samples will be frozen until it is determined if you will receive the vaccine as part of your treatment. This tumor cell collection is not part of your standard of care treatment

If you are assigned to Treatment Group A after your transplant, your bone marrow will be thawed and used to make your vaccine. If you are assigned to Treatment Group B or C, your

bone marrow sample will not be used in this study. Your doctor may talk to you about other uses for your stored bone marrow samples not related to this study.

We will also talk with you about giving extra blood and marrow samples for future research (see section **17: Blood and Marrow Samples for Future Research**). The extra blood and marrow samples are also completely optional.

Initial Treatment for Multiple Myeloma (MM)

The first step (initial) is standard chemotherapy treatment (chemo). Chemo lowers the number of MM cells in your body. Different chemotherapy drugs can be used to treat MM. Your doctor will decide which drug or combination of drugs is best for you.

You will be asked to sign a separate consent form that will explain the side effects of whichever drug your doctor chooses. This step may last several months.

Blood-Forming Cell Collection

Next, we'll collect blood-forming cells from your blood stream. This is known as apheresis. These cells will be frozen and stored until your transplant day.

Conditioning Regimen Before Transplant

The third step is the conditioning regimen, or the chemotherapy, you will get before your transplant. The conditioning regimen helps the blood-forming cells start to grow and make new cells in your bone marrow (engraft). The regimen includes a chemo drug called **melphalan**, given by intravenous infusion (IV) in your arm.

Your doctor will decide when you are ready to get melphalan. You will start this treatment step no more than 12 months after you start on this study. Melphalan works better for some patients than it does for others. If the drug does not work well for you, you may not be able to have a transplant. Your doctor will talk to you about other treatment options.

Infusion of Blood Forming Cells (Transplant)

On your transplant day (Day 0), your cells will be given to you through your catheter. The cells will travel to your bone marrow where they will start to make healthy, new blood cells (engraft).

Your Treatment Group

Approximately 2 months after transplant, we will check to see if you're healthy enough to start maintenance treatment. If you are found healthy enough, we will randomize you to 1 of the 3 treatment groups. **"Randomize"** means that you will be put in one or another group by chance, just like flipping a coin. We will use a computer program to assign you by chance to Treatment

Group A, B, or C. You won't be able to choose your group. Once you are assigned to a group, you can't change to another group. The study doctor can't change your group either.

Half of the patients on this study (about 66), will be assigned to Treatment Group A. One-fourth of the patients (about 33) will be assigned to Treatment Group B; and one-fourth of patients (about 33) will be assigned to Treatment Group C. **See the section below, Your Maintenance Treatment After Transplant, for a description of the treatments.**

Once all these tests are done, you will be randomized to Treatment Group A, B or C. See **Table 2** for a schedule of these tests.

If you're not healthy enough after transplant, you won't be randomized. If this happens, your Follow up on this study is complete

Your Maintenance Treatment After Transplant

You will start maintenance treatment about 3 months after your transplant.

➤ **Treatment Group A: Lenalidomide, Vaccine, and GM-CSF injections**

If you are assigned to Treatment Group A, you will get: lenalidomide, vaccines, and GM-CSF injections.

We will collect extra blood samples to make the vaccine. We will insert a catheter into a large vein in your neck or chest. The catheter will collect some blood cells and the rest are returned to your body. This process is called leukapheresis. We will use the blood cells and the marrow that was collected and frozen to create your vaccine. The vaccine will be made in a laboratory and frozen until you're ready for maintenance treatment.

You will take lenalidomide as a pill every day for 2 years or until your disease returns. You will receive enough pills for 1 cycle (28 days in each cycle) at a time. It's important to take the pill at the same time every day. Be sure to talk with your doctor to figure out a good time to take the pill every day.

You will also receive 3 doses of the vaccine. This will be given to you on the first day of your 2nd, 3rd, and 4th cycle of lenalidomide. The vaccine is the part of the treatment we are testing in this study. This is also called the research treatment.

With each vaccine, you will also get an injection of GM-CSF in your upper thigh. GM-CSF helps to boost the effect of the vaccine. You will receive the GM-CSF injection on the day you get the vaccine and every day for 3 days after the vaccine. This will either be given to you in the clinic or you will be taught to give the injections to yourself at home.

See **Table 2** for a timeline of the drugs for Treatment Group A.

Table 2: Treatment Group A

Treatment Group A drugs:	Amount:	Timeline for Maintenance Treatment Drugs											
		Cycle: (28 days in each cycle)	1	2	3	4	6	9	12	15	18	21	24
Lenalidomide	1 dose every day for 2 years		X	X	X	X	X	X	X	X	X	X	X
Vaccine	3 doses total			X	X	X							
GM-CSF	12 doses total			X (+3 days)	X (+3 days)	X (+3 days)							

➤ **Treatment Group B: Lenalidomide and GM-CSF**

If you are assigned to Treatment Group B, you will get: lenalidomide and GM-CSF injections.

You will take lenalidomide every day for 2 years or until your disease returns. You will receive the enough pills for 1 cycle (28 days in each cycle) at a time. It's important to take the pill at the same time every day. Be sure to talk with your doctor to figure out a good time to take the pill every day.

You will also get an injection of GM-CSF in upper thigh. You will get the GM-CSF injection every day for 4 days starting the first day of cycles 2, 3 and 4. This will either be given to you in the clinic or you will be taught to give the injections to yourself at home.

See **Table 3** for a timeline of the drugs for Treatment Group B.

Table 3: Treatment Group B

Treatment Group B drugs:	Amount:	Timeline for Maintenance Treatment Drugs											
		Cycle: (28 days in each cycle)	1	2	3	4	6	9	12	15	18	21	24
Lenalidomide	1 dose every day for 2 years		X	X	X	X	X	X	X	X	X	X	X
GM-CSF	12 doses total			X (+ 3 days)	X (+ 3 days)	X (+ 3 days)							

➤ Treatment Group C: Lenalidomide

If you are assigned to Treatment Group C, you will take lenalidomide every day for 2 years or until your disease returns. You will receive enough pills for 1 cycle (28 days in each cycle) at a time. It's important to take the pill at the same time every day. Be sure to talk with your doctor to figure out a good time to take the pill every day.

Maintenance treatment dose (All treatment groups)

How We Will Give You Lenalidomide

Lenalidomide is available only from a certified pharmacy through the Revlimid REMS® program. During maintenance therapy, only a 28-day supply (1 cycle) will be given to you at a time.

If this is the first time you've taken lenalidomide, you'll have to sign-up, or register, for the program. This will include a separate consent process, which calls out the reproductive risks of taking this medicine. You'll have to give your name, address, phone number, birth date, and social security number to sign-up for the program. This information will be provided to Celgene Corporation and Biologics Incorporated to show you're taking part in this study.

After you're registered, you'll be counseled at the site or through the program:

- When you're first given the drug
- At least every 28 days while you're taking it, and

- When you stop taking it

You will be counseled about:

- Side effects
- Not sharing lenalidomide (or other study drugs)
- Risks of exposing a fetus (unborn baby)
- Donating blood
- How to take the pills

We'll give you the, "*Lenalidomide Information Sheet for Patients Enrolled in Clinical Research Studies*" with each new supply of lenalidomide to remind you of these safety issues.

How to Take the Lenalidomide Pills

- Swallow the whole lenalidomide pill with water at the same time each day. Do not break, chew or open the capsules.
- If you miss a dose, take it as soon as you remember on the same day. If you miss taking your dose for the entire day, take your regular dose the next scheduled day (do not take a double dose to make up for the missed dose).
- If you take more than the prescribed dose you should seek emergency medical care if needed and contact study staff right away.
- Women who can become pregnant that care for you should wear gloves whenever they need to touch the lenalidomide pills or bottles.

Checking Your Health

We will watch your health closely during your maintenance treatment, including how well your organs work. We will lower your dose if your organs don't handle the treatment well.

We won't start a new cycle until your organs work well again. If we lower your dose and then your organs start working normally, we may raise your dose again.

We will stop the maintenance treatment if you:

- Have a serious side effect

- Have low blood cell counts
- Are a woman and become pregnant, or there is a chance that you are pregnant
- Go more than 56 days before starting a new maintenance treatment cycle
- Don't follow the study directions, or
- Choose to leave the study.

You will need to visit your clinic for several check-ups and tests during your maintenance treatment. These tests are shown in **Table 4. These assessments are standard of care for patients receiving lenalidomide maintenance therapy.**

Table 4. Timeline of Tests Before and During Your Maintenance Treatment

Tests	Before you're Random ized	Timeline for Maintenance Treatment Tests (CYCLES)											Timeline for Post-Transplant Tests (Months)		
		1	2	3	4	6	9	12	15	18	21	24	6T	12T	24T
Physical exam, height, and weight	X		X	X	X	X	X	X	X	X	X	X	X	X	X
Tests for toxicities, and infections	X	X	X	X	X	X	X		X		X	X			
Blood tests for cell counts, liver and kidney function	X	X	X	X	X	X	X	X	X	X	X	X			
Tests to see how much cancer you still have	X			X		X	X	X	X	X	X	X	X	X	X
Pregnancy test (women able to have children)	X	X	X	X	X	X	X	X	X	X	X	X			

Blood and Bone Marrow Samples

Throughout the study, we will collect blood and marrow samples to check if the treatments are working. These samples are being done as a part of the standard of care disease assessment. In addition, blood and marrow samples are also collected as part of this research study to check your immune response to the vaccine and are not part of your standard of care. The maximum amount of blood we will collect for this study in one day is 70mL (approximately 4.5 tablespoons). If more blood is required at a particular time, we will collect blood on 2 or more days spread out over 1-2 weeks.

The maximum amount of bone marrow sample we will collect is 37mL (approximately 2.5 tablespoons). **Table 5** shows the timeline and amount of samples we may collect from you during the study.

Table 5: Timeline and Sample Amount Collected during Study (in approximate tablespoons (Tbsp) or teaspoons (tsp))

Sample	When you join the study	Before you're randomized	Maintenance Cycles						
			1	2	2 (day 7)	3	4	4 (day 7)	9
Bone marrow	2.5 Tbsp	1 tsp							2 tsp
Optional bone marrow samples for future research	<1 tsp	<1 tsp							<1 tsp
Blood-forming cells	6.5 Tbsp *over 10 days	2 Tbsp	4.5 Tbsp	4.5 Tbsp	4 Tbsp	4.5 Tbsp	4.5 Tbsp	4 Tbsp	4.5 Tbsp
Optional blood samples for future research	1.5 tsp	1.5 tsp							1.5 tsp

6. Risks and Discomforts

You may have side effects while on the study. Side effects can range from mild to serious. The risks and discomforts of autologous transplant are the same if you join this study, or if you don't join this study.

You might do better or worse than you would with a standard transplant. Your healthcare team may give you medicines to help with side effects like nausea (feeling sick to your stomach). In some cases, side effects can last a long time or may never go away.

Risks of Making the Vaccine

We will collect bone marrow samples on all patients.

We will take the bone marrow samples from your hip bone with a large needle. This is called a bone marrow aspirate. After, you may feel pain and bleed at the site where the needle went in your hip. A small number of patients will have pain that runs down their legs, pain that lasts more than a few days, and/or get an infection.

If you are assigned to Treatment Group A, we will collect blood and bone marrow samples to make the vaccine (see **Table 5. Timeline and Sample Amount Collected during Study**).

We will also collect extra samples of blood through a catheter in your neck or chest (leukaphereis). The risks of leukapheresis include:

- Bleeding
- Punctured lung (hole in lung)
- Infection
- Low blood cell counts.

Risks of Medications

The risks of the chemotherapy and maintenance drugs you will get as part of the treatment are listed below.

Melphalan - Conditioning Regimen Drug (Before Transplant)

Likely (May happen in more than 20% of patients)	Less Likely (May happen in less than 20% of patients)	Rare, but Serious (May happen in less than 2% of patients)
<ul style="list-style-type: none"> • Loss of appetite • Constipation • Diarrhea • Nausea (feeling sick to your stomach) • Vomiting (throwing up) • Temporary hair loss • Sensitive skin • Infection • Low number of white blood cells ▪ Low number of platelets in the blood with increased risk of bleeding • Anemia (low number of red blood cells) • Mouth sores • Sore throat (red with swelling) • Skin breakdown (if drug leaks from vein) 	<ul style="list-style-type: none"> • Changes in heart beat • Dizzy • Feeling faint • Shortness of breath • Hepatitis (swelling of the liver) • Kidney failure • Weight loss • Feeling weak 	<ul style="list-style-type: none"> • Allergic reaction • Lung infection • Scarring of lung tissue • Seizure • Vasculitis (inflammation of blood vessels) • Low blood pressure • Sweating too much • Sterility (unable to have children) • Liver damage • Heart stops beating • New cancer of bone marrow cells

Vaccine

Likely (May happen in more than 20% of patients)	Less Likely (May happen in less than 20% of patients)	Rare, but Serious (May happen in less than 2% of patients)
<ul style="list-style-type: none"> • Twitching (at injection site) • Itching (general or at injection site) • Discomfort (at injection site) • Swelling (at injection site) • Bruising at the injection site 	<ul style="list-style-type: none"> • Fluid retention (too much water in your body) • Fatigue • Muscle aches • Itching • Fever • Abnormal Antinuclear Antibody (ANA) blood test • Decreased white blood cell count (including neutrophils) • Headache • Diarrhea • Feeling light headed • Flu-like symptoms • Night sweats 	<ul style="list-style-type: none"> • Allergic reaction leading to rashes, joint pain, kidney, heart or lung damage, drop in blood counts • Tumor growth or the spread of your cancer • Infection at the vaccine site • Change in thyroid function (fatigue, weight gain, sensitivity to cold and lack of interest or emotion) • Pulmonary embolus (blood clot in the lungs)

Likely (May happen in more than 20% of patients)	Less Likely (May happen in less than 20% of patients)	Rare, but Serious (May happen in less than 2% of patients)
	<ul style="list-style-type: none">• Elevated TSH (TSH is a thyroid hormone)• Candida infection	

GM-CSF (Leukine)

Likely (May happen in more than 20% of patients)	Less Likely (May happen in less than 20% of patients)	Rare, but Serious (May happen in less than 2% of patients)
<ul style="list-style-type: none"> • Fever • Chills • Nausea • Vomiting • Diarrhea • Fatigue • Feeling weak • Headache • Loss of appetite • Flushed face (red or pink cheeks) • Bone and muscle pain in arms, legs and feet • Red skin, swelling, or itching at needle injection site • Low blood pressure • Shortness of Breath 	<ul style="list-style-type: none"> • Rapid, irregular heartbeats • Feeling lightheaded or dizzy • Allergic reaction • Fluid in lungs or around your heart • Changes in vision • Low blood cell counts • Infection at needle injection site 	<ul style="list-style-type: none"> • Blood clots

Lenalidomide- Maintenance Treatment Drug

Likely (May happen in more than 10% of patients)	Less Likely (May happen in less than 10% of patients)	Rare, but Serious (May happen in less than 2% of patients)
<ul style="list-style-type: none"> • Low number of white blood cells (with or without fever) • Anemia; Decrease in cells that help your blood clot • Vision Blurred • Diarrhea • Pain, Constipation • Indigestion • Nausea • Vomiting • Feeling weak and unwell • Tired • Swelling • Fever • Chills • Pneumonia or other infections • Sore throat • Stuffy nose • Weight loss • Decreased appetite • High blood sugar 	<ul style="list-style-type: none"> • Abnormally low number of blood cells • Destruction of red blood cells • Heart attack • Abnormal heart beats • Heart stops working • Low oxygen to heart tissue • Dry mouth • Decreased action of intestine • Bile flow from liver slowed or blocked • Gout • Fall • Bruise • Lowered level of consciousness with drowsiness, listlessness, and apathy • Abnormal liver lab tests • Increase in liver protein that indicates inflammation in body • Loss of fluid • Diabetes • High uric acid in blood • Iron build up in body 	<ul style="list-style-type: none"> • Swelling of lungs • Over and underactive thyroid • Severe allergic conditions including: Swelling under skin; Severe skin reactions involving lining of the nose, mouth, stomach and intestines or rash leading to the separation of the top layer of skin • Tumor Lysis Syndrome (TLS) is caused by the sudden, rapid death of cancer cells in response to treatment. When cancer cells die they may spill their inner (intracellular) contents, which accumulate faster than they can be eliminated. This debris from the cancer cells can change the balance of the chemistry of the body, which can be dangerous.

<ul style="list-style-type: none"> • Chemical imbalance in blood • Pain including muscles, joints, and non-cardiac chest pain • Dizziness • Altered sense of taste • Headache • Eye lens cloudy • Abnormal sense of touch • Pain and decreased sensation in nerves • Shaking • Cough • Shortness of breath • Nosebleed • Blood clot in lower extremities, lungs, heart, brain, and other organs • Dry skin • Itching • Allergic reaction • Feeling sad • Not sleeping well 	<ul style="list-style-type: none"> • Muscle weakness • Cancer • Stroke • Tingling of skin • Fainting • Moody • Kidney failure • Breathing disorder • Excessive sweating • Night sweats • Skin redness • Swelling of skin filled with blood • Swelling of blood vessels • High or low blood pressure • Blood not getting to extremities • Clot in vein • Sudden increase in tumor size • Rapid death of cancer cells where the accumulating contents of dying cancer cells cause an imbalance in the chemistry of the body which can lead to kidney damage • Blood cancer that causes decreased number of red blood cells, white blood cells, and platelets because they do not develop normally. 	
--	---	--

You shouldn't donate blood while you're in the study and for 28 days after you stop taking lenalidomide.

Risk to the Unborn

Lenalidomide can cause severe birth defects or death of a baby if the mother or the father is taking this medicine at the time of conception or during pregnancy. **Because of this, it is extremely important that you don't get pregnant while you're taking lenalidomide.**

If you're pregnant or nursing, you're not eligible to take part in this study. Women who can become pregnant must use at least 2 forms of effective birth control while in the study or abstain from all reproductive sexual intercourse. Effective birth control is defined as the following:

1. Refraining from all acts of vaginal sex (abstinence)
2. Consistent use of birth control pills
3. Injectable birth control methods (Depo-Provera, Norplant)
4. Tubal sterilization or male partner who has undergone a vasectomy
5. Placement of an IUD (intrauterine device)
6. Use of a cervical cap or a diaphragm with contraceptive jelly and/or condoms with contraceptive foam every time you have sex.

Females taking lenalidomide have blood clots more often. Because of this, you should talk to your doctor about birth control pills and hormone replacement therapy, and the risks and benefits.

You do not need to use effective birth control only if you are a woman and cannot have children because you:

- Had a hysterectomy (your ovaries and uterus were removed), OR
- Had a bilateral oophorectomy (your ovaries were removed), OR
- Went through menopause (post-menopausal).

Reproductive Risks

The drugs used in this research study may damage your reproductive organs, affect your ability to have children, or cause birth defects if you take them while you are pregnant or nursing.

Both women who can become pregnant and their male partners should use birth control while on this study and for 28 days after maintenance treatment is stopped. **If you or your partner becomes pregnant during this study, you must tell the study doctor immediately.**

Your doctor will discuss the risks to your unborn child and options with you.

It is important that females who aren't pregnant or nursing don't become pregnant while part of the study. If you are a woman and become pregnant while on this study, we will stop the maintenance treatment drug right away.

Your study doctor will watch your health closely while you are pregnant and for 30 days after the pregnancy ends.

- **Females who join the study**

If you are female and can become pregnant, you will need to take a pregnancy test before you start the study. You should discuss ways to prevent pregnancy while you're in the study. Women who have gone through puberty might experience irregular menstrual cycles or their cycle might stop forever. This doesn't mean that you can't become pregnant. You must still use 2 effective forms of birth control during the study and continue with it for 28 days after you finish maintenance treatment.

Be sure to talk with your doctor about options for fertility planning, like storing your eggs, before starting chemotherapy treatment.

- **Males who join the study**

If you are male, your body may not be able to produce sperm (become sterile). Be sure to talk with your doctor about options for fertility planning, like banking your sperm, before starting chemotherapy treatment.

Damage to the vital organs in your body

Your vital organs include your heart, lungs, liver, intestines, kidneys, bladder and brain. The chemotherapy drugs may hurt these organs. You may develop lung problems from chemotherapy or an infection.

Some patients can have veno-occlusive disease (VOD) of the liver. Patients with VOD become jaundiced (yellow skin), have problems with their liver, retain too much water (feel swollen and uncomfortable), and have stomach swelling and pain.

If there is serious damage to your vital organs, you may have to stay in the hospital longer or return to the hospital after your transplant. Many patients get better, but these complications can cause permanent damage to your organs or death.

Relapse (return) of disease or a new blood cancer

Your disease may come back even if the transplant was successful at first.

We don't know if new blood cancers are caused by lenalidomide or other drugs. Other research looked at the number of patients who got new blood cancers after taking lenalidomide for:

- Diseases other than multiple myeloma, AND
- Relapsed multiple myeloma.

In these studies, no difference was shown in the number of patients who got new blood cancers.

Researchers for other studies of lenalidomide are still watching patients to see if they get new blood cancers. We will give you any new information that we learn about new blood cancers.

All Patients Taking Lenalidomide

In order to participate in this study you must register into and follow the requirements of the REVLIMID REMS™ program of Celgene Corporation. This program provides education and counseling on the risks of fetal exposure, blood clots and reduced blood counts. You will be required to receive counseling every 28 days during treatment with lenalidomide, follow the pregnancy testing and birth control requirements of the program that are appropriate for you and take telephone surveys regarding your compliance with the program.

You have been informed of the risk of birth defects. If you are female, you agree not to become pregnant while taking lenalidomide. For this reason, lenalidomide is provided to patients under a special distribution program called REVLIMID REMS™.

Other Information:

There may be some unknown or unanticipated discomforts or risks associated with this treatment in addition to those specified above, but every precaution will be taken to assure your personal safety and to minimize discomforts.

Throughout the study, the researchers will tell you of new information that might affect your decision to remain in the study.

If you wish to discuss the information above or any other discomforts you may experience, you may ask questions now or call your doctor _____, the Principal Investigator or contact person listed on the front page of this form.

Other RisksSerious infections

It may take many months for your immune system to recover from the chemotherapy and maintenance therapy drugs. There is an increased risk of infection during this time when your body is healing. We will give you drugs to reduce the chance of infection, but they may not work. If you have an infection, you may have to stay in the hospital longer or return to the hospital after transplant. Many patients get better, but some infections can cause death.

Unforeseen Risks

Chemo drugs can damage your blood cells, which may cause a new blood cancer to grow. We know from other MM research studies, that more patients had a second cancer after chemo and/or autologous transplant with maintenance lenalidomide than those who didn't get lenalidomide. We don't know if taking lenalidomide for a long time raises the risk of having a second cancer.

Other new risks might appear at any time during the study. These risks might be different from what is listed in this Consent Form. There may be some unknown or unanticipated discomforts or risks associated with this treatment in addition to those specified above, but every precaution will be taken to assure your personal safety and to minimize discomforts.

Other Treatments or Medicines

Some medicines react with each other, and it is important that you tell the study doctor or staff about any other drugs, treatments, or medicines you are taking. This includes non-prescription or over-the-counter medicines, vitamins, and herbal treatments.

It is also important that you tell the study staff about any changes to your medicines while you're in the study.

For more information about risks and side effects, ask your study doctor.

7. Other Treatments

Participation in this study is optional. If you choose not to take part, you may still receive non-transplant treatments or an autologous or an allogeneic transplant to treat your disease. The treatment and evaluations you would receive could be very similar to what you would receive if you join this study.

Your study doctor will talk with you about your options. If you decide not to participate in this study, your medical care will not be affected in any way.

Your other options may include:

- Treatment with other drugs, radiation, or a combination of drugs and radiation without a transplant.
- An allogeneic (donor) blood or marrow transplant that is not part of the study, or another type of transplant
- Participation in another clinical trial, if available (check with your doctor)
- No treatment for your blood cancer at this time
- Comfort care

Every treatment option has benefits and risks. Talk with your doctor about your treatment choices before you decide if you will take part in this study.

8. Possible Benefits

Taking part in this study may or may not make your health better. The information from this study will help doctors learn more about drugs used to treat MM.

This information could help people with multiple myeloma who may need a transplant in the future.

9. New Information Available During the Study

During this research study, the study doctors may learn about new information about the study drugs or the risks and benefits of the study. If this happens, they will tell you about the new information. The new information may mean that you can no longer participate in the study, or that you may not want to continue in the study.

If this happens, the study doctor will stop your participation in the study and will offer you all available care to suit your needs and medical conditions.

10. Privacy, Confidentiality and Use of Information

Your privacy is very important to us. The study doctors will make every effort to protect it. The study doctors have a privacy permit to help protect your records if there is a court case. However, some of your medical information may be given out if required by law. If this should happen, the study doctors will do their best to make sure that any information that goes out to others will not identify who you are.

Data regarding your clinical situation, including follow-up after 2 years, may be obtained from the CIBMTR, which captures information on all US transplants.

All your medical and demographic information (such as race and ethnicity, gender and household income) will be kept private and confidential. *(Name of Transplant Center)* and the organizations listed below will not disclose your participation by any means of communication to any person or organization, except by your written request, or permission, or unless required by federal, state or local laws, or regulatory agencies.

The individuals below will have access to your research and medical information. They may use this information for inspections or audits to study the outcomes of your treatment. By agreeing to participate, you consent to such inspections and to the copying of parts of your records, if required by these organizations.

We may give out your personal information if required by law. If information from this study is published or presented at scientific meetings, your name and other personal information will not be used.

Information about your transplant from your original medical records may be seen or sent to national and international transplant registries, including:

1. /Institution/
2. The Center for International Blood and Marrow Transplant Research (CIBMTR)
3. The National Marrow Donor Program (NMDP)
4. The Food and Drug Administration (FDA)
5. The National Institutes of Health (NIH), which include the National Heart, Lung, and Blood Institute (NHLBI) and the National Cancer Institute (NCI)
6. Data and Coordinating Center of the Blood and Marrow Transplant Clinical Trials Network (BMT CTN)
7. Data and Safety Monitoring Board (DSMB), not part of /Institution/
8. Study investigators.
9. Celgene, the manufacturer of lenalidomide

A description of this clinical trial will be available on <http://www.ClinicalTrials.gov>, as required by U.S. Law. This Web site will not include information that can identify you. At most, the Web site will include a summary of the results. You can search this Web site at any time. For questions about access to your medical records, please contact /name/ at /number/.

11. Ending Your Participation

The study doctor or the study sponsor may stop the study at any time, and we may ask you to leave the study. We may ask you to leave the study if you do not follow directions or if you suffer from side effects of the treatment. If we ask you to leave the study, the reasons will be discussed with you. Possible reasons to end your participation in this study include:

1. You do not meet the study requirements.
2. You need a medical treatment not allowed in this study.
3. The study doctor decides that it would be harmful to you to stay in the study.

4. You are having serious side effects.
5. You become pregnant.
6. You cannot keep appointments or take study drugs as directed.
7. The study is stopped for any reason.

You could have serious health risks if you stop treatment during the conditioning process before you receive your transplant. If you stop taking the immune suppressing drugs (see **Section 6: Risks and Discomforts**) too soon after transplant, your body could reject the stem cells or you could develop serious complications and possibly die.

We ask that you talk with the research doctor and your regular doctor before you leave the study. Your doctors will tell you how to stop safely and talk with you about other treatment choices.

If you decide to leave this study after getting the study treatment, or are asked to leave by your doctor for medical reasons, you will need to come back to the doctor's office for tests for your safety. Even if you leave the study, the information collected from your participation will be included in the study evaluation, unless you specifically ask that it not be included.

12. Physical Injury as a Result of Participation

It is important that you tell your doctor, _____ *[investigator's name(s)]* or study staff if you feel that you have been injured because of taking part in this study. You can tell the doctor in person or call him/her at _____ *[telephone number]*.

You will get all available medical treatment if you are injured from taking part in this study. You and/or your health plan will be charged for this treatment. There is no provision for free medical care or monetary compensation from the study sponsor, The National Institutes of Health or the study contributor, Celgene Corporation.

In case you are injured in this study, you do not lose any of your legal rights to ask for or receive payment by signing this form.

13. Compensation or Payment

You will not be paid for your participation in this research study. You will not get compensation or reimbursement for any extra expenses (travel, meals, etc.) you may have through your participation on this trial.

Taking part in this study might help researchers make products to sell. Celgene or others may profit from these products. You will not have any rights to the patents or discoveries that could happen from this research, and you will not receive any payments from it.

14. Costs and Reimbursements

Most of the visits for this research study are standard medical care for your autologous transplant and will be billed to your insurance company. You and/or your health plan/insurance company will need to pay for some or all of the costs of standard treatment in this study.

You or your insurance will not be charged for blood and marrow samples for research on this study. You will not pay for any extra tests that are being done for the study. Lenalidomide will be provided to you for free for two years. After that your doctor will discuss what treatment is best for you.

Some health plans will not pay these costs for people taking part in studies. Check with your health plan or insurance company to find out if they will pay.

For questions about your costs, financial responsibilities, and/or medical insurance coverage for your transplant and this study, please contact **/Center/ Financial Counselor at /Number/**.

For more information on clinical trials and insurance coverage, you can visit the National Cancer Institute's Web site at <http://cancer.gov/clinicaltrials/understanding/insurance-coverage>. You can print a copy of the "Clinical Trials and Insurance Coverage" information from this Web site.

Another way to get the information is to call 1-800-4-CANCER (1-800-422-6237) and ask them to send you a free copy.

15. For More Information

If you need more information about this study, or if you have problems while taking part in this study, you can contact the study doctor or his/her staff.

They can be reached at the telephone numbers listed here:

[Insert name and contact details]

16. Contact Someone about Your Rights

If you wish to speak to someone not directly involved in the study, or if you have any complaints about any aspect of the project, the way it is being conducted or any questions about your rights as a research participant, then you may contact:

[Insert appropriate contact details]

The ethical aspects of this study have been reviewed and approved by **[name of IRB]**.

17. Blood and Bone Marrow Samples for Future Research (Optional)

This section of the informed consent form is about future research studies that will use blood and bone marrow samples from people who are taking part in the main study. You may choose to give samples for these future research studies if you want to. You can still be a part of the main study even if you say 'no' to give samples for future research studies.

Researchers are trying to learn more about how the human body processes the drugs used for transplant and how the body recovers after transplant. This research is meant to gain knowledge that may help people in the future and make transplants even more successful.

If you agree to provide blood and bone marrow samples, here is what will happen:

- We will collect 3 extra blood samples at the same time you have routine blood tests done (Table 3):
 - Around the time of your enrollment
 - About 2 months after your transplant
 - Before the start of your 9th cycle of maintenance therapy
- The amount of blood collected from you is about 1 teaspoon (6 ml) each time.
- We will collect 3 bone marrow samples at the same time you have routine bone biopsies done. The amount of tissue collected from you is about a half of a teaspoon (3 mL) each time. We will collect samples at 3 different dates in the study (see **Table 3**):
 - Around the time of your enrollment
 - About 2 months after your transplant
 - Before the start of your 9th cycle of maintenance therapy.
- The blood and bone marrow samples will be sent to the BMT CTN Repository for processing and storage. A repository is a place that protects, stores and sends out samples for approved research studies. All research samples will be given a bar code that cannot be linked to you by future researchers testing your samples.
- Materials stored in the Repository will be used mainly by clinicians and researchers in the BMT CTN network. In the future, the unused research samples and clinical data will be made available outside of this network.

- Researchers can apply to study the materials stored in the Repository. The BMT CTN Steering Committee and/or the BMT CTN Executive Committee must approve each request before they will share samples or information with researchers. This is to make sure that the investigators requesting the samples are qualified, and that the research is of high quality.
- DNA from your stored blood samples might be used in genome-wide association (GWA) studies for a future project either done or supported by the National Institutes of Health (NIH). Genome-wide association studies are a way for scientists to find genes that have a role in human disease or treatment. Each study can look at hundreds of thousands of genetic changes at the same time.

If your coded samples are used in such a study, the researcher is required to add your test results and sample information into a shared, public research database. This public database is called the NIH Genotype and Phenotype Database and it is managed by the National Center for Biotechnology Information (NCBI). The NCBI will never have any information that would identify you, or link you to your information or research samples.

Some general things you should know about letting us store your blood samples for research are:

1. We will only store samples from people who give us permission.
2. Research is meant to gain knowledge that may help people in the future. You will not get any direct benefit from taking part. Additionally, you or your doctor will not be given results and they will not be added to your medical record.
3. A possible risk is the loss of confidentiality about your medical information. We will use safety measures with both your samples and clinical information to make sure that your personal information will be kept private. The chance that this information will be given to someone else is extremely small.
4. Your blood will be used only for research and will not be sold. The research done with your blood may help to develop new products in the future. You will not get paid for any samples or for any products that may be developed from current or future research.

You can change your mind at any time about allowing us to use your samples and health information for research.

We ask that you contact **[Principal Investigator]** in writing and let him/her know you do not want us to use your research samples or health information for research. His/her mailing address is on the first page of this form. However, samples and information that have already been shared with other researchers cannot be taken back or destroyed.

Making Your Choice

Please read each sentence below and think about your choice. After reading each sentence, please indicate your choice below. If you have any questions, please talk to your doctor or nurse, or call our research review board at [\[contact information\]](#).

No matter what you decide to do, it will not affect your care.

Statement of Consent for Research Samples

The purpose of storing blood and tissue samples, the procedures involved, and the risks and benefits have been explained to me. I have asked all the questions I have at this time and I have been told whom to contact if I have more questions. I have been told that I will be given a signed copy of this consent form to keep.

I understand that I do not have to allow the use of my blood and tissue for research. If I decide to not let you store research samples now or in the future, it will not affect my medical care in any way.

I voluntarily agree that my blood, tissue, and information can be stored indefinitely by the BMT CTN and/or NHLBI Repositories for research to learn about, prevent, or treat health problems. I also understand that my DNA and health information may or may not be used in genome-wide association studies.

Blood

- ☐ I agree to allow my blood samples to be stored for research.
- ☐ I do not agree to allow my blood samples to be stored for research.

Bone marrow

- ☐ I agree to allow my bone marrow samples to be stored for research.
- ☐ I do not agree to allow my bone marrow samples to be stored for research.

Signature

Date

Health Insurance Portability and Accountability Act 1 (HIPAA12) Authorization to use and disclose individual health information for research purpose

- **Purpose:**

As a research participant, I authorize the Principal Investigators and the researcher's staff to use and disclose my individual health information for the purpose of conducting the research study:

Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions

- **Individual Health Information to be Used or Disclosed:**

My individual health information that may be used or disclosed to do this research includes:

1. Demographic information (for example, date of birth, sex, weight)
2. Medical history (for example, diagnosis, complications with prior treatment)
3. Findings from physical exams
4. Laboratory test results obtained at the time of work up and after transplant (for example, blood tests, biopsy results)

- **Parties Who May Disclose My Individual Health Information:**

The researcher and the researcher's staff may collect my individual health information from:

[List hospitals, clinics or providers from which health care information can be requested]:

- **Parties Who May Receive or Use My Individual Health Information:**

The individual health information disclosed by parties listed in item c and information disclosed by me during the course of the research may be received and used by the following parties:

1. Dr. David Avigan, Co-Principal Investigator
2. Dr. Nina Shah, Co-Principal Investigator
3. Dr. David Chung, Co-Principal Investigator

12 HIPAA is the Health Insurance Portability and Accountability Act of 1996, a federal law related to privacy of health information

4. Celgene, its collaborators or designees
5. National Heart, Lung, and Blood Institute (NHLBI) and the National Cancer Institute (NCI), both of the National Institutes of Health (NIH),
6. Blood and Marrow Transplant Clinical Trials Network (BMT CTN) Data and Coordinating Center
7. U.S. government agencies that are responsible for overseeing research such as the Food and Drug Administration (FDA) and the Office of Human Research Protections (OHRP)
8. U.S. government agencies that are responsible for overseeing public health concerns such as the Centers for Disease Control (CDC) and federal, state and local health departments.

- **Right to Refuse to Sign this Authorization:**

I do not have to sign this Authorization. If I decide not to sign the Authorization, I will not be allowed to participate in this study or receive any treatment related to research that is provided through the study.

My decision not to sign this authorization will not affect any other treatment, payment, or enrollment in health plans or eligibility for benefits.

- **Right to Revoke:**

I can change my mind and withdraw this authorization at any time by sending a written notice to the Principal Investigator to inform the researcher of my decision.

If I withdraw this authorization, the researcher may only use and disclose the protected health information already collected for this research study. No further health information about me will be collected by or disclosed to the researcher for this study.

- **Potential for Re-disclosure:**

My individual health information disclosed under this authorization may be subject to re-disclosure outside the research study and no longer protected.

Examples include potential disclosures for law enforcement purposes, mandated reporting or abuse or neglect, judicial proceedings, health oversight activities and public health measures.

- **Genetic Information Nondiscrimination Act (GINA)**

A new federal law (2009), called the Genetic Information Nondiscrimination Act (GINA) generally makes it illegal for health insurance companies, group health plans, and employers of 15 or more persons to discriminate against you based on your genetic information.

Health insurance companies and group health plans may not request your genetic information that we get from this research. This means that they must not use your genetic information when making decisions regarding insurability. Be aware that this new federal law will not protect you against genetic discrimination by companies that sell life insurance, disability insurance, or long-term care insurance.

- **This authorization does not have an expiration date.**

TITLE: Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions**PROTOCOL NUMBER:** BMT CTN #1401**PRINCIPAL INVESTIGATOR:****Name:****Address:****Email:****Phone:****Fax:**

I have read and understood this Consent Form. The nature and purpose of the research study has been explained to me.

- I have had the chance to ask questions, and understand the answers I have been given. I understand that I may ask questions at any time during the study.
- I freely agree to be a participant in the study.
- I understand that I may not directly benefit from taking part in the study.
- I understand that, while information gained during the study may be published, I will not be identified and my personal results will stay confidential.
- I have had the chance to discuss my participation in this research study with a family member or friend.
- I understand that I can leave this study at any time, and doing so will not affect my current care or prevent me from receiving future treatment.
- I understand that I will be given a copy of this signed consent form.

Participant Name_____
Date_____
Signature_____
Date

I certify that I have provided a verbal explanation of the details of the research study, including the procedures and risks. I believe the participant has understood the information provided.

Name of Counseling Physician_____
Date_____
Signature of Counseling Physician_____
Date

APPENDIX B-2

**INFORMED CONSENT DOCUMENTS FOR BMT CTN 1401 OPTIONAL
CORRELATIVE LABORATORY STUDY PARTICIPATION**

Informed Consent to Participate in Research

Optional Correlative Laboratory Study Participation for patients enrolled on BMT CTN 1401 entitled “*Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions*”

Your Name: _____

Study Title: Optional Correlative Laboratory Study Participation for patients enrolled on BMT CTN 1401 “*Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions*”

Parent Study Title: Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions

Protocol: BMT CTN 1401 v4.0

Principal Investigator: *Insert local PI information*

Sponsor: The National Institutes of Health (NIH) is sponsoring this study by providing financial support for the coordination of this study through the Blood and Marrow Transplant Clinical Trials Network (BMT CTN).

1. Introduction

We invite you to participate in one or both of the optional correlative laboratory studies associated with the BMT CTN 1401 trial. You're being asked to join these optional studies because you were enrolled on BMT CTN 1401 and are being evaluated for randomization. Participation in these studies will involve you agreeing to allow investigators to either use previously collected research samples required by the primary trial and/or provide additional blood and marrow samples during your participation on the primary BMT CTN 1401 trial.

It is your choice to participate in these optional studies and to provide additional blood or bone marrow samples for this optional research. Even if you decide not to participate in these optional research studies, you can still participate on the primary BMT CTN 1401 study.

This Consent Form will tell you about the purpose of the optional research studies, the clinical samples needed for this research, the possible risks and benefits, other options available to you, and your rights as a research participant.

Everyone who takes part in research at [insert facility name] should know that:

- Being in any research study is voluntary.
- You will not directly benefit from being in the study. Knowledge we gain from this study may benefit others.
- If you give blood and bone marrow samples for research, you can change your mind at any time. The blood and marrow samples that you already provided will be used for research but no further research will be done after you change your mind.
- If you decide to quit the study, it will not affect your at [insert name of facility or institution].
- Please ask the study staff questions about anything that you do not understand, or if you would like to have more information.
- You can ask questions now or any time during the study.
- Please take the time you need to talk about the study with your doctor, study staff, and your family and friends. It is your decision to provide samples for research. If you decide to join, please sign and date the end of the Consent Form.

2. Study Background

The National Institutes of Health (NIH), through the Blood and Marrow Transplant Clinical Trials Network (BMT CTN), are providing staff support and money for this research study. The BMT CTN and the NIH will make decisions about how to manage the study.

For participation in one or both of these optional research studies, you will be asked to provide blood or bone marrow samples that will be shipped to a lab for further testing. Your doctor will discuss the research studies that are available to you as a part of this study.

3. Study Purpose

We are conducting these additional laboratory studies to further our understanding and knowledge about how the treatments you receive as part of the parent study (BMT CTN 1401) work.

4. Rights to Ask Questions and/or Withdraw

You have the right to ask questions about the study at any time. If you have questions about your rights as a participant or you want to leave the study, please contact:

[insert contact info]

Participating in either of these optional research studies is voluntary. You can choose not to be in these studies or leave either study at any time. However, some of the blood and bone marrow already donated may have already been used for research. If you choose not to take part or leave one of these studies, it will not affect your regular medical care in any way.

Your study doctor and study staff will be available to answer any questions that you may have about taking part in or leaving these optional research studies.

5. Study Treatment and Tests

There are two different research studies associated with the primary parent trial (BMT CTN 1401) that you are participating in. Your doctor will explain which studies you are being asked to participate on at the end of this consent form and you will be presented with one or two more informed consent forms. These consent forms will go over the study tests in detail. The schedule of blood or bone marrow collections will depend on which studies you agree to participate in. If you would like to participate in this optional research, you will be asked to sign this informed consent form along with the other consent form(s) associated with the optional research your doctor is asking you to be a part of.

If you agree to participate on either of these research studies, your blood and or bone marrow will be shipped to laboratories at either Beth Israel Deaconess Medical Center and/or the University of Wisconsin. All research samples will be given a bar coded ID that cannot be linked to you by the researchers testing your samples. Samples will not be used for purposes other than those specified in this consent form.

6. Risks and Discomforts

Information about your demographics, disease, and response to the treatments you receive as part of the main clinical study may be provided to the investigators listed in the privacy portion of this consent form. This information is used to help the investigators understand how the research samples correlate to your disease. There are no major risks to having your blood or bone marrow drawn. It can be uncomfortable to have your blood and bone marrow taken and it can sometimes leave a bruise. You might faint, but this is unlikely to happen. Only trained people will take your blood and bone marrow.

A possible risk is the loss of confidentiality about your medical information. We will use safety measures with both your samples and clinical information to make sure that your personal information will be kept private. The chance that this information will be given to someone else is extremely small.

For more information about risks and side effects, ask your study doctor.

7. Other Treatments

Participation in this study is optional. The alternative to participating on this research sample study, is not agreeing to provide blood or bone marrow samples. If you choose not to take part, your treatment on BMT CTN 1401 will not be affected

This information could help people with multiple myeloma who may need a transplant in the future

8. Possible Benefits

Taking part in this study will not make your health better. You will not get any direct benefit from taking part in this study. The information from this study will help doctors and researchers learn more about how well unrelated transplant works as treatment for people with a blood disease.

9. Privacy, Confidentiality and Use of Information

Your privacy is very important to us. The study doctors will make every effort to protect it. The study doctors have a privacy permit to help protect your records if there is a court case. However, some of your medical information may be given out if required by law. If this should happen, the study doctors will do their best to make sure that any information that goes out to others will not identify who you are.

Data regarding your clinical situation, including follow-up after 2 years, may be obtained from the CIBMTR, which captures information on all US transplants.

All your medical and demographic information (such as race and ethnicity, gender and household income) will be kept private and confidential. **(Name of Transplant Center)** and the organizations listed below will not disclose your participation by any means of communication to any person or

organization, except by your written request, or permission, or unless required by federal, state or local laws, or regulatory agencies.

The individuals below will have access to your research and medical information. They may use this information for inspections or audits to study the outcomes of your treatment. By agreeing to participate, you consent to such inspections and to the copying of parts of your records, if required by these organizations.

We may give out your personal information if required by law. If information from this study is published or presented at scientific meetings, your name and other personal information will not be used.

Information about your transplant, research samples, and treatment on BMT CTN 1401 from your original medical records may be seen or sent to national and international transplant registries, including:

1. /Institution/
2. The Center for International Blood and Marrow Transplant Research (CIBMTR)
3. The National Marrow Donor Program (NMDP)
4. The Food and Drug Administration (FDA)
5. The National Institutes of Health (NIH), which include the National Heart, Lung, and Blood Institute (NHLBI) and the National Cancer Institute (NCI)
6. Data and Coordinating Center of the Blood and Marrow Transplant Clinical Trials Network (BMT CTN)
7. Data and Safety Monitoring Board (DSMB), not part of /Institution/
8. Study investigators.
9. Dr. David Avigan and laboratory staff at Beth Israel Deaconess Medical Center/Dana Farber Cancer Institute
10. Dr. Fotis Asimakopoulos and laboratory staff at University of Wisconsin and at Medical College of Wisconsin.
11. Celgene, the manufacturer of lenalidomide

DNA from your blood samples might be used in genome-wide association (GWA) studies for a future project either done or supported by the National Institutes of Health (NIH). Genome-wide

association studies are a way for scientists to find genes that have a role in human disease or treatment. Each study can look at hundreds of thousands of genetic changes at the same time.

If your coded samples are used in such a study, the researcher is required to add your test results and sample information into a shared, public research database. This public database is called the NIH Genotype and Phenotype Database and it is managed by the National Center for Biotechnology Information (NCBI). The NCBI will never have any information that would identify you, or link you to your information or research samples.

A description of this clinical trial will be available on <http://www.ClinicalTrials.gov>, as required by U.S. Law. This Web site will not include information that can identify you. At most, the Web site will include a summary of the results. You can search this Web site at any time.

For questions about access to your medical records, please contact **/name/** at **/number**.

10. Ending Your Participation

The study doctor or the study sponsor may stop the study at any time, and we may ask you to leave the study. We may ask you to leave the study if you do not follow directions or if you suffer from side effects of the treatment. If we ask you to leave the study, the reasons will be discussed with you. Possible reasons to end your participation in this study include:

1. You do not meet the study requirements.
2. The study doctor decides that it would be harmful to you to stay in the study.
3. You become unable to donate blood or bone marrow for any reason
4. The study is stopped for any reason.

Even if you leave the study, the information collected from your participation will be included in the study evaluation, unless you specifically ask that it not be included.

11. Physical Injury as a Result of Participation

It is important that you tell your doctor, _____ *[investigator's name(s)]* or study staff if you feel that you have been injured because of taking part in this study. You can tell the doctor in person or call him/her at _____ *[telephone number]*.

You will get all available medical treatment if you are injured from taking part in this study. You and/or your health plan will be charged for this treatment. There is no provision for free medical care or monetary compensation from the study sponsor, The National Institutes of Health or the study contributor, Celgene Corporation.

In case you are injured in this study, you do not lose any of your legal rights to ask for or receive payment by signing this form.

12. Payment and Study Costs

You will not be paid for donating your blood and/or bone marrow.. You will not be paid or reimbursed for any extra expenses (travel, meals, etc.) you may have to donate blood and bone marrow.

Your blood and marrow will be used only for research and will not be sold. The research done with your blood may help to develop new products in the future. You will not get paid for any samples or for any products that may be developed from current or future research.

The visits for this research study are standard medical care for your autologous transplant and will be billed to your insurance company. You and/or your health plan/insurance company will need to pay for some or all of the costs of standard treatment in this study.

For questions about your costs, financial responsibilities, and/or medical insurance coverage for your transplant and this study, please contact **/Center/ Financial Counselor at /Number/**.

13. For More Information

If you need more information about these studies, or if you have problems while taking part in either of these studies, you can contact the study doctor or his/her staff.

They can be reached at the telephone numbers listed here:

[Insert name and contact details]

14. Contact Someone about Your Rights

If you wish to speak to someone not directly involved in the study, or if you have any complaints about any aspect of the project, the way it is being conducted or any questions about your rights as a research participant, then you may contact:

[Insert appropriate contact details]

The ethical aspects of this study have been reviewed and approved by **[name of IRB]**.

Health Insurance Portability and Accountability Act 1 (HIPAA 13) Authorization to use and disclose individual health information for research purpose

13 HIPAA is the Health Insurance Portability and Accountability Act of 1996, a federal law related to privacy of health information

- **Purpose:**

As a research participant, I authorize the Principal Investigators and the researcher's staff to use and disclose my individual health information for the purpose of conducting either of the research studies described in this consent:

Optional Correlative Laboratory Study Participation **for patients enrolled on “Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions”**

- **Individual Health Information to be Used or Disclosed:**

My individual health information that may be used or disclosed to do this research includes:

1. Demographic information (for example, date of birth, sex, weight)
2. Medical history (for example, diagnosis, complications with prior treatment)
3. Findings from physical exams
4. Laboratory test results obtained at the time of work up and after transplant (for example, blood tests, biopsy results)

- **Parties Who May Disclose My Individual Health Information:**

The researcher and the researcher's staff may collect my individual health information from:

[List hospitals, clinics or providers from which health care information can be requested]:

- **Parties Who May Receive or Use My Individual Health Information:**

The individual health information disclosed by parties listed in item c and information disclosed by me during the course of the research may be received and used by the following parties:

1. Dr. David Avigan, Co-Principal Investigator
2. Dr. Nina Shah, Co-Principal Investigator
3. Dr. David Chung, Co-Principal Investigator
4. Laboratory staff at Beth Israel Deaconess Medical Center/Dana Farber Cancer Institute
5. Dr. Fotis Asimakopoulos and laboratory staff at University of Wisconsin and at Medical College of Wisconsin.

6. Celgene, its collaborators or designees
7. National Heart, Lung, and Blood Institute (NHLBI) and the National Cancer Institute (NCI), both of the National Institutes of Health (NIH),
8. Blood and Marrow Transplant Clinical Trials Network (BMT CTN) Data and Coordinating Center
9. U.S. government agencies that are responsible for overseeing research such as the Food and Drug Administration (FDA) and the Office of Human Research Protections (OHRP)
10. U.S. government agencies that are responsible for overseeing public health concerns such as the Centers for Disease Control (CDC) and federal, state and local health departments.

- **Right to Refuse to Sign this Authorization:**

I do not have to sign this Authorization. If I decide not to sign the Authorization, I will not be allowed to participate in this study.

My decision not to sign this authorization will not affect any other treatment, payment, or enrollment in health plans or eligibility for benefits.

- **Right to Revoke:**

I can change my mind and withdraw this authorization at any time by sending a written notice to the Principal Investigator to inform the researcher of my decision.

If I withdraw this authorization, the researcher may only use and disclose the protected health information already collected for this research study. No further health information about me will be collected by or disclosed to the researcher for this study.

- **Potential for Re-disclosure:**

My individual health information disclosed under this authorization may be subject to re-disclosure outside the research study and no longer protected.

Examples include potential disclosures for law enforcement purposes, mandated reporting or abuse or neglect, judicial proceedings, health oversight activities and public health measures.

- **Genetic Information Nondiscrimination Act (GINA)**

A federal law (2009), called the Genetic Information Nondiscrimination Act (GINA) generally makes it illegal for health insurance companies, group health plans, and employers of 15 or more persons to discriminate against you based on your genetic information.

Health insurance companies and group health plans may not request your genetic information that we get from this research. This means that they must not use your genetic information when making decisions regarding insurability. Be aware that this federal law will not protect you against genetic discrimination by companies that sell life insurance, disability insurance, or long-term care insurance.

- **This authorization does not have an expiration date.**

TITLE: Optional Correlative Laboratory Study Participation for Patients Enrolled on BMT CTN 1401 entitled, “Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions”

PROTOCOL NUMBER: BMT CTN #1401

PRINCIPAL INVESTIGATOR:

Name:

Address:

Email:

Phone:

Fax:

I have read and understood this Consent Form. The nature and purpose of the research study has been explained to me.

- I have had the chance to ask questions, and understand the answers I have been given. I understand that I may ask questions at any time during the study.
- I freely agree to be a participant in the study.
- I understand that I may not directly benefit from taking part in the study.
- I understand that, while information gained during the study may be published, I will not be identified and my personal results will stay confidential.
- I have had the chance to discuss my participation in this research study with a family member or friend.
- I understand that I can leave this study at any time, and doing so will not affect my current care or prevent me from receiving future treatment.
- I understand that I will be given a copy of this signed consent form.

Participant Name

Date

Signature

Date

I certify that I have provided a verbal explanation of the details of the research study, including the procedures and risks. I believe the participant has understood the information provided.

Name of Counseling Physician

Date

Signature of Counseling Physician

Date

Supplemental Informed Consent Form to Participate in Research

Study Title: Characterization of the T cell immune response and myeloma genomic signature as predictors of immunologic and clinical response

Parent Study Title: Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions

A. What is this Document?

The purpose of this document is to give you more information on the specific research sample study your physician is asking for your consent to join. This document will explain further details of what samples are collected, why they are being collected, and where they are being sent. Details about risks, study participation, and your rights are explained in the primary informed consent form that your doctor already presented to you.

B. Why am I here?

You are being asked to participate in this study because you are participating on BMT CTN 1401 and are approaching the randomization portion of the parent study.

If you give us your permission, we would like to collect extra samples of your blood and marrow at several time points during your participation on the parent trial.

C. Why are you doing this study?

We are collecting samples to learn more about how your body responds to the treatment on the parent trial. We will collect samples on 60 patients across all treatment arms. Your participation on this study is expected to last up to 3 years.

D. What will happen to me if I join the study?

If you say you want to be in the study, we will ask you for the following samples:

- **Bone Marrow Aspirate Samples:** About 1 teaspoon of bone marrow will be collected at the following timepoints:
 - Just before you are randomized on the parent trial (about 50-80 days after your bone marrow transplant). This is one teaspoon more of bone marrow being collected at the time of a required bone marrow aspirate being performed as part of the primary trial.
 - Just before you start your 9th Cycle of Maintenance (about 1 year after your bone marrow transplant). This is one teaspoon more of bone marrow being collected at

the time of a required bone marrow aspirate being performed as part of the primary trial.

- If your disease comes back, an additional bone marrow aspirate sample will be collected at that time. This is an additional aspirate not required as part of the primary trial.
- Blood Samples:
 - About 2 teaspoons of blood will be collected just before you start Cycles 1, 2, and 4 of Maintenance Therapy. This is extra blood being collected at the time of a required blood sample collection being performed as part of the primary trial.
 - If your disease comes back, about 10 teaspoons of blood will be collected at that time. This is an additional collection not required as part of the primary trial.
- We will also use a portion of the bone marrow aspirate sample that you already provided at the time of agreeing to participate on the BMT CTN 1401 study as part of this research sample study.

E. Where will my samples be sent?

Your samples will be sent to a laboratory at Beth Israel Deaconess Medical Center.

All research samples will be tied to a number. This number will not be linked to your name or other identifying information.

Statement of Consent for Research Samples

The purpose of providing blood and bone marrow samples, the procedures involved, and the risks and benefits have been explained to me. I have asked all the questions I have at this time and I have been told whom to contact if I have more questions. I have been told that I will be given a signed copy of this consent form to keep.

I understand that I do not have to allow the use of my blood and bone marrow for this research. If I decide to not let collect research samples now or in the future, it will not affect my medical care in any way.

I voluntarily agree that my blood, bone marrow, and information can be used for the study entitled “*Characterization of the T cell immune response and myeloma genomic signature as predictors of immunologic and clinical response*”. I also understand that my DNA and health information may or may not be used in genome-wide association studies.

Signature

Date

Supplemental Informed Consent Form to Participate in Research

Study Title: VCAN Proteolysis: Investigation of a Potential Novel Immune Biomarker in Myeloma

Parent Study Title: Phase II Multicenter Trial of Single Autologous Hematopoietic Cell Transplant Followed by Lenalidomide Maintenance for Multiple Myeloma with or without Vaccination with Dendritic Cell /Myeloma Fusions

A. What is this Document?

The purpose of this document is to give you more information on the bone marrow sample study your physician is asking for your consent to join. This document will explain further details of what samples are collected, why they are being collected, and where they are being sent. Details about risks, study participation, and your rights are explained in the primary informed consent form that your doctor already presented to you. If you want to participate in this research you will be asked to sign the primary informed consent form as well as this document.

B. Why am I here?

You are being asked to participate in this study because you are participating on BMT CTN 1401 and are approaching the randomization portion of the parent study.

If you give us your permission, we would like to send a portion of a clinical bone marrow sample to study investigators at the time of randomization. The sample we are asking for is part of a clinical sample collection procedure you will have already undergone as part of the parent trial. We will not collect any additional samples as part of this study. We are asking for your consent to ship a portion of this clinical bone marrow sample to a different lab and study it for research purposes.

C. Why are you doing this study?

We are collecting bone marrow samples to learn more about how your body responds to the treatment on the parent trial. We will collect samples on 20 patients across all treatment arms. Your participation on this study is expected to last 1 month. We may use data collected on the parent trial for this study for up to 3 years after you are randomized on the parent trial.

D. What will happen to me if I join the study?

The bone marrow sample we are asking for is part of a procedure you will already undergo as part of the parent trial. We will not collect any additional samples as part of this study. We are

asking for your consent to ship a portion of this bone marrow sample to a different lab and study it for research purposes.

If you say you want to be in the study, we will not collect any additional samples. The clinical pathology lab at /Institution/ will cut a small piece of your bone marrow biopsy that you provided for the BMT CTN 1401 parent trial and ship it to investigators at University of Wisconsin. If the lab is not able to obtain enough sample from the bone marrow biopsy you provided for the parent trial (BMT CTN 1401), you will not be asked to provide additional samples and you will be removed from the study.

E. Where will my samples be sent?

Your samples will be sent to a laboratory at University of Wisconsin.

All research samples will be tied to a number. This number will not be linked to your name or other identifying information.

Statement of Consent for Research Samples

The purpose of providing bone marrow biopsy samples, the procedures involved, and the risks and benefits have been explained to me. I have asked all the questions I have at this time and I have been told whom to contact if I have more questions. I have been told that I will be given a signed copy of this consent form to keep.

I understand that I do not have to allow the use of my bone marrow biopsy samples for this research. If I decide to not let collect research samples now or in the future, it will not affect my medical care in any way. It will also not affect my participation in the Parent Trial.

I voluntarily agree that my bone marrow biopsy samples and information can be used for the study entitled “*VCAN Proteolysis: Investigation of a Potential Novel Immune Biomarker in Myeloma*”. I also understand that my DNA and health information may or may not be used in genome-wide association studies.

Signature

Date

APPENDIX C

LABORATORY PROCEDURES

APPENDIX C

LABORATORY PROCEDURES

Collection of MANDATORY Samples for TC-DC/Myeloma Fusion Vaccine Production and Immunologic Endpoints

Bone marrow aspirate and peripheral blood will be collected for patients who consent to the BMT CTN 1401 study. Sample collection will be performed periodically throughout the study and will provide clinical samples required for: (1) DC/Myeloma fusion cell vaccine production and (2) correlative laboratory testing associated with critical immunologic study endpoints.

- **TC-DC/Myeloma Fusion Vaccine Production:** Upon enrollment on the BMT CTN 1401 study, all patients will have both peripheral blood and marrow aspirate samples collected, and delivered to the clinical center's cell processing facility for the isolation and storage of myeloma tumor cells as summarized in the table below. A sample of the processed myeloma tumor cells will be shipped to the BIDMC Central Laboratory for phenotypic characterization.

For patients randomized to the cellular vaccine arm of the study, a therapeutic apheresis product will be collected and delivered to the clinical center's cell processing facility for the isolation of therapeutic cell dendritic cells (TC-DC) and the manufacturing of the TC-DC/Myeloma fusion cell vaccine as summarized in the table below. A sample of the processed TC-DC cells and manufactured TC-DC/Myeloma fusion cell vaccine will be shipped to the BIDMC Central Laboratory for phenotypic characterization. And finally, additional samples of the manufactured TC-DC/Myeloma fusion cell vaccine will be shipped to Labs, Inc. Reference Laboratory for final product release sterility testing, endotoxin testing and mycoplasma screening.

- **Research Samples for Laboratory Correlatives Associated with Immunologic Endpoints:** Required research samples for study-specific immunologic correlatives include the collection of blood and marrow aspirate samples as summarized in the table below (and in table 4.3B). The correlative studies include MRD Assessment, measurement of Plasmablast Response, NK Cell Reconstitution and the evaluation of Myeloma Reactive T-Cells. Once the samples are collected at specified time points they will be shipped on the day of collection directly to specified ancillary study laboratory for processing and testing. These samples will be tracked through GlobalTrace. Detailed procedures regarding specimen collection schedules, procedures and shipping instructions will be found in the BMT CTN 1401 Research Sample Information Guide.

Collection of Research Samples Associated with OPTIONAL Ancillary Correlative Laboratory Studies

Additional marrow and/or peripheral blood samples, often linked to mandatory research sample collections for laboratory correlatives associated with protocol-defined immunologic endpoints, will be collected from patients who have agreed to participate in one or both of the optional

laboratory studies described in Appendices J and K of this protocol. Sample collection will be done at the transplant center and will require minimum processing. Once the samples are collected at specified time points they will be shipped on the day of collection directly to specified ancillary study laboratory for processing and testing. The collection, processing and shipping of these biospecimens are summarized in the table below (and in table 4.3B). These samples will be tracked through GlobalTrace. Detailed procedures regarding specimen collection schedules, procedures and shipping instructions will be found in the BMT CTN 1401 Research Sample Information Guide.

- **Bone Marrow Aspirate or Marrow Biopsy Samples**
 - Patients participating in the **VCAN Proteolysis Assessment Study**: Once correlative study consent has been signed **and** patient has been randomized to one of the maintenance therapy arms of the trial, the clinical site will request 4 slides of decalcified marrow biopsy material to be made by the center's Clinical Pathology laboratory using stored paraffin blocks containing the processed clinical marrow tissue.
 - Patients participating in the **T Cell/Genomic Signature Study**: Once correlative study consent has been signed **and** patient has been randomized to one of the maintenance therapy arms of the trial, the clinical site will collect:
 - Additional 4 mL of marrow aspirate at two of the mandatory sample collection time points on the primary trial (1) at time of randomization, and (2) just prior to the initiation of cycle 9 of maintenance therapy.
 - **Event Driven Sample Collection**: At disease progression collect 5 mL of marrow aspirate when possible.
- **Peripheral Blood Samples**
 - Patients participating in the **T Cell/Genomic Signature Study**: Once correlative study consent has been signed **and** patient has been randomized to one of the maintenance therapy arms of the trial, the clinical site will collect:
 - Additional 10 mL of peripheral blood at three of the mandatory sample collection time points on the primary trial: just prior to the initiation of cycles 1, 2 and 4 of maintenance therapy.
 - **Event Driven Sample Collection**: At disease progression collect 40 mL of peripheral blood.

Collection of OPTIONAL Samples for Future Research

Bone marrow aspirate and peripheral blood will be collected from patients who signed consent to provide additional samples for future research. Sample collection will be done at the transplant center and will require minimum processing. Once the samples are collected at specified time

points they will be shipped on the day of collection to the BMT CTN Central Processing Laboratory for final processing and storage at the BMT CTN Research Repository. The collection, processing and shipping of these biospecimens are summarized in the table below (and in table 4.3B). These samples will be tracked through GlobalTrace. Detailed procedures regarding specimen collection schedules, procedures and shipping instructions will be found in the BMT CTN 1401 Research Sample Information Guide.

- **Bone Marrow Aspirate Samples**

Patients who consented for marrow sample collection for future research studies will have 3 mL of bone marrow aspirate collected at three time points (1) time of enrollment, (2) time of randomization, and (3) just prior to the initiation of cycle 9 of maintenance (~1 year post transplant)

- **Peripheral Blood Samples**

Patients who consented for peripheral blood sample collection for future research studies will have 6 mL blood samples collected at three time points (1) time of enrollment, (2) following randomization onto the study, and (3) just prior to the initiation of cycle 9 of maintenance (~1 year post transplant)

Mandatory Research Samples & Testing Associated with TC-DC/Myeloma Fusion Vaccine Production					
Purpose	Sample Type	Sample Collection Summary	Dates Samples Obtained	Cell Processing Laboratory	Shipping Location
Myeloma Tumor Cell Collection	30 mL Bone Marrow Aspirate	Bone marrow aspirates will be collected using sodium-heparin anticoagulant, following standard institutional procedures and promptly delivered to the center's cell processing facility for the isolation and storage of myeloma tumor cells.	At time of enrollment for all study patients	Bone marrow aspirate sample will be processed and myeloma tumor cells stored for possible future use for DC/Myeloma cell fusion vaccine production. Processing will be performed according to protocol-specific standardized procedures used by all BMT CTN 1401 center cell processing laboratories.	Transplant Center
Autologous Plasma for Vaccine Production	50 mL Peripheral Blood	Collect blood sample in five 10 mL fill green top plastic BD Vacutainer® tubes, containing Sodium-Heparin anticoagulant, and promptly deliver to the center's cell processing facility.	At time of enrollment for all study patients	Peripheral blood samples will be processed and plasma recovered for use in tumor cell processing/storage as well as subsequent DC/Myeloma fusion cell vaccine manufacturing steps.	Transplant Center
Myeloma Tumor Cell Phenotype Characterization	3 mL Processed Myeloma Tumor Cells	<i>No additional sample collection needed from patient</i>	N/A	5-7 million processed myeloma tumor cells in 3 mL tumor media will be shipped at 2-8° C by priority overnight FED EX delivery to BIDMC Laboratory for tumor lysate and immunohistochemical staining.	BIDMC Central Project Laboratory
Isolation of Therapeutic Dendritic Cells	Peripheral Blood Therapeutic Cells, Apheresis Product	A therapeutic peripheral blood apheresis product will be collected following standard institutional procedures and promptly delivered to the center's cell processing facility for the isolation of dendritic cells to be used in cellular vaccine preparation.	At time of randomization to only the DC/Myeloma Vaccine arm of the study	Apheresis products will be processed and TC-Dendritic cells recovered for manufacturing the DC/Myeloma fusion cell vaccine. Processing will be performed according to protocol-specific standardized procedures.	Transplant Center

Mandatory Research Samples & Testing Associated with TC-DC/Myeloma Fusion Vaccine Production (continued)					
Purpose	Sample Type	Sample Collection Summary	Dates Samples Obtained	Cell Processing Laboratory	Shipping Location
TC– DC Cell Phenotype Characterization	Processed TC-Dendritic cells	<i>No additional sample collection needed from patient</i>	N/A	2-5 x 10 ⁶ processed dendritic cells in media will be shipped at 2-8° C by priority overnight FED EX delivery to BIDMC Laboratory for immunohistochemical staining.	BIDMC Central Project Laboratory
TC-DC/Myeloma Fusion Cell Phenotype Characterization	Processed TC-DC/Myeloma Fusion cells	<i>No additional sample collection needed from patient</i>	N/A	1-2 x 10 ⁶ processed TC-DC/Myeloma Fusion cells in media will be shipped at 2-8° C by priority overnight FED EX delivery to BIDMC Laboratory for immunohistochemical staining.	BIDMC Central Project Laboratory
TC-DC/Myeloma Fusion Cell Vaccine Final Release Microbiological Assessment	Processed TC-DC/Myeloma Fusion cells	<i>No additional sample collection needed from patient</i>	N/A	2.0-5.0 x10 ⁶ cells (fusion cell vaccine) in 3 mL of media will be shipped at ambient temperature by priority overnight FED EX delivery to Labs, Inc., for product release sterility testing.	Labs, Inc. Reference Laboratory
				1 mL of fusion cell product media will be shipped at 2-8° C by priority overnight FED EX delivery to Labs, Inc. for endotoxin testing.	
				Two sample aliquots, each containing 0.5-1.0x10 ⁶ cells (fusion cell vaccine) in 0.5 mL media will be shipped at 2-8° C by priority overnight FED EX delivery to Labs, Inc. for Mycoplasma screening.	

Mandatory Research Samples Associated with Immunologic Study Endpoints					
Purpose	Sample Type	Sample Collection Summary	Dates Samples Obtained	Shipping Specifications	Shipping Location
MRD Assessment	6 mL Bone Marrow Aspirate	Collect bone marrow aspirated sample and place 3 mL into each of two green top plastic BD Vacutainer® tube, containing Sodium-Heparin anticoagulant. Gently mix sample by inversion 8-10 times to mix sample well with heparin anticoagulant.	Following randomization onto the study; and following the completion of cycle 9 of maintenance (~ 1 year post transplant)	One 3 mL bone marrow aspirate sample will be shipped at ambient temperature on the day of collection to RPCI by priority overnight FED EX delivery for MRD assessment.	RPCI Laboratory
				The remaining 3 mL bone marrow sample will be sent to the local pathology department for MRD assessment.	Transplant Center
Myeloma Reactive T Cells	4 mL Bone Marrow Aspirate	Collect bone marrow aspirated sample and place into a green top plastic BD Vacutainer® tube, containing Sodium-Heparin anticoagulant.	At time of enrollment, following randomization onto the study; and just prior to the initiation of cycle 9 of lenalidomide maintenance	Bone marrow aspirate sample will be shipped at at 2-8° C on the day of collection to BIDMC by priority overnight FED EX delivery.	BIDMC Laboratory
Plasmablast Response	30 mL Peripheral Blood	Collect blood sample and place into three 10 mL fill green top plastic BD Vacutainer® tubes, containing Sodium-Heparin anticoagulant. Gently mix sample by inversion 8-10 times to mix sample well with heparin anticoagulant.	<p><i>For all patients:</i> at time of randomization; and 7 days after the initiation of the 4th cycle (cycle 4+7) of maintenance therapy.</p> <p><i>Additional collections for patients in the vaccine arm:</i> immediately prior the first dose of the vaccine (cycle 2) and 7 days later (cycle 2+7); and immediately prior to the third dose of vaccine (cycle 4).</p>	Blood sample tubes will be shipped on the day of collection at ambient temperature, to the Wrammert Laboratory at Emory University by priority overnight FED EX delivery.	Emory University Laboratory
Myeloma Reactive T Cells	40 mL Peripheral Blood	Collect blood sample and place into four 10 mL fill green top plastic BD Vacutainer® tubes, containing Sodium-Heparin anticoagulant. Gently mix sample by inversion 8-10 times to mix sample well with heparin anticoagulant.	At time of enrollment, and just prior to initiation of cycles 1, 2, 3, 4, and 9 of maintenance.	Blood sample tubes will be shipped at at 2-8° C on the day of collection to BIDMC by priority overnight FED EX delivery.	BIDMC Laboratory

Mandatory Research Samples Associated with Immunologic Study Endpoints (continued)					
Purpose	Sample Type	Sample Collection Summary	Dates Samples Obtained	Shipping Specifications	Shipping Location
NK Cell Reconstitution	30 mL Peripheral Blood	Collect blood sample and place into three 10 mL fill green top plastic BD Vacutainer® tube, containing Sodium-Heparin anticoagulant. Gently mix sample by inversion 8-10 times to mix sample well with heparin anticoagulant.	Immediately prior to initiation of cycle 1 of maintenance; 7 days after initiation of cycle 2 (cycle 2+7); immediately prior to initiation of cycle 3; 7 days after initiation of cycle 4 (cycle 4+7); and just prior to the initiation of cycle 9.	Blood sample tubes will be shipped at ambient temperature on the day of collection, to MDACC by priority overnight FED EX delivery	MDACC Laboratory

OPTIONAL Undefined Future Research Sample Collections					
Purpose	Sample Type	Sample Collection Summary	Dates Samples Obtained	Shipping Specifications	Shipping /Storage Location
OPTIONAL Future Research	3 mL bone marrow aspirate	Collect bone marrow aspirated sample and place into a green top plastic BD Vacutainer® tube, containing Sodium-Heparin anticoagulant. Gently mix sample by inversion 8-10 times to mix sample well with heparin anticoagulant.	At time of enrollment, following randomization onto the study, and just prior to the initiation of cycle 9 of maintenance	Bone marrow aspirate tube will be shipped at ambient temperature on the day of collection, to the BMT CTN Research Repository by priority overnight FED EX delivery for processing and final frozen storage of marrow aliquots.	BMT CTN Research Repository
OPTIONAL Future Research	6 mL peripheral blood	Collect blood sample in a Red/Gray Top BD SST™ Tube with Silica Clot Activator & Polymer Gel. Let sample sit upright in rack for 30-60 minutes. Centrifuge for 10 minutes. Gel barrier will form separating the serum specimen from clot.	At time of enrollment, following randomization onto the study, and just prior to the initiation of cycle 9 of maintenance	Serum blood tube will be shipped at ambient temperature on the day of collection, to the BMT CTN Research Repository by priority overnight FED EX delivery for processing and final frozen storage of serum aliquots.	BMT CTN Research Repository

Research Samples Associated with Optional Correlative Laboratory Studies					
VCAN Proteolysis: Investigation of a Potential Novel Immune Biomarker in Myeloma (Appendix J)					
Purpose	Sample Type	Sample Collection Summary	Dates Samples Obtained	Shipping Specifications	Shipping Location
VCAN Proteolysis Assessment	4 slides paraffin-embedded, decalcified bone marrow biopsy	Slides of decalcified clinical marrow biopsy material will be made by the center's Clinical Pathology laboratory using stored paraffin blocks containing the processed marrow tissue.	<p>Core bone marrow biopsy required to be collected to confirm CR <i>in a subset</i> of study patients prior to randomization.</p> <p>Slides will be requested once patient is consented for this correlative study <u>and</u> has been randomized to an appropriate maintenance arm of the 1401 trial.</p>	The 4 marrow biopsy slides will be packaged and shipped at ambient temperature to the Asimakopoulos Laboratory by priority overnight FED EX delivery.	University of Wisconsin-Madison Laboratory
Characterization of the T cell Immune Response and Myeloma Genomic Signature as Predictors of Immunologic and Clinical Response (Appendix K)					
T Cell/Genomic Signature Study	4-5 mL Bone Marrow Aspirate	Collect 4-5 mL bone marrow aspirated sample (<i>based on visit requirements</i>) and place into a green top plastic BD Vacutainer® tube, containing Sodium-Heparin anticoagulant. Gently mix sample by inversion 8-10 times to mix sample well with heparin anticoagulant.	<p>4 mL marrow collected at time of randomization; and just prior to the initiation of cycle 9 of maintenance therapy.</p> <p>EVENT DRIVEN SAMPLE: At disease progression collect 5 mL of marrow when possible.</p>	Bone marrow aspirate sample will be shipped at 2-8° C on the day of collection to the BIDMC laboratory by priority overnight FED EX delivery.	BIDMC Laboratory
	10-40 mL Peripheral Blood	Collect blood sample and place into 1-4 10 mL fill green top plastic BD Vacutainer® tubes, (<i>based on visit requirements</i>), containing Sodium-Heparin anticoagulant. Gently mix sample by inversion 8-10 times to mix sample well with heparin anticoagulant.	<p>Collect 10 mL blood just prior to initiation of cycles 1, 2, and 4 of maintenance therapy.</p> <p>EVENT DRIVEN SAMPLE: At time of disease progression, collect 40 mL of blood.</p>	Blood sample tubes will be shipped at 2-8° C on the day of collection to BIDMC laboratory by priority overnight FED EX delivery.	BIDMC Laboratory

APPENDIX D

LENALIDOMIDE RISKS OF FETAL EXPOSURE, PREGNANCY TESTING GUIDELINES AND ACCEPTABLE BIRTH CONTROL METHODS

APPENDIX D

LENALIDOMIDE RISKS OF FETAL EXPOSURE, PREGNANCY TESTING GUIDELINES AND ACCEPTABLE BIRTH CONTROL METHODS

Risks Associated with Pregnancy

Lenalidomide is structurally related to thalidomide. Thalidomide is a known human teratogenic active substance that causes severe life-threatening birth defects. An embryofetal development study in animals indicates that lenalidomide produced malformations in the offspring of female monkeys who received the drug during pregnancy. The teratogenic effect of lenalidomide in humans cannot be ruled out. Therefore, a risk minimization plan to prevent pregnancy must be observed.

The use of lenalidomide in pregnant females and nursing mothers has not been studied nor has the effect of the lenalidomide on human eggs and sperm. The risks to a fetus are not known. However, because lenalidomide is related to thalidomide, and thalidomide is known to cause severe birth defects, the following requirements must be observed.

Females of childbearing potential (FCBP)[†] must agree to use two reliable forms of contraception simultaneously or to practice complete abstinence from heterosexual intercourse during the following time periods related to this study: 1) for at least 28 days before starting study drug; 2) while participating in the study; and, 3) for at least 28 days after discontinuation from the study. The two methods of reliable contraception must include one highly effective method (i.e. intrauterine device (IUD), hormonal [birth control pills, injections, or implants], tubal ligation, partner's vasectomy) and one additional effective (barrier) method (i.e. latex condom, diaphragm, cervical cap). FCBP must be referred to a qualified provider of contraceptive methods if needed.

Criteria for females of childbearing potential (FCBP)

This protocol defines a female of childbearing potential as a sexually mature female who: 1) has not undergone a hysterectomy or bilateral oophorectomy or 2) has not been naturally postmenopausal (amenorrhea following cancer therapy does not rule out childbearing potential) for at least 24 consecutive months (i.e., has had menses at any time in the preceding 24 consecutive months).

[†] A female of childbearing potential is a sexually mature female who: 1) has not undergone a hysterectomy or bilateral oophorectomy; or, 2) has not been naturally postmenopausal for at least 24 consecutive months (i.e., has had menses at any time in the preceding 24 consecutive months).

Counseling

All counseling will be conducted through the Revlimid REMS program.

For a female of childbearing potential, lenalidomide is contraindicated unless all of the following are met (i.e., all females of childbearing potential must be counseled concerning the following risks and requirements prior to the start of lenalidomide):

- She understands the potential teratogenic risk to the unborn child
- She understands the need for effective contraception, without interruption, 28 days before starting study treatment, throughout the entire duration of study treatment, dose interruption and 28 days after the end of study treatment
- She should be capable of complying with effective contraceptive measures
- She is informed and understands the potential consequences of pregnancy and the need to notify her study doctor immediately if there is a risk of pregnancy
- She understands the need to commence the study treatment as soon as study drug is dispensed following a negative pregnancy test
- She understands the need and accepts to undergo pregnancy testing based on the frequency outlined in this protocol
- She acknowledges that she understands the hazards and necessary precautions associated with the use of lenalidomide

The investigator must ensure that for females of childbearing potential:

- Complies with the conditions for pregnancy risk minimization, including confirmation that she has an adequate level of understanding
- Acknowledge the aforementioned requirements

For a female NOT of childbearing potential, lenalidomide is contraindicated unless all of the following are met (i.e., all females NOT of childbearing potential must be counseled concerning the following risks and requirements prior to the start of lenalidomide):

- She acknowledges that she understands the hazards and necessary precautions associated with the use of lenalidomide

Traces of lenalidomide have been found in semen. Male patients taking lenalidomide must meet the following conditions (i.e., all males must be counseled concerning the following risks and requirements prior to the start of lenalidomide):

- Understand the potential teratogenic risk if engaged in sexual activity with a pregnant female or a female of childbearing potential
- Understand the need for the use of a condom even if he has had a vasectomy, if engaged in sexual activity with a pregnant female or a female of childbearing potential.

Contraception

Females of childbearing potential (FCBP) enrolled in this protocol must agree to use two reliable forms of contraception simultaneously or to practice complete abstinence from heterosexual

contact during the following time periods related to this study: 1) for at least 28 days before starting study drug; 2) while participating in the study; 3) during dose interruptions; and, 4) for at least 28 days after study treatment discontinuation.

The two methods of reliable contraception must include one highly effective method and one additional effective (barrier) method. FCBP must be referred to a qualified provider of contraceptive methods if needed. The following are examples of highly effective and additional effective methods of contraception:

Highly effective methods:

- Intrauterine device (IUD)
- Hormonal (birth control pills, injections, implants)
- Tubal ligation
- Partner's vasectomy

Additional effective methods:

- Male condom
- Diaphragm
- Cervical Cap

Because of the increased risk of venous thromboembolism in patients with multiple myeloma taking lenalidomide and dexamethasone, combined oral contraceptive pills are not recommended. If a patient is currently using combined oral contraception the patient should switch to one of the effective method listed above. The risk of venous thromboembolism continues for 4 to 6 weeks after discontinuing combined oral contraception. The efficacy of contraceptive steroids may be reduced during co-treatment with dexamethasone.

Implants and levonorgestrel-releasing intrauterine systems are associated with an increased risk of infection at the time of insertion and irregular vaginal bleeding. Prophylactic antibiotics should be considered particularly in patients with neutropenia.

Pregnancy testing

Medically supervised pregnancy tests with a minimum sensitivity of 50 mIU/mL must be performed for females of childbearing potential, including females of childbearing potential who commit to complete abstinence, as outlined below.

Before starting study drug*Female Patients:*

FCBP must have two negative pregnancy tests (minimum sensitivity of 50mIU/mL) prior to starting study drug. The first pregnancy test must be performed within 10 to 14 days prior to the start of study drug and the second pregnancy test must be performed within 24 hours prior to the start of study drug (prescriptions must be filled within 7 days as required by the Revlimid REMS® program). The patient may not receive study drug until the study doctor has verified that the results of these pregnancy tests are negative.

Male Patients:

Must practice complete abstinence or agree to use a condom during sexual contact with a pregnant female or a female of childbearing potential while participating in the study, during dose interruptions and for at least 28 days following study drug discontinuation, even if he has undergone a successful vasectomy.

During study participation and for 28 days following study drug discontinuation*Female Patients:*

- FCBP with regular or no menstrual cycles must agree to have pregnancy tests weekly for the first 28 days of study participation and then every 28 days while on study, at study discontinuation, and at day 28 following study drug discontinuation. If menstrual cycles are irregular, the pregnancy testing must occur weekly for the first 28 days and then every 14 days while on study, at study discontinuation, and at days 14 and 28 following study drug discontinuation.
- At each visit, the Investigator must confirm with the FCBP that she is continuing to use two reliable methods of birth control.
- Counseling about pregnancy precautions and the potential risks of fetal exposure must be conducted at a minimum of every 28 days.
- If pregnancy or a positive pregnancy test does occur in a study patient, study drug must be immediately discontinued.
- Pregnancy testing and counseling must be performed if a patient misses her period or if her pregnancy test or her menstrual bleeding is abnormal. Study treatment must be discontinued during this evaluation.
- Females must agree to abstain from breastfeeding during study participation and for at least 28 days after study drug discontinuation.

Male Patients:

- Counseling about the requirement for complete abstinence or condom use during sexual contact with a pregnant female or a female of childbearing potential and the potential risks of fetal exposure to lenalidomide must be conducted at a minimum of every 28 days.
- If pregnancy or a positive pregnancy test does occur in the partner of a male study patient during study participation, the investigator must be notified immediately.

Additional precautions

- Patients should be instructed never to give this medicinal product to another person and to return any unused capsules to the study doctor at the end of treatment.
- Female patients should not donate blood during treatment and for at least 28 days following discontinuation of lenalidomide.
- Male patients should not donate blood, semen or sperm during treatment or for at least 28 days following discontinuation of lenalidomide.
- Only enough study drug for 28 days or one cycle of therapy (whichever is shorter) may be dispensed with each cycle of therapy.

APPENDIX E

Revlimid REMS ® PROGRAM



APPENDIX E

Revlimid Overview Revlimid REMS® for Study Participants

Revlimid REMS® description:

A program that allows patients enrolled in authorized clinical trials access to free Revlimid® through the Revlimid REMS® program.

Access to the Revlimid REMS Program:

- 1. All physicians must be registered prescribers of Revlimid® in the Revlimid REMS® Program All clinical sites must have access to the Revlimid REMS® software to enroll patients in the Revlimid REMS® program**
 - 1) Prescriber submits Registration Form via fax or RevAssist® Online (**RAO** for Revlimid) to Celgene Customer Care.
 - 2) Prescriber is registered within 15 minutes.
 - 3) Registration confirmation fax is sent to prescriber's office via fax. (For RAO, the confirmation notification is displayed on the screen immediately)
 - 4) Starter Kit is sent to prescriber's office (overnight). The starter kit will contain the following:
 - Instructions For Prescribers
 - Patient Resource Packs
 - Guide to English and Non-English Materials
 - Computer software used to generate Patient-Physician Agreement Forms (PPAF)
- 2. All studies must have an FDA letter of IND exemption or an active IND, active IRB approval and Celgene required regulatory documents.**
- 3. Patients must sign the research specific IRB-approved informed consent and be enrolled in a Celgene-approved Medical Affairs clinical trial using Revlimid®**
- 4. Celgene Customer Care Center must be contacted to confirm if a patient needs to be registered by calling 1-888-423-5436**
- 5. Patients must also sign the appropriate PPAF form and follow all the procedures of the Revlimid REMS® Program**
 - 1) Patient and Prescriber complete the PPAF together.
 - 2) The form is faxed to Celgene Customer Care or submitted electronically through **RAO**.
 - 3) Patient is registered within 15 minutes.
 - 4) Confirmation fax is sent to prescribing office notifying them that the patient is now registered. For RAO, the confirmation notification is displayed on the screen immediately.
- 6. Patients and prescribers must take the phone surveys as required by the Revlimid REMS® Program (The PPAF generated for the patient determines which phone survey questions**

will be asked.) An authorization number is provided at the completion of the phone survey, the authorization number should be noted on the prescription form.

Patient Survey requirements:

- For men: Do not need to call Celgene the first month but must call monthly starting the second month.
- For females of non - child bearing potential: Must call for the first month and then call every 6 months after.
- For females of child bearing potential: Must call for the first month and then every month after.

Prescribing Revlimid® in the Revlimid REMS® program

- Celgene Medical Affairs Operations will activate the study with Biologics upon receipt of all required regulatory documents.
- Biologics will not dispense or ship Revlimid® prior to Celgene's notification of activation.
- Prescription information **MUST BE** entered using **the BMT CTN 1401 Revlimid REMS® study specific electronic prescription form**. This form can be found on the BMT CTN SharePoint website (<https://bmtctnsp.net>)
- **An authorization number** must be on the prescription form at the time of faxing.
- Prescriptions for Revlimid® must be sent to Biologics Clinical Trial Division at the following FAX number: 919-256-0794
- Only a 28-day supply of Revlimid® may be provided per cycle sent to the actual address noted on the **Revlimid REMS® electronic study specific prescription form**.
- **Biologics will verify the authorization number** and complete the patient counseling.

Protocol compliance and drug return

- Patients will be required to return unused drug to the study site for destruction per institutional guidelines.
- Sites may request that patients maintain a diary and/or to bring their bottles in for a pill count at each visit in order to review **"patient compliance."**

IMPORTANT INFORMATION ABOUT Revlimid REMS®

- To avoid fetal exposure REVLIMID®(lenalidomide) is only available under a special restricted distribution program called Revlimid REMS®
- Only prescribers registered with Revlimid REMS® can prescribe REVLIMID®(lenalidomide)
- Only Revlimid REMS® contract pharmacies can dispense REVLIMID®(lenalidomide)
- In order to receive REVLIMID® (lenalidomide), patients must enroll in Revlimid REMS® and agree to comply with the requirements of the Revlimid REMS® program
- Information about REVLIMID®(lenalidomide) and the Revlimid REMS® program can be obtained by calling the Celgene Customer Care Center toll-free at 1-888-423-5436, or at [www. REVLIMID.com](http://www.REVLIMID.com)

How to Fill a REVLIMID® (lenalidomide) Prescription

1. Healthcare provider (HCP) instructs patient to complete patient survey
2. HCP completes survey
3. HCP completes patient prescription form
4. HCP obtains Revlimid REMS® authorization number
5. HCP provides authorization number on patient prescription form
6. HCP faxes form, including prescription
7. HCP advises patient that a representative from a Revlimid REMS® contract pharmacy will contact them
8. Revlimid REMS® contract pharmacy conducts patient education
9. Revlimid REMS® contract pharmacy calls for confirmation number
10. Revlimid REMS® contract pharmacy ships REVLIMID® with the FDA-approved MEDICATION GUIDE

APPENDIX F

HUMAN SUBJECTS

APPENDIX F

HUMAN SUBJECTS

1. Subject Consent

A conference will be held with the patient, donor and family to discuss this study and alternative treatments available for the treatment of multiple myeloma. The conference will be conducted by the Principal Investigator or other designated physician. Potential risks associated with the study interventions should be discussed as objectively as possible. Consent will be obtained using an IRB-approved consent.

The BMT CTN will provide a template of the consent form to each center. Each center will customize the template according to their local requirements and submit it for review by the local Internal Review Board (IRB). The DCC will verify the adequacy of the consent forms prior to submission to the IRB. Each center must provide evidence of IRB approval.

2. Confidentiality

Confidentiality will be maintained by individual names being masked and assigned a patient identifier code. The code relaying the patient's identity with the ID code will be kept separately at the center. The ID code will be transmitted to the network.

3. Participation of Women and Minorities and Other Populations

Women and ethnic minorities and other populations will be included in this study. Accrual of women and minorities at each center will be monitored to determine whether their rates of enrollment are reflective of the distribution of potentially eligible women and minorities expected from data reported to the CIBMTR and from published data on incidence of MM in these groups. Centers will be notified if their rates differ significantly from those expected and asked to develop appropriate recruitment strategies.

APPENDIX G
KARNOFSKY PERFORMANCE STATUS SCALE

APPENDIX G

KARNOFSKY PERFORMANCE STATUS SCALE

<u>Index</u>	<u>Specific Criteria</u>	<u>General</u>
100	Normal, no complaints, no evidence of disease.	Able to carry on normal activity; no special care needed.
90	Able to carry on normal activity, minor signs or symptoms of disease.	
80	Normal activity with effort, some signs or symptoms of disease.	
70	Care for self, unable to carry on normal activity or to do work.	Unable to work, able to live at home and care for most personal needs, varying amount of assistance needed.
60	Requires occasional assistance from others but able to care for most needs.	
50	Requires considerable assistance from others and frequent medical care	
40	Disabled, requires special care and assistance.	Unable to care for self, requires institutional or hospital care or equivalent, disease may be rapidly progressing.
30	Severely disabled, hospitalization indicated, but death not imminent.	
20	Very sick, hospitalization necessary, active supportive treatment necessary.	
10	Moribund	
0	Dead	

APPENDIX H

NEW YORK HEART ASSOCIATION (NYHA) CLASSIFICATION OF CARDIAC DISEASE

APPENDIX H**NEW YORK HEART ASSOCIATION (NYHA)
CLASSIFICATION OF CARDIAC DISEASE**

The following table presents the NYHA classification of cardiac disease.

Class	Functional Capacity	Objective Assessment
I	Patients with cardiac disease but without resulting limitations of physical activity. Ordinary physical activity does not cause undue fatigue, palpitation, dyspnea, or anginal pain.	No objective evidence of cardiovascular disease.
II	Patients with cardiac disease resulting in slight limitation of physical activity. They are comfortable at rest. Ordinary physical activity results in fatigue, palpitation, dyspnea, or anginal pain.	Objective evidence of minimal cardiovascular disease.
III	Patients with cardiac disease resulting in marked limitation of physical activity. They are comfortable at rest. Less than ordinary activity causes fatigue, palpitation, dyspnea, or anginal pain.	Objective evidence of moderately severe cardiovascular disease.
IV	Patients with cardiac disease resulting in inability to carry on any physical activity without discomfort. Symptoms of heart failure or the anginal syndrome may be present even at rest. If any physical activity is undertaken, discomfort is increased.	Objective evidence of severe cardiovascular disease.

APPENDIX I

ADVERSE EVENTS

APPENDIX I

ADVERSE EVENTS

I.1. Adverse Event Reporting

Adverse events (AEs) will be collected on calendar-driven forms and event-driven forms in AdvantageEDC..

The calendar-driven forms are those that appear in the AdvantageEDC Forms grid for each enrolled patient at designated timepoints (e.g. Day 28 post transplant) throughout the course of the study. Completion of calendar-driven forms is expected by the target date for the given assessment period.

Calendar Driven forms for the BMT CTN 1401 study are as follows:

- **Toxicity Form:** this form documents all expected toxicities for the BMT CTN 1401 study; each toxicity is also assigned a grade, based on the NCI CTCAE Version 4.0.
- **Follow Up Status Form:** this form documents the status of each patient at various intervals on the study.

Event-driven forms must be completed when a certain event triggers the appearance of the form in the AdvantageEDC Forms Grid. Most often the event-driven form is triggered by information entered on the Follow-up Status Form. Event-driven forms for the BMT CTN 1401 study are as follows:

- **Re-Admission/Hospitalization Form:** this form documents all hospital admissions, *including* the admission for transplant for this study.
- **Infection Form:** this form documents infections from Day 0 (date of transplant) through the 1-year post-transplant follow-up period.
- **Progression/Relapse Form:** the form captures detailed information associated with progression or relapse of the primary disease. DO NOT report progression or relapse as an Unexpected, Grade 3-5 Adverse Event.
- **Death Form:** this form documents the death of a patient from the time of study enrollment and randomization through the 1-year post-transplant follow-up period.
- **Adverse Event Forms:** this series of forms captures details on adverse events that are *both* unexpected and grades 3-5, based on the NCI CTCAE Version 4.0, **regardless of attribution** to any of the study interventions. These forms are also used to collect information on any SAE event required by the additional adverse event reporting requirements. These events require expedited reporting and will be reviewed by the Medical Monitor associated with the BMT CTN Data and Coordinating Center (DCC) within 2 business days of receiving the summary of the adverse event from the transplant center. If the Medical Monitor requires additional information to make his/her assessment, the transplant center will have 4 business days to respond to the request for additional information.

I.2 Reporting Expected Toxicities

Expected toxicities for all patients enrolled on BMT CTN 1401 will be collected on the BMT CTN 1401 Calendar-Driven Toxicity Form and some of the event-driven forms (Infection, Readmission/Hospitalization, Progression/Relapse, Graft Failure and Death forms). Any grade 4 expected event not collected on the calendar-driven toxicity or specified event-driven form must be reported through the expedited AE reporting system in AdvantageEDC.

I.3 Reporting Unexpected, Grade 3-5 Adverse Events

All Unexpected, Grade 3-5 Adverse Events should be reported for every patient enrolled on the study from the time of randomization until 3 years post maintenance initiation.

I.4 Reporting Secondary Malignancies

All secondary primary malignancies (SPMs), excluding non-melanoma skin cancers, experienced by participants from the time of enrollment until 3 years post maintenance initiation will be reported using the Adverse Event forms (AE1-AE6) in AdvantageEDC and must be reported within three business days of knowledge of the event. The Event Description of the Adverse Event forms should include histologic type.

I.5 Reporting Adverse Events from enrollment until randomization

No adverse event reporting is required from the time of enrollment until randomization. All patient deaths, SPMs, and progressions must still be reported during this time.

I.6 Reporting Adverse Events after the start of Lenalidomide

Additional Adverse Event reporting applies to patients who receive lenalidomide. Refer to Chapter 4 for additional instructions for reporting of Second Primary Malignancies and other special events of interest. Determination of the expectedness of adverse events should be differentiated between the initial transplant, maintenance, or vaccine therapy at the discretion of the investigator. For example, oral mucositis would be an expected risk associated with transplantation, but the investigator should assess the expectedness for oral mucositis during maintenance with or without vaccine therapy.

TABLE I-1: ADVERSE EVENTS FOR BMT CTN 1401 FROM FIRST DOSE OF LENALIDOMIDE THROUGH 28 DAYS AFTER LAST DOSE¹ BY ORGAN SYSTEM

Adverse Event	Collection Type	Collection Form ²
AUDITORY DISORDERS		
Hearing loss	Calendar-Driven	Toxicity
BLOOD AND LYMPHATIC DISORDERS		
Anemia	Calendar-Driven	Toxicity

Adverse Event	Collection Type	Collection Form ²
Disseminated intravascular coagulation ³	Event-Driven	Adverse Event Form
Febrile neutropenia ⁵	Event-Driven	Adverse Event Form
Lymphopenia	Calendar-Driven	Toxicity
Neutropenia	Calendar-Driven	Toxicity
Thrombocytopenia	Calendar-Driven	Toxicity
Thrombotic thrombocytopenic purpura/ Thrombotic microangiopathy	Calendar-Driven	Toxicity
CARDIAC DISORDERS		
Asystole ³	Event-Driven	Adverse Event Form
Atrial fibrillation ³	Event-Driven	Adverse Event Form
Atrial flutter ³	Event-Driven	Adverse Event Form
Atrioventricular block ³	Event-Driven	Adverse Event Form
Left ventricular systolic dysfunction	Calendar-Driven	Toxicity
Myocardial infarction ³	Event-Driven	Adverse Event Form
New or worsening heart failure ³	Event-Driven	Adverse Event Form
Pericardial effusion ³	Event-Driven	Adverse Event Form
Pericarditis ³	Event-Driven	Adverse Event Form
Sinus bradycardia ³	Event-Driven	Adverse Event Form
Ventricular arrhythmia	Calendar-Driven	Toxicity
Ventricular tachycardia ³	Event-Driven	Adverse Event Form
ENDOCRINE DISORDERS		
Hyperthyroidism ³	Event-Driven	Adverse Event Form
Hypothyroidism	Calendar-Driven	Toxicity
GASTROINTESTINAL DISORDERS		
Abdominal pain	Calendar-Driven	Toxicity
Anorexia	Calendar-Driven	Toxicity
Constipation	Calendar-Driven	Toxicity
Diarrhea	Calendar-Driven	Toxicity
Diverticulitis ³	Event-Driven	Adverse Event Form
Dry mouth	Calendar-Driven	Toxicity
Dysgeusia (taste alteration)	Calendar-Driven	Toxicity
Dyspepsia (heartburn)	Calendar-Driven	Toxicity
GI perforation ³	Event-Driven	Adverse Event Form
Intestinal obstruction ³	Event-Driven	Adverse Event Form
Ischemic bowel ³	Event-Driven	Adverse Event Form
Nausea	Calendar-Driven	Toxicity
Oral mucositis	Calendar-Driven	Toxicity
Vomiting	Calendar-Driven	Toxicity

Adverse Event	Collection Type	Collection Form ²
GENERAL DISORDERS		
Chest pain- non-cardiac	Calendar-Driven	Toxicity
Chills	Calendar-Driven	Toxicity
Edema- generalized	Calendar-Driven	Toxicity
Fatigue	Calendar-Driven	Toxicity
Fever	Calendar-Driven	Toxicity
Sudden death ³	Event-Driven	Adverse Event Form
HEPATOBIILIARY/PANCREAS DISORDERS		
Abnormal liver function tests	Calendar-Driven	Toxicity
Hepatitis ³	Event-Driven	Adverse Event Form
Liver failure ³	Event-Driven	Adverse Event Form
Pancreatitis ³	Event-Driven	Adverse Event Form
HEMORRHAGIC DISORDERS		
Intracranial ³	Event-Driven	Adverse Event Form
Gastrointestinal ³	Event-Driven	Adverse Event Form
Genitourinary	Calendar-Driven	Toxicity
Pulmonary/Upper respiratory ³	Event-Driven	Adverse Event Form
IMMUNE SYSTEM DISORDERS		
Allergic reaction ³	Event-Driven	Adverse Event Form
Anaphylaxis (swelling of the skin and/or swelling of the face or throat) ³	Event-Driven	Adverse Event Form
INFECTIONS		
Infections/ Sepsis	Event-Driven	Infection
INVESTIGATIONS		
Increased ALT $\geq 3.0 \times \text{ULN}^4$	Event-Driven	Adverse Event Form
Increased AST $\geq 3.0 \times \text{ULN}^4$	Event-Driven	Adverse Event Form
Increased bilirubin $\geq 3.0 \times \text{ULN}^4$	Event-Driven	Adverse Event Form
Increased creatinine	Calendar-Driven	Toxicity
Prolongation of QTc interval ⁴	Event-Driven	Adverse Event Form
Weight loss	Calendar-Driven	Toxicity
METABOLISM AND NUTRITION DISORDERS		
Dehydration	Calendar-Driven	Toxicity
Hypercalcemia	Calendar-Driven	Toxicity
Hyperglycemia	Calendar-Driven	Toxicity
Hypoglycemia	Calendar-Driven	Toxicity
Hypokalemia	Calendar-Driven	Toxicity
Hyponatremia	Calendar-Driven	Toxicity
Tumor lysis syndrome ³	Event-Driven	Adverse Event Form
MUSCULOSKELETAL AND TISSUE DISORDERS		
Arthralgia	Calendar-Driven	Toxicity

Adverse Event	Collection Type	Collection Form ²
Myalgia	Calendar-Driven	Toxicity
Muscle weakness (generalized or specific area)	Calendar-Driven	Toxicity
Musculoskeletal pain	Calendar-Drive	Toxicity
NERVOUS SYSTEM DISORDERS		
Anxiety	Calendar-Driven	Toxicity
Ataxia	Calendar-Driver	Toxicity
Confusion	Calendar-Driven	Toxicity
Cranial palsy ³	Event-Driven	Adverse Event Form
Depression	Calendar-Driven	Toxicity
Depressed level of consciousness ³	Event-Driven	Adverse Event Form
Dizziness	Calendar-Driven	Toxicity
Edema cerebral ³	Event-Driven	Adverse Event Form
Encephalopathy ³	Event-Driven	Adverse Event Form
Headache	Calendar-Driven	Toxicity
Insomnia	Calendar-Driven	Toxicity
Neuralgia ³	Event-Driven	Adverse Event Form
Neuropathy ³	Event-Driven	Adverse Event Form
Reversible posterior leukoencephalopathy syndrome (PRES) ³	Event-Driven	Adverse Event Form
Seizure ³	Event-Driven	Adverse Event Form
Severe muscle weakness/paralysis ³	Event-Driven	Adverse Event Form
Somnolence	Calendar-Driven	Toxicity
Spinal cord compression (non-malignant) ³	Event-Driven	Adverse Event Form
Stroke	Calendar-Driver	Toxicity
Syncope (fainting) ³	Event-Driven	Adverse Event Form
Tremor	Calendar-Drive	Toxicity
OCULAR/VISUAL DISORDERS		
Blurred vision ³	Event-Driven	Adverse Event Form
Conjunctivitis	Calendar-Driven	Toxicity
Sudden loss of vision ³	Event-Driven	Adverse Event Form
RENAL DISORDERS		
Cystitis Non-infective	Calendar-Driven	Toxicity
Acute kidney injury ³	Event-Driven	Adverse Event Form
Chronic kidney disease ³	Event-Driven	Adverse Event Form
RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS		
Adult respiratory distress syndrome ³	Event-Driven	Adverse Event Form
Bronchitis	Calendar-Driven	Infection
Bronchopulmonary hemorrhage ³	Event-Driven	Adverse Event Form

Adverse Event	Collection Type	Collection Form ²
Cough	Calendar-Driven	Toxicity
Dyspnea	Calendar-Driven	Toxicity
Hypoxia	Calendar-Driven	Toxicity
Pleural effusion (non-malignant) ³	Event-Driven	Adverse Event Form
Pneumonia	Calendar-Driven	Infection
Pneumonitis ³	Event-Driven	Adverse Event Form
Pulmonary Hypertension ³	Event-Driven	Adverse Event Form
Sinusitis	Calendar-Driven	Toxicity
Sore throat	Calendar-Driven	Toxicity
SKIN AND SUBCUTANEOUS TISSUE DISORDERS		
Dry skin	Calendar-Driven	Toxicity
Erythema multiforme ³	Event-Driven	Adverse Event Form
Pruritis	Calendar-Driven	Toxicity
Pyoderma gangrenosum ³	Event-Driven	Adverse Event Form
Rash	Calendar-Driven	Toxicity
Sweet's syndrome (acute neutrophilic dermatosis) ³	Event-Driven	Adverse Event Form
VASCULAR DISORDERS		
Capillary leak syndrome ³	Event-Driven	Adverse Event Form
Edema	Calendar-Driven	Toxicity
Hypertension	Calendar-Drive	Toxicity
Hypotension	Calendar-Driven	Toxicity
Thromboembolic event (DVT, pulmonary emboli) ³	Event-Driven	Adverse Event Form
Vasculitis ³	Event-Driven	Adverse Event Form
OTHER		
Pregnancy ³	Event-Driven	Adverse Event Form
Secondary malignancies ⁶	Event-Driven	Adverse Event Form
Other unexpected grade 3-5 AE	Event-Driven	Adverse Event Form

¹Last dose of lenalidomide refers to the last dose prior to permanent discontinuation of study lenalidomide.

²The form listed for collection may not be the only form that the toxicity should be reported on, but the lists the highest/most expeditious level of reporting.

³Any AEs that meets one of the criteria of a serious adverse event (SAE) as defined by the 21 CFR (312.32) and occurs from the first dose through 28 days after the last dose of lenalidomide, the Adverse Event Forms are required to be submitted.

⁴Any elevation in AST, ALT or bilirubin ≥ 3.0 times the upper limit of normal (ULN) and occurs from the first dose through 28 days after the last dose of lenalidomide, requires completion of the Adverse Event forms regardless of whether the event meets one of the seriousness criteria as defined by 21 CFR (312.32).

⁵Febrile neutropenia will be collected on the event-driven Infection Form. If febrile neutropenia meets one of the criteria of a serious adverse event (SAE) as defined by the 21 CFR (312.32) and occurs from the first dose through 28 days after the last dose of lenalidomide, the Adverse Event Forms are required to be submitted.

⁶Secondary malignancies will be collected on the adverse event forms at any time from study enrollment through 3 years post-maintenance.

APPENDIX J
VCAN ANCILLARY STUDY



VCAN Proteolysis: Investigation of a Potential Novel Immune Biomarker in Myeloma

Version 1.0

Principal Investigator

Fotis Asimakopoulos, MD, PhD

Co-Principal Investigator

Binod Dhakal, MD, MS

BMT CTN 1401 Study Co-Chairs and Protocol Officer

David Avigan, MD, Nina Shah, MD, David Chung, MD, Marcelo Pasquini, MD, MS

CHAPTER 1

1 BACKGROUND AND RATIONALE

1.1 Background

Myeloma tumor cells are critically dependent on their interactions with their cellular and non-cellular microenvironment (the extracellular matrix) in both bone-marrow and extramedullary niches (1-5). We have recently focused on the unique interplay between extracellular matrix, myeloid and stromal cells in fine-tuning the inflammatory and immune microenvironment of the myeloma niche (6). Myeloma progression is characterized by accumulation of the large matrix proteoglycan, versican (VCAN) (7). VCAN contributes to myeloma homeostasis through the activation of tumor-infiltrating myeloid cells [and resultant production of trophic cytokines, such as IL-1 β and IL-6] but also by generating immune privilege (through interference with the functionality of antigen-presenting dendritic cells (DC)) (8). By contrast, the regulated proteolysis of VCAN by ADAMTS proteases generates fragments (VCAN-matrikines, including versikine) with opposing, *immunostimulatory* activities (9). Thus VCAN proteolysis intensity (and the resultant balance between parent, tolerogenic VCAN and its immunogenic fragments) may help determine the immunogenicity of the tumor microenvironment. Because efficacy of immunotherapies depends on prior cancer-immunity “set points” (10), the VCAN proteolysis status may provide a novel predictive immune biomarker.

With regard to the relevance of the VCAN pathway to BMT CTN 1401’s scientific premise, we hypothesize that the more immunogenic marrow microenvironment generated through VCAN proteolysis will potentiate the primary effector response generated through CTN1401 vaccination. Specifically, we have shown that versikine enhances Batf3-DC differentiation and maturation (IL-12 production)(6, 9, 11). Batf3-DC are a key DC subset that is essential for tumor antigen cross-presentation (12) and for efficacy of immunotherapies such as checkpoint inhibition (13). Intratumoral Batf3-DC are also critical for the efferent arm of the immune response: Gajewski has shown that Batf3-DC in the tumor microenvironment are responsible for T-cell effector recruitment through chemokine networks involving CXCL9 and CXCL10 (14). Thus, active VCAN proteolysis will be predicted to potentiate effector responses post-ASCT (15). Improved infiltration of primary effector cells and enhanced abundance and activation of (versikine-promoted) Batf3-DC at the tumor site (or, residual disease site) will result in secondary, in situ vaccination effects that will amplify the primary effector response. Therefore, patients with active VCAN proteolysis in the bone marrow are predicted to be more likely to activate a beneficial cancer immunity cycle (16), to generate more robust effector responses and experience better clinical outcomes.

Consistent with this hypothesis, at ASH 2017 we presented data that corroborated a correlation between bone marrow VCAN proteolysis and increased CD8+ infiltration and activation post-ASCT for myeloma (Pagenkopf et al., ASH abstract #1756, ASH Annual Meeting 2017). Our findings on the immunoregulatory role of VCAN in myeloma were independently corroborated by Bruno Paiva’s group: they demonstrated that VCAN-producing macrophages expand post-autologous SCT and may be associated with MRD persistence through tolerogenic actions of intact VCAN (17).

CHAPTER 2

2 STUDY DESIGN AND PROCEDURES

2.1 Study Overview

This study is a laboratory ancillary study that will investigate the versican proteolysis status on randomized patients for BMT CTN 1401. The study will examine samples provided from 10 patients on the Lenalidomide/Vaccine/GM-CSF arm and 10 patients on either the Lenalidomide alone or the Lenalidomide/GM-CSF arms

2.2 Specific Study Aims

To investigate VCAN proteolysis as a predictive biomarker for dendritic cell (DC) vaccine-based immunotherapy post- ASCT for myeloma.

The study will correlate versican proteolysis status with study-mandated immunological endpoints (myeloma-specific T cells). It is hypothesized that versican proteolysis-*predominant* status will predict those patients that meet or exceed the primary immunologic endpoint criteria (expansion of myeloma-specific T cells).

Additionally, the study will correlate versican proteolysis status with the primary and relevant secondary clinical endpoints. It is hypothesized that versican proteolysis-*predominant* status will correlate with improved clinical outcomes.

2.3 Patient Eligibility

Patients must meet specified eligibility criteria outlined in section 2.3.1 and 2.3.2 to be registered on the study. All questions regarding eligibility criteria should be directed to the protocol coordinator at 301-251-1161.

2.3.1 Patient Inclusion Criteria

1. Patients successfully enrolled in Segment A of BMT CTN 1401
2. Patients eligible for randomization per Patient Eligibility Criteria for Randomization in Chapter 2 of BMT CTN 1401
3. Patients with an available core biopsy at time of randomization

2.3.2 Patient Exclusion Criteria

1. Patients unable or unwilling to provide informed consent for participation on the ancillary study

2.4 Study Procedures

Patients will be consented at the time of randomization.

Patients who agree to participate in the ancillary study will provide permission for the BMT CTN DCC to obtain a portion of the pre-randomization clinical bone marrow biopsy that was collected for the BMT CTN 1401 parent trial. Following randomization, 4 slides of paraffin-embedded, decalcified bone marrow tissue will be prepared from the core biopsy sample at the clinical site and shipped to University of Wisconsin for analysis.

No additional samples are required for this study.

2.5 Patients for whom adequate sample cannot be provided for the planned research study analysis will be removed from the study.

Shipment of the samples will be tracked in GlobalTrace and on the Specimen Acquisition Form in AdvantageEDC for all required samples. Detailed procedures regarding specimen collection schedules and data entry will be found in the BMT CTN 1401 Research Sample Information Guide and the BMT CTN 1401 Forms Guide.

2.6 Participant Risks

Risks of the BMT CTN 1401 procedures are described in the parent clinical trial protocol. There are no additional risks associated with this ancillary study.

2.7 Study Analysis

This is a pilot study of a small initial group and all analyses are exploratory. Results from this pilot study will inform plans for expansion into larger cohorts.

ANCILLARY STUDY REFERENCES

1. Kawano Y, Moschetta M, Manier S, Glavey S, Gorgun GT, Roccaro AM, Anderson KC, Ghobrial IM. Targeting the bone marrow microenvironment in multiple myeloma. *Immunol Rev*. 2015;263(1):160-72. Epub 2014/12/17. doi: 10.1111/imr.12233. PubMed PMID: 25510276.
2. Lemaire M, Deleu S, De Bruyne E, Van Valckenborgh E, Menu E, Vanderkerken K. The microenvironment and molecular biology of the multiple myeloma tumor. *Adv Cancer Res*. 2011;110:19-42. Epub 2011/06/28. doi: B978-0-12-386469-7.00002-5 [pii]10.1016/B978-0-12-386469-7.00002-5. PubMed PMID: 21704227.
3. Mitsiades CS, McMillin DW, Klippel S, Hideshima T, Chauhan D, Richardson PG, Munshi NC, Anderson KC. The role of the bone marrow microenvironment in the pathophysiology of myeloma and its significance in the development of more effective therapies. *Hematol Oncol Clin North Am*. 2007;21(6):1007-34, vii-viii. Epub 2007/11/13. doi: S0889-8588(07)00115-3 [pii] 10.1016/j.hoc.2007.08.007. PubMed PMID: 17996586.
4. Podar K, Chauhan D, Anderson KC. Bone marrow microenvironment and the identification of new targets for myeloma therapy. *Leukemia*. 2009;23(1):10-24. Epub 2008/10/10. doi: leu2008259 [pii] 10.1038/leu.2008.259. PubMed PMID: 18843284; PMCID: 3418600.
5. Ribatti D, Nico B, Vacca A. Importance of the bone marrow microenvironment in inducing the angiogenic response in multiple myeloma. *Oncogene*. 2006;25(31):4257-66. Epub 2006/03/07. doi: 1209456 [pii]10.1038/sj.onc.1209456. PubMed PMID: 16518413.
6. Asimakopoulos F, Hope C, Johnson MG, Pagenkopf A, Gromek K, Nagel B. Extracellular matrix and the myeloid-in-myeloma compartment: balancing tolerogenic and immunogenic inflammation in the myeloma niche. *J Leukoc Biol*. 2017. doi: 10.1189/jlb.3MR1116-468R. PubMed PMID: 28254840.
7. Hebron E, Hope C, Kim J, Jensen JL, Flanagan C, Bhatia N, Maroulakou I, Mitsiades C, Miyamoto S, Callander N, Hematti P, Asimakopoulos F. MAP3K8 kinase regulates myeloma growth by cell-autonomous and non-autonomous mechanisms involving myeloma-associated monocytes/macrophages. *Br J Haematol*. 2013;160(6):779-84. Epub 2012/12/21. doi: 10.1111/bjh.12175. PubMed PMID: 23252623; PMCID: 3594344.
8. Tang M, Diao J, Gu H, Khatri I, Zhao J, Cattral MS. Toll-like Receptor 2 Activation Promotes Tumor Dendritic Cell Dysfunction by Regulating IL-6 and IL-10 Receptor Signaling. *Cell Rep*. 2015;13(12):2851-64. doi: 10.1016/j.celrep.2015.11.053. PubMed PMID: 26711349.
9. Hope C, Foulcer S, Jagodinsky J, Chen SX, Jensen JL, Patel S, Leith C, Maroulakou I, Callander N, Miyamoto S, Hematti P, Apte SS, Asimakopoulos F. Immunoregulatory roles of versican proteolysis in the myeloma microenvironment. *Blood*. 2016;128(5):680-5. doi: 10.1182/blood-2016-03-705780. PubMed PMID: 27259980.
10. Chen DS, Mellman I. Elements of cancer immunity and the cancer-immune set point. *Nature*. 2017;541(7637):321-30. doi: 10.1038/nature21349. PubMed PMID: 28102259.
11. Hope C, Emmerich PB, Papadas A, Pagenkopf A, Matkowskyj KA, Van De Hey DR, Payne SN, Clipson L, Callander NS, Hematti P, Miyamoto S, Johnson MG, Deming DA, Asimakopoulos F. Versican-Derived Matrikines Regulate Batf3-Dendritic Cell Differentiation and Promote T Cell Infiltration in Colorectal Cancer. *J Immunol*. 2017;199(5):1933-41. doi: 10.4049/jimmunol.1700529. PubMed PMID: 28754680; PMCID: PMC5568487.
12. Vanpouille-Box C, Galluzzi L. CD103(+) cells at the forefront of anticancer immunity. *Oncoimmunology*. 2017;6(11):e1356154. doi: 10.1080/2162402X.2017.1356154. PubMed PMID: 29147613; PMCID: PMC5674943.
13. Sanchez-Paulete AR, Cueto FJ, Martinez-Lopez M, Labiano S, Morales-Kastresana A, Rodriguez-Ruiz ME, Jure-Kunkel M, Azpilikueta A, Aznar MA, Quetglas JI, Sancho D, Melero I. Cancer Immunotherapy with Immunomodulatory Anti-CD137 and Anti-PD-1 Monoclonal Antibodies Requires BATF3-Dependent Dendritic Cells. *Cancer Discov*. 2016;6(1):71-9. doi: 10.1158/2159-8290.CD-15-0510. PubMed PMID: 26493961; PMCID: PMC5036540.
14. Spranger S, Dai D, Horton B, Gajewski TF. Tumor-Residing Batf3 Dendritic Cells Are Required for Effector T Cell Trafficking and Adoptive T Cell Therapy. *Cancer Cell*. 2017;31(5):711-23 e4. doi: 10.1016/j.ccell.2017.04.003. PubMed PMID: 28486109.
15. Chung DJ, Pronschinske KB, Shyer JA, Sharma S, Leung S, Curran SA, Lesokhin AM, Devlin SM, Giralt SA, Young JW. T-cell Exhaustion in Multiple Myeloma Relapse after Autotransplant: Optimal Timing of

- Immunotherapy. *Cancer Immunol Res.* 2016;4(1):61-71. doi: 10.1158/2326-6066.CIR-15-0055. PubMed PMID: 26464015; PMCID: PMC4703436.
16. Chen DS, Mellman I. Oncology meets immunology: the cancer-immunity cycle. *Immunity.* 2013;39(1):1-10. doi: 10.1016/j.immuni.2013.07.012. PubMed PMID: 23890059.
 17. Arana P, Zabaleta A, Lasa M, Maiso P, Alignani D, Jelinek T, Segura V, van den Bossche W, Teodosio C, Damasceno D, Rodriguez-Otero P, Prosper F, Mateos M-V, Lahuerta JJ, Bladé J, van Dongen JJM, San Miguel J, Orfao A, Paiva B. High-Throughput Characterization and New Insight into the Role of Tumor Associated Macrophages (TAMs) in Multiple Myeloma (MM). *Blood.* 2016;128(22):482-.
 18. Hope C, Emmerich PB, Papadas A, Pagenkopf A, Matkowskyj KA, Van De Hey DR, Payne SN, Clipson L, Callander NS, Hematti P, Miyamoto S, Johnson MG, Deming DA, Asimakopoulos F. Versican-Derived Matrikines Regulate Batf3-Dendritic Cell Differentiation and Promote T Cell Infiltration in Colorectal Cancer. *J Immunol.* 2017. doi: 10.4049/jimmunol.1700529. PubMed PMID: 28754680.

APPENDIX K

T CELL/MYELOMA GENOMIC SIGNATURE ANCILLARY STUDY



**Characterization of the T cell immune response and
myeloma genomic signature as predictors of immunologic
and clinical response**

Version 1.0

Principal Investigator

David Avigan, MD

Co-Principal Investigator

Jacalyn Rosenblatt, MD

BMT CTN 1401 Study Co-Chairs and Protocol Officer

David Avigan, MD, Nina Shah, MD, David Chung, MD, Marcelo Pasquini, MD, MS

PROTOCOL SYNOPSIS

Characterization of the T cell immune response and myeloma genomic signature as predictors of immunologic and clinical response

Study Design:

This is a laboratory ancillary study to the BMT CTN 1401 parent clinical trial. The study will further characterize the T cell response to vaccination with respect to targeting of neoantigens, assessment of T cell clonality, and characterization of the immune landscape via CYTOF analysis. In addition, genomic characterization of immunoregulatory pathways in the myeloma cells at baseline and presence of immunoregulatory noncoding RNAs will be correlated with immunologic response. Samples will consist of additional bone marrow and peripheral blood sampling at the time of collection for other indications outlined in the parent protocol.

Primary Objective:

Determine the impact of therapy on the expansion of T cells recognizing tumor associated neo-antigens and compare response between patients randomized to the Vaccine/GM-CSF/Lenalidomide arm and patients randomized to the Lenalidomide/GM-CSF or Lenalidomide Alone arms.

Secondary Objective:

Correlate of neoantigen T cell response with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant, characterize T cell clonality and TCR identity following therapy, assess measures of the immune landscape and correlated with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant, characterize myeloma cell expression of immunoregulatory pathways by genomic analysis and correlate with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant, to characterize the presence of immunoregulatory noncoding RNAs and correlate with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant

Eligibility Criteria:

Eligible patients are patients enrolled on BMT CTN 1401 that are eligible to proceed to randomization within the BMT CTN 1401 protocol specified window. Patients must also have consented to provide optional future research samples and have provided a bone marrow aspirate for future research at time of enrollment on

BMT CTN 1401. Patients must sign the additional research sample consent prior to enrollment on this study.

Treatment Description:

Patients will be consented at time of randomization on the BMT CTN 1401 parent trial. Eligible patients will undergo sample collections at the following time points:

- **Randomization: Additional** 4mL Bone Marrow Aspirate sample, collected at the time of this required bone marrow aspirate being performed as part of the primary 1401 trial.
- **Prior to Cycles 1, 2, and 4 of Maintenance: Additional** 10mL Peripheral Blood sample, collected at the time of these required research blood collections are being performed as part of the primary 1401 trial.
- **Prior to Cycle 9 of Maintenance: Additional** 4mL Bone Marrow Aspirate sample, collated at the time of this required bone marrow aspirate being performed as part of the primary 1401 trial.
- **At time of Disease Progression:** 5mL Bone Marrow Aspirate collection and 40mL Peripheral Blood collection

Additionally, a portion of the research bone marrow aspirate sample provided to the protocol-defined lab and/or BMT CTN Biorepository at time of initial enrollment on to BMT CTN 1401 will be used for this correlative laboratory study.

Accrual Objective:

60 patients randomized on BMT CTN 1401 (30 randomized to Vaccine/Lenalidomide/GM-CSF and 30 randomized to either Lenalidomide/GM-CSF or Lenalidomide). The estimated accrual period is 2 years

Accrual Period:

The estimated accrual period is 2 years.

Study Duration:

The estimated study duration is 5 years.

CHAPTER 1

1. BACKGROUND AND RATIONALE

1.1. Background

Despite the advances in biologic therapy for Multiple Myeloma (MM), curative outcomes remain elusive and patients ultimately experience life threatening disease progression. Tumor heterogeneity results in significant plasticity in response to therapy and the emergence of resistant clones[1,2]. In a paradigm shift for cancer therapeutics, cellular immunotherapy has emerged as a highly promising strategy with the potential to eradicate diverse tumor clones and provide ongoing surveillance against disease relapse. Two critical areas of investigation in cellular immunotherapy are the development of chimeric antigen receptor (CAR) T cells and cancer vaccines.

One vital area of investigation is the development of tumor vaccines to elicit the expansion of tumor specific effector cells, maintain regulatory mechanism of host immunity to minimize off target effects, and develop long term memory response to protect against relapse. Vaccines targeting individual antigens have been developed, but clinical efficacy may be limited by the inability to target the diversity of tumor clones and the development of antigen negative variants as a means of resistance[22, 23]. The group of Avigan and Rosenblatt has developed a promising cancer vaccine in which patient derived tumor cells are fused with autologous dendritic cells (DCs), presenting a broad array of antigens that capture the heterogeneity of the tumor clonal populations. In diverse animal models, vaccination with DC/tumor fusions results in eradication of established metastatic disease[24-26]. In phase I clinical studies of patients with hematological malignancies and solid tumors, vaccination with DC/tumor fusions was well tolerated, induced the expansion of tumor specific T cells, and resulted in disease regression or stabilization in patients with advanced disease[27-30]. In a recently reported trial, vaccination of patients with acute leukemia with DC/AML fusions following chemotherapy induced durable expansion in the peripheral blood and marrow of tumor reactive lymphocytes as defined by recognition of whole tumor cells and shared tumor antigens. The role of neoantigens in driving this response is also being explored. Remarkably, 71% of vaccinated patients demonstrating ongoing remission at nearly 5 years of follow up despite a median age of 63. These results are in stark contrast to previously reported outcomes where 4-year survival of patients over the age of 55 approximates 10%.

The DC/tumor vaccine platform has been studied as a strategy to induce tumor specific immunity and disease response in patients with myeloma. In a phase I trial of patients with advanced MM (median of 4 prior lines of therapy) vaccination with DC/MM fusions was feasible, was not associated with clinical evidence of autoimmunity or cytokine release syndrome, induced the 10 fold expansion of CD4 and CD8 myeloma specific T cells, and resulted in prolonged disease stabilization in nearly 70% of patients. In a subsequent study, vaccination of patients with MM following autologous transplantation induced the persistent expansion of myeloma specific T cells and was associated with the near doubling of patients achieving complete remission between day 100 and 1 year post-transplant in the absence of other maintenance therapy. Of note, based on its immunologic potency and efficacy in targeting post-transplant residual disease in a post-transplant

phase II study in MM, the fusion vaccine was chosen for a multicenter randomized trial (CTN 1401). Patients undergoing autologous transplantation are randomized in the post-transplant period in a 2:1:1 pattern to one of three arms A) vaccination with DC/MM fusions and GM-CSF in conjunction with lenalidomide maintenance; B) lenalidomide maintenance and GM-CSF; c) lenalidomide maintenance alone. Patients undergo tumor cell collection at time of initial presentation, standard induction, stem cell mobilization and collection, and high dose melphalan with stem cell rescue. Patients undergo randomization day 56-80 post-transplant. Patients assigned to the vaccine arm undergo leukapheresis, DC generation, and vaccine production. Patients begin maintenance therapy day 90-100 post-transplant. The primary aim of the study is to compare the vaccine and nonvaccine arms with respect to patients in CR at 1 year. Secondary aims of the study include assessment of PFS at 2 years, and post-transplant conversion from PR to CR. Immunologic endpoints include an assessment of the impact of therapy on presence of myeloma specific T cells in the peripheral blood and bone marrow. Serial samples of peripheral blood and bone marrow are conducted to provide material for this analysis. 17 leading cancer centers who have been trained and validated for vaccine production as a part of an FDA approved protocol. This study provides an important precedent demonstrating feasibility and the potential for expanding the fusion vaccine platform to a multicenter setting to definitively demonstrate vaccine efficacy.

A vital aspect of the design of immunotherapy clinical trials is the development of effective platforms to further characterize the nature of the immunologic response, establish assays that reflect the effect of therapy on the immunologic milieu, and most, importantly, determine those measures that most closely correlate with clinical efficacy. In addition, the identification of biomarkers predictive of immune responsiveness is critical to apply these therapies to the appropriate patient subsets. Personalized cellular therapy offers a unique platform to assess the relative contribution of immunologic response to shared as compared to neo-antigens as determinative of clinical outcome. While mutational burden has been defined as predictive of response to checkpoint inhibition in solid tumor settings, the relevance of neo-antigen burden and associated immunologic response has not been well defined for myeloma or in patients following vaccine therapy.

In the present accessory study, we will examine the impact of therapy on the functional and phenotypic assessment on the T cell repertoire including clonal nature of the T cell response, the relative role of shared vs. neoantigens, the functional competence of tumor-specific T cells, and the durability of T cell expansion. In addition, we will characterize landscape of the tumor microenvironment to identify tumor- and immune-related cell subsets (effector, antigen presenting, and accessory cell populations, MM subpopulations) associating with response or resistance to the different immunotherapies. We will utilize whole genome sequencing (WGS) and RNA seq analysis to further interrogate the impact of tumor immunobiology, patterns of T cell clonality, identification of T cell receptors, and characterization of the immune repertoire on response. In this context, utilizing a platform of 60 microRNAs that impact immune presentation and regulation, including miR200c and miR34a, we will determine if levels are predictive of immune responsiveness.

CHAPTER 2

2. STUDY DESIGN

2.1. Study Overview

The study is a laboratory ancillary study that will further characterize the nature of the T cell response to vaccination and provide for genomic characterization of immunoregulatory aspects of the tumor and microenvironment. These assessments will occur with tumor cells acquired at time of study entry, immune cells obtained from bone marrow prior to maintenance therapy and cycle 9; T cells obtained from peripheral blood prior to cycles 1, 2, 4 and at time of disease progression (if applicable); and characterization of the immune landscape from peripheral blood obtained prior to cycle 5 of maintenance. Each of the blood and bone marrow samples are being obtained at an already designated time of bone marrow aspiration or peripheral blood draw in the parent 1401 protocol.

2.2. Hypothesis and Specific Objectives

2.2.1. Hypothesis

Evidence of tumor specific immunity as manifested by response to neo-antigens, oligoclonal expansion and emergence of tumor directed TCR will be more common with the vaccinated cohort and will correlate with expansion of myeloma specific T cells in the blood and achievement and maintenance of complete response at 1 year post-transplant. Similarly, genomic analysis of myeloma cells and tumor microenvironment will demonstrate a correlation between presence of immune activation pathways and microRNA and better response to vaccination.

2.2.2. Primary Objective

Determine the impact of therapy on the expansion of T cells recognizing tumor associated neo-antigens and compare response between the Vaccine/GM-CSF/Lenalidomide arm and patients randomized to the Lenalidomide/GM-CSF or Lenalidomide Alone arms.

2.2.3. Secondary Objectives

Correlate of neoantigen T cell response with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant

Characterize T cell clonality and TCR identity following therapy

Assess measures of the immune landscape and correlated with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant

Characterize myeloma cell expression of immunoregulatory pathways by genomic analysis and correlate with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant

Characterize the presence of immunoregulatory noncoding RNAs and correlate with expansion of

myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant

2.3. Patient Eligibility

Patients must meet specified eligibility criteria outlined in section 2.3.1 and 2.3.2 to be registered on the study. All questions regarding eligibility criteria should be directed to the protocol coordinator at 301-251-1161.

2.3.1. Patient Inclusion Criteria

1. Patients successfully enrolled in Segment A of BMT CTN 1401
2. Patients eligible for randomization per Patient Eligibility Criteria for Randomization in Chapter 2 of BMT CTN 1401
3. Patients with and available bone marrow aspirate material remaining at the BIDMC protocol-defined laboratory and/or BMT CTN Biorepository that was collected at the time of initial enrollment on BMT CTN 1401

2.3.2. Patient Exclusion Criteria

1. Patients unable or unwilling to undergo bone marrow aspiration or peripheral blood sampling
2. Patients unable or unwilling to provide informed consent for participation on the ancillary study

2.4. Study Procedures

Patients who agree to participate in the ancillary study will provide additional blood and bone marrow samples during the course of the parent clinical trial as described below. For additional information on collection and shipment of the required samples, refer to the BMT CTN 1401 Research Sample Information Guide.

2.4.1. Peripheral Blood Collection

Prior to cycles 1, 2, and 4 of maintenance, 10 mL of peripheral blood will be analyzed for Characterization of peripheral blood T cells with respect to clonality, TCR analysis, and neoantigen recognition and plasma analysis for immunoregulatory microRNA signature. This sample is collected in addition to the correlative study samples collected at these time points for the BMT CTN 1401 parent trial.

At the time of progression, 40 mL of peripheral blood will be analyzed for characterization of marrow infiltrating T cells with respect to clonality, TCR analysis quantifying T cell subsets, and neoantigen recognition and plasma analysis for immunoregulatory microRNA signature. This sample is an additional collection time point for the ancillary study only. Samples at time of disease progression should only be collected on patients who have not yet completed their follow up on the BMT CTN 1401 parent trial.

2.4.2. Bone Marrow Aspirate Collection

At time of initial tumor collection 4 mL of bone marrow aspirate are stored in the BMT CTN Bioprepository for patients that consent to provide samples for future research. These samples will be obtained from the biorepository for genomic analysis of tumor cells for immune regulatory signature and determination of neoantigen targets. No additional samples are required at this timepoint.

At time of randomization & prior to cycle 9 of maintenance 4 mL of bone marrow aspirate will be collected for characterization of marrow infiltrating T cells with respect to clonality, TCR analysis quantifying T cell subsets, and neoantigen recognition, assessment of density of antigen expression by myeloma cells and subsets of immunoregulatory cells and, plasma analysis for immunoregulatory microRNA signature. This sample is collected in addition to the correlative study samples collected at these time points for the BMT CTN 1401 parent trial.

At the time of disease progression *5 mL of bone marrow aspirate will be collected for genomic analysis of tumor cells for immune regulatory signature and determination of neoantigen targets, characterization of marrow infiltrating T cells with respect to clonality, TCR analysis quantifying T cell subsets, and neoantigen recognition, plasma analysis for immunoregulatory microRNA signature.* This sample is an additional collection time point for the ancillary study only. Samples at time of disease progression should only be collected on patients who have not yet completed their follow up on the BMT CTN 1401 parent trial.

2.5. Participant Risks

Risks of the BMT CTN 1401 procedures are described in the parent clinical trial protocol. This ancillary study involves additional sampling of bone marrow and peripheral blood at the time these procedures are performed for other indications. Risks of additional sampling include bruising and pain at the site of sample acquisition.

CHAPTER 3

3. STUDY ENDPOINTS

3.1. Ancillary Testing

3.1.1. Characterization of the immunologic response to the neoantigens

Bone marrow mononuclear cells will be isolated from aspirate samples obtained at enrollment to the parent study (1401) by ficoll density centrifugation and segregated into CD38+ plasma cell fraction and CD38- nonmalignant cells. Whole Genome Sequencing (WGS) and RNA sequencing will be performed on malignant plasma cell and normal cell fractions tumor samples to characterize the tumor associated mutational burden. Putative neoantigens will be identified based on the computational algorithms established by Dr. Bhasin. Putative antigens arising from mutational events are further evaluated by algorithms predicting proteasome cleavage, transport through TAP transporter and MHC affinity. The peptides with a proteasome cleavage site at their C terminal but not in the center and with high/moderate TAP binding affinity will be used for MHC binding analysis. The MHC binding analysis will be performed using HLA Pred (combination of Propred/Propred1) and IEDB consensus prediction methods. The peptides that are predicted high/moderate binders by both IEDB ($IC_{50} < 500nM$) and HLA Pred (Score > Threshold 3%) algorithms will be used for CTL epitope prediction using CTL Pred. Candidate immunogenic peptides will be synthesized and used as antigens for pulsing autologous antigen presenting cells and testing their immunogenic potential *in vitro* using standard T cell assays. Peptides will be pulsed onto PBMCs collected prior to cycle 1, 2, and 4 of maintenance. The prevalence of antigen specific T cells will be quantified by measuring the percentage of CD4 and CD8 T cells that express IFN γ by intracellular FACS analysis. The potential role of antigenic drift in tumor immune escape will be assessed with respect to presence of the identified neo-antigens in samples obtained at disease progression. Immune response to neo-antigens and shared antigens will be compared between patients undergoing vaccination and lenalidomide maintenance alone, and who do and do not achieve CR following vaccination to correlate response with clinical outcome.

3.1.2. Assessment of T cell clonality and function

T cells will be isolated at baseline and following treatment and will be subjected to RNA seq to identify patterns of clonality as previously described. We will perform a computational algorithm to extract of TCR repertoires from bulk RNA-Seq data for quantitative analysis. Single cell analysis of cytokine expression will be assessed to further characterize the functional status of T cells.

3.1.3. Characterization of the immune landscape

The immunologic landscape in the peripheral blood and bone marrow will be interrogated for the bone marrow and peripheral blood sample drawn prior to cycle 1 maintenance therapy and prior to cycle 9 of maintenance. We will use a quantitative flow cytometry based analysis (QIFIKIT, Agilent) to determine the actual antigen density on the surface of MM cells. We will assess specific immune and tumor related cellular parameters. Effector cells will be characterized including CD4⁺ and CD8⁺ α/β T cells, γ/δ T cells, NK (CD3⁺CD56⁺) cells and B cells (CD19). Effector T cells will

be further analyzed for the presence and temporal changes in naïve (Tn: CD45RA⁺,CCR7⁺) effector memory (Tem: CD45RA⁺CCR7⁺) central memory (Tcm; CD45RA⁺CCR7⁺) and terminally differentiated effector T cells (TEMRA; CD45RA⁺CCR7⁺) T cell activation markers including HLA-DR, CD25, CD69, CD38, CD137, Granzyme B. Regulatory cells including Tregulatory (CD25⁺CD25⁺CD127^{dim}Foxp3⁺) B regulatory cells (CD19⁺CD38⁺CD24⁺IL-10⁺) and Myeloid derived Suppressor Cells (MDSC; CD11b⁺CD33⁺ CD38⁺ CD14⁺ or CD15⁺) will be quantified as well as expression levels of immune checkpoint receptors PD-1, LAG3, TIM-3, CTLA-4, BTLA, TIGIT, CD47 and GITR. MM cells will be isolated from the bone marrow samples by their typical markers CD138, CD38, CD56 and analyzed for the potential markers for immune escape (PDL-1 PDL-2 (ligand PD-1), GAL-9 (ligand TIM3), HVEM (ligand BTLA), Sirp1a (ligand CD47) and CD155 (ligand TIGIT)) and immune resistance (survivin, Xiap1, MCL-1 and NOTCH).

3.1.4. Genomic analysis to assess myeloma associated immunoregulatory pathways and clonal evolution

Malignant plasma cells will be isolated from bone marrow aspirate samples obtained at time of enrollment in the parent 1401 study as previously described. Expression patterns of a previously identified panel of immune regulatory pathways will be determined. Similarly clonal evolution as evidenced by mutational pattern will be determined in patients demonstrating evidence of disease relapse.

3.1.5. Noncoding RNAs as predictors of response

Noncoding RNAs such as microRNAs (miRNA) are critical mediators of oncogenesis and are now being investigated as immunoregulatory molecules. We have demonstrated that miR200c and miR34a modulate tumor expression of PDL1 and recruitment of MDSCs into the bone marrow microenvironment, respectively. Using a panel of 60 miRNA families identified by the Slack lab as having immunoregulatory properties, we will examine how their pattern of expression at baseline is predictive of subsequent immune and clinical response and how alterations in expression correlated with disease progression and immune escape. Serum obtained from bone marrow and peripheral blood samples will be assessed using the Abcam Firefly platform to quantify levels of the target miRNA families.

3.2. Definition of Disease Status

3.2.1. Assessment of Disease Status

Patient disease status will be assessed per Chapter 3 of the BMT CTN 1401 parent trial.

3.2.2. Disease Assessment Comparison

The parent trial will assess the rates of VGPR or better (VGPR, nCR, CR and sCR) responses at specific time points in all arms. The ancillary study will perform immunologic and genomic analysis to further characterize the immunologic response, immune landscape, and immunoregulatory features of the myeloma cells at baseline. These findings will be compared between patients randomized to the Vaccine/GM-CSF/Lenalidomide arm and patients randomized to the Lenalidomide/GM-CSF or Lenalidomide Alone arms and will be correlated with achievement of CR, maintenance of CR at 1 year and PFS.

3.2.3. Progression-Free and Overall Survival

The event for PFS and OS are defined in the BMT CTN 1401 parent clinical, and the same definition is used for the ancillary study.

CHAPTER 4

4. PATIENT ENROLLMENT AND EVALUATION

4.1. Screening and Enrollment Procedures

Patients meeting eligibility criteria defined in Chapter 2 will be consented prior to Randomization on BMT CTN 1401. Patients that are eligible will proceed with randomization on the BMT CTN 1401 parent trial. The Segment B enrollment form will include a question regarding participation on this ancillary study. A separate ancillary study enrollment form is not required. Once the patient is randomized on the BMT CTN 1401 parent trial, a schedule of assessments will be generated which will include any required data collection for the ancillary study.

4.2. Patient Evaluations and Assessments

4.2.1. Data Reporting

Sample collection will be tracked in GlobalTrace and on the Specimen Acquisition Form in AdvantageEDC for all required samples. Detailed procedures regarding specimen collection schedules and data entry will be found in the BMT CTN 1401 Research Sample Information Guide and the BMT CTN 1401 Forms Guide.

4.2.2. Adverse Event Reporting

Any adverse events that may occur as a result of a bone marrow aspiration or peripheral blood draw performed as a part of the Ancillary study will be captured through the mechanism specified in the parent trial.

4.2.3. Follow Up Schedule

4.2.3: FOLLOW UP SCHEDULE

Study Visit	Target Day
Tumor Cell Collection ¹	<i>No additional sample needed at this timepoint²</i>
Randomization ¹	±14 days from date of randomization
Start of Maintenance ¹ Cycle 1	≤ 3 ³ days prior to initiation of maintenance
Cycle 2 ¹	≤ 3 days prior to Day 1 of Cycle 2
Cycle 4 ¹	≤ 3 days prior to Day 1 of Cycle 4
Cycle 9 ¹	≤ 3 days prior to Day 1 of Cycle 9
Disease Progression ⁴	≤ 14 days after date of progression

¹Corresponds to existing BMT CTN 1401 Study Visit

²Sample will be obtained from BMT CTN Biorepository

³In the case that a sample is collected and then Maintenance therapy is delayed more than 14 days, the Prior to Cycle 1 sample should be recollected if possible.

⁴Corresponds to new study visit for this ancillary study

4.2.4. Schedule of Assessments

The sample collection schedule for this ancillary study is outlined below and in Table 4.3.4.

1. Bone Marrow Aspirate: A bone marrow aspirate will be collected and shipped to the BIDMC Lab as outlined below:
 - a. Prior to randomization (4mL)
 - b. Prior to Cycle 9 of maintenance (4mL)
 - c. At the time of disease progression (5mL)
2. Peripheral Blood: A peripheral blood sample will be collected and shipped to the BIDMC Lab as outlined below:
 - a. Prior to cycles 1, 2, and 4 of maintenance (10mL)
 - b. At time of disease progression (40mL)

TABLE 4.2.4: Sample Collection Time Points

Sample Collection	Tumor Cell Collection (At Enrollment)	Pre-Randomization	Cycles of maintenance				Disease Progression
			1	2	4	9	
Bone Marrow Aspirate	X ¹	X				X	X
Total Marrow Volume (mL)	4mL	4mL				4mL	5mL
Peripheral Blood			X	X	X		X
Total Blood Volume (mL)			10 mL	10 mL	10 mL		40mL

¹ No additional sample required. Sample will be obtained from remaining material at BIDMC lab and/or BMT CTN Biorepository

CHAPTER 5

5. STATISTICAL CONSIDERATIONS

5.1. Study Design and Objectives

5.1.1. Primary Objective

Determine the impact of therapy on the expansion of T cells recognizing tumor associated neo-antigens and compare response between patients randomized to the Vaccine/GM-CSF/Lenalidomide arm and patients randomized to the Lenalidomide/GM-CSF or Lenalidomide Alone arms.

5.1.2. Secondary Objectives

Correlate of neoantigen T cell response with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant

Characterize T cell clonality and TCR identity following therapy

Assess measures of the immune landscape and correlated with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant

Characterize myeloma cell expression of immunoregulatory pathways by genomic analysis and correlate with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant

Characterize the presence of immunoregulatory noncoding RNAs and correlate with expansion of myeloma specific T cell and achievement and maintenance of complete remission at 1 year post-transplant

5.2. Sample Size and Power Calculations

Accrual to this study is limited to patients participating in the BMT CTN 1401 trial. The target accrual for this study is 188 patients. Assuming 30% of patients will not proceed to randomization, we plan for approximately 66 patients randomized to the vaccine arm and 33 randomized to one of two control arms. Based on the prior cooperative group studies, we anticipate approximately 40% of patients will achieve CR and sCR at 1 year. In the prior phase II study of patients undergoing post-transplant vaccination in the absence of lenalidomide maintenance, CR was achieved in 29% and 54% of patients at 100 days and 1 year post-transplant respectively. Given that we anticipate that lenalidomide maintenance will enhance vaccine efficacy, we will consider the experimental arm of vaccine/GM-CSF + maintenance promising if the CR rate at 1 year is improved from 40% to 60%. A sample size of 66 patients (132 total) randomized to the vaccine arm and the combined no-vaccine arms will have 85% power to detect this improvement in the CR rate at 1 year from 40% to 60%, using a two-sample comparison of proportions with a one-sided type I error of 10%.

5.3. Analysis of the Primary Endpoint

The Ancillary study will compare the expansion of peripheral blood T cells recognizing neo-antigens (prior to vaccination to peak treatment levels) in patients receiving the DC/MM fusion vaccine as compared to standard maintenance patients. We will quantify neo-antigen specific T cells by determining expression of IFN γ in response to exposure to neoantigen peptides in samples obtained prior to and following study therapy. We hypothesize that vaccination will result in the significant expansion of these T cells in contrast to patients receiving nonvaccine maintenance therapy. Of note, lenalidomide may amplify the presence of the T cell population somewhat via nonspecific immune stimulation of tumor specific populations that are observed during post-transplant lymphopoietic reconstitution.

The primary endpoint of log 10 peak change in tumor reactive T cells from baseline will be compared across the 3 groups using analysis of variance, and if significant then two-sample t-tests will be conducted for each pairwise comparison to determine whether the vaccine and nonvaccine groups are different from one another. A secondary analysis of peak change in tumor reactive T cells will be done by comparing the proportions of patients who experience >10-fold increase in IFN γ expression, using a chi-squared test. Myeloma reactive T- cell response profiles over time will be compared between the treatment groups using linear mixed models for repeated measures data. Peak immune response as well as most recent immune response will be compared between those in CR and those not in CR at each assessment, using a two-sample t-test, to assess the relationship between immune response and clinical response. Peak and most recent immune response will also be considered as time-dependent covariates in a Cox proportional hazards model to assess their impact on progression-free survival.

We will assess whether there is a 2 fold difference in the percentage of the vaccine and nonvaccine groups achieving a minimum 10-fold increase of T-cell recognizing neoantigens, keeping power = 85%, $\alpha = 0.10$, using 2-sided Fisher's exact test.

For instance, if 3 out of 15 pts (20%) in non-vaccine group have at least 10-fold increase of T-cell recognizing neoantigens", then with 85% power and at $\alpha = 0.10$ we can say vaccine group will have 66% of patients that have "at least 10-fold increase of T-cell recognizing neoantigens" and detecting a difference of about 46% (20% vs 66%) between non-vaccine group ($n_1 = 15$) and vaccine group ($n_2 = 30$).

Table 5.3: Power = 85%, $\alpha = 0.10$ and 2-sided Fisher exact test, $d =$ at least 2-fold increase (post / pre ≥ 2) of T-cell recognizing neoantigens in non-vaccine group.

	S1	S2	S3	S4	S5	S6	S7	S8
Vaccine/GM-CSF/Lenalidomide	31%	46%	57%	66%	73%	78%	84%	89%
Non-Vaccine Group ¹	$d = 0/15$ (0%)	$d=1/15$ (7%)	$d=2/15$ (13%)	$d=3/15$ (20%)	$d=4/15$ (27%)	$d=5/15$ (33%)	$d=6/15$ (40%)	$d=7/15$ (47%)

¹Includes patents randomized to both the Lenalidomide/GM-CSF arm and the Lenalidomide Alone arm.

5.4. Analysis of the Secondary Endpoints

Additional exploratory analyses will be conducted in a similar fashion to examine a number of secondary immunologic endpoints, including: 1) clonality assessment and TCR identification of T cells, 2) T cell cytokine expression 3) immune landscape analysis 4) genomic analysis of tumor immunoregulatory pathways 5) patterns of immunoregulatory microRNA expression. In each case, log transformations will be considered to induce normality, and if still non-normal then nonparametric tests will be used. Profiles of these secondary immunologic endpoints will be described at each time point for each group using summary statistics and compared between the vaccine and non-vaccine groups using mixed models for repeated measures data. The correlation between the immune environment measures and the myeloma specific T-cell response will be assessed and reported using Spearman rank correlation at each time point.

Correlation between immune parameters and clinical outcome as measured by achievement and maintenance of CR at 1 year and PFS will be similarly determined. After the PFS and CR data is analyzed in the parent trial, they will be available for analysis in this ancillary study.

ANCILLARY STUDY REFERENCES

1. Bolli, N., et al., *Heterogeneity of genomic evolution and mutational profiles in multiple myeloma*. Nat Commun, 2014. **5**: p. 299
2. Melchor, L., et al., *Single-cell genetic analysis reveals the composition of initiating clones and phylogenetic patterns of branching and parallel evolution in myeloma*. Leukemia, 2014. **28**(8): p. 1705-15.
3. Uttenthal, B., et al., *Wilms' Tumour 1 (WT1) peptide vaccination in patients with acute myeloid leukaemia induces short-lived WT1-specific immune responses*. Br J Haematol, 2014. **164**(3): p. 366-75.
4. Kuball, J., et al., *Pitfalls of vaccinations with WT1-, Proteinase3- and MUC1-derived peptides in combination with MontanideISA51 and CpG7909*. Cancer Immunol Immunother, 2011. **60**(2): p. 161-71.
5. Gong, J., et al., *Activation of antitumor cytotoxic T lymphocytes by fusions of human dendritic cells and breast carcinoma cells*. Proc Natl Acad Sci U S A, 2000. **97**(6): p. 2715-8.
6. Avigan, D., *Dendritic cell-tumor fusion vaccines for renal cell carcinoma*. Clin Cancer Res, 2004. **10**(18 Pt 2): p. 6347S-52S.
7. Rosenblatt J et al. Individualized vaccination of AML patients in remission is associated with induction of antileukemia immunity and prolonged remissions. Sci Transl Med. 2016 Dec
8. Rosenblatt, J., et al., *Vaccination with dendritic cell/tumor fusion cells results in cellular and humoral antitumor immune responses in patients with multiple myeloma*. Blood. **117**(2): p. 393-402. 19
9. Rosenblatt, J., et al., *Vaccination with dendritic cell/tumor fusions following autologous stem cell transplant induces immunologic and clinical responses in multiple myeloma patients*. Clin Cancer Res, 2013. **19**(13): p. 3640-8.

APPENDIX L

REFERENCES

-
- ¹ Ratta, M. *et al.* Dendritic cells are functionally defective in multiple myeloma: the role of interleukin-6. *Blood* 100, 230-237 (2002).
- ² Sharma, A., Khan, R., Joshi, S., Kumar, L. & Sharma, M. Dysregulation in T helper 1/T helper 2 cytokine ratios in patients with multiple myeloma. *Leuk Lymphoma* 51, 920-927, doi:10.3109/10428191003699563 (2010).
- ³ Joshua, D. E., Brown, R. D., Ho, P. J. & Gibson, J. Regulatory T cells and multiple myeloma. *Clin Lymphoma Myeloma* 8, 283-286, doi:10.3816/CLM.2008.n.039 (2008).
- ⁴ Beyer, M. *et al.* In vivo peripheral expansion of naive CD4+CD25high FoxP3+ regulatory T cells in patients with multiple myeloma. *Blood* 107, 3940-3949, doi:10.1182/blood-2005-09-3671 (2006).
- ⁵ Prabhala, R. H. *et al.* Elevated IL-17 produced by TH17 cells promotes myeloma cell growth and inhibits immune function in multiple myeloma. *Blood* 115, 5385-5392, doi:10.1182/blood-2009-10-246660 (2010).
- ⁶ Nielsen, H. *et al.* Immune dysfunction in multiple myeloma. Reduced natural killer cell activity and increased levels of soluble interleukin-2 receptors. *APMIS* 99, 340-346 (1991).
- ⁷ Avigan, D. *et al.* Fusion cell vaccination of patients with metastatic breast and renal cancer induces immunological and clinical responses. *Clin Cancer Res* 10, 4699-4708 (2004).
- ⁸ Gong, J. *et al.* Activation of antitumor cytotoxic T lymphocytes by fusions of human dendritic cells and breast carcinoma cells. *Proc Natl Acad Sci U S A* 97, 2715-2718. (2000).
- ⁹ Gong, J., Chen, D., Kashiwaba, M. & Kufe, D. Induction of antitumor activity by immunization with fusions of dendritic and carcinoma cells. *Nat Med* 3, 558-561. (1997).
- ¹⁰ Vasir, B. *et al.* Fusion of dendritic cells with multiple myeloma cells results in maturation and enhanced antigen presentation. *Br J Haematol* 129, 687-700 (2005).
- ¹¹ Gong, J. *et al.* Reversal of tolerance to human MUC1 antigen in MUC1 transgenic mice immunized with fusions of dendritic and carcinoma cells. *Proc Natl Acad Sci U S A* 95, 6279-6283. (1998).
- ¹² Gong, J. *et al.* Immunization against murine multiple myeloma with fusions of dendritic and plasmacytoma cells is potentiated by interleukin 12. *Blood* 99, 2512-2517. (2002).
- ¹³ Rosenblatt, J. *et al.* Vaccination with dendritic cell/tumor fusion cells results in cellular and humoral antitumor immune responses in patients with multiple myeloma. *Blood* 117, 393-402, doi:10.1182/blood-2010-04-277137 (2011).

-
- ¹⁴ Rosenblatt, J. *et al.* Vaccination with dendritic cell/tumor fusions following autologous stem cell transplant induces immunologic and clinical responses in multiple myeloma patients. *Clin Cancer Res* 19, 3640-3648, doi:10.1158/1078-0432.CCR-13-0282 (2013).
- ¹⁵ McCarthy, P. L. *et al.* Lenalidomide after stem-cell transplantation for multiple myeloma. *N Engl J Med* 366, 1770-1781, doi:10.1056/NEJMoa1114083 (2012).
- ¹⁶ Luptakova, K. *et al.* Lenalidomide enhances anti-myeloma cellular immunity. *Cancer Immunol Immunother* 62, 39-49, doi:10.1007/s00262-012-1308-3 (2013).
- ¹⁷ Davies, F.E. *et al.* Thalidomide and immunomodulatory derivative augment natural killer cell cytotoxicity in multiple myeloma. *Blood*, 210-216 (2001)
- ¹⁸ WAI, L.E. *et al.* Distinct roles for the NK cell-activating receptors in mediating interactions with dendritic cells and tumor cells. *J Immunol*, 222-229, doi: 10.4049/jimmunol.1002597 (2010).
- ¹⁹ BGM Durie *et al.* [International uniform response criteria for multiple myeloma.](#) *Leukemia* (2006) 1-7.